

In the multivariate Cox analysis, the covariates treatment strategy (PDS versus IDS) HR = 0.57 CI 95% [0.44–0.74],  $p < 0.0001$ , and residual tumor after surgery HR = 1.78 CI 95% [1.25–2.53],  $p < 0.0001$  remain significant as a PFS prognostic factors. The OS prognostic factors was the covariates treatment strategy (PDS versus IDS) ( $p < 0.002$ ), residual tumor after surgery ( $p < 0.0001$ ), age at diagnosis ( $p < 0.02$ ) and BRCA mutation ( $p < 0.02$ ).

**Conclusion** Our data of real-world are in line with those reported in clinical trial for patient with advanced ovarian cancer in 1st line setting with surgical treatment.

### 2022-RA-704-ESGO TREATMENT PATTERNS AND TIME TO NEXT TREATMENT AMONG PATIENTS WITH OC IN A REAL-LIFE SETTING IN FINLAND: THE OCRWE-FINLAND STUDY

<sup>1</sup>Heini Rauhamaa, <sup>1</sup>Fredrik Herse, <sup>1</sup>Otti Isomeri, <sup>2</sup>Juhana Idänpään-Heikkilä, <sup>2</sup>Sari Käkälä, <sup>3</sup>Sakari Hietanen, <sup>4</sup>Mikko Loukovaara, <sup>5</sup>Annika Auranen. <sup>1</sup>Nordic Healthcare Group, Helsinki, Finland; <sup>2</sup>GlaxoSmithKline, Espoo, Finland; <sup>3</sup>Turku University Hospital, Turku, Finland; <sup>4</sup>Helsinki University Hospital, Helsinki, Finland; <sup>5</sup>Tays Cancer Centre, Tampere University Hospital and Tampere University, Tampere, Finland

10.1136/ijgc-2022-ESGO.549

**Introduction/Background** Ovarian cancer (OC) is a disease characterized by a dynamic treatment landscape in the real-life setting. The OCRWE-Finland study aims at describing the real-life burden of patients with OC, including treatment patterns, time to next treatment, disease characteristics and progression, survival, and healthcare resource utilization. This abstract reports on the observed treatment patterns.

**Methodology** OCRWE-Finland is a multicentre, retrospective, noninterventional study collecting hospital medical records from university hospitals in Helsinki, Turku, and Tampere. Patients with ovarian, fallopian tube, or primary peritoneal cancer who were newly diagnosed as part of routine clinical care and received all OC treatments in these hospitals from 2014–2019 were included. Registry data were collected and combined by Findata (authorization holder), operating under the performance guidance of the Finnish Ministry of Social Affairs and Health.

**Abstract 2022-RA-704-ESGO Table 1** Baseline characteristic and first-line treatment patterns among patients with HGSOc

	Patients with HGSOc N=621
FIGO stage at diagnosis	
Stage IV	20%
Stage III	55%
Stage II	5%
Stage I	9%
No data	11%
First-line treatment	
Primary debulking surgery	59%
Neoadjuvant chemotherapy and interval debulking surgery	21%
Chemotherapy only	12%
Residual tumour status after primary debulking surgery	
Optimal debulking	37%
Suboptimal debulking	44%
Residual <1cm	27%
Residual ≥1cm	17%
Unknown residual tumour status	19%

FIGO, International Federation of Gynaecology and Obstetrics; HGSOc, high-grade serous ovarian cancer.

**Results** In total, 1711 patients with OC (mean age=65.9 y, StDev=13.4 y) and 621 patients with high-grade serous OC

(HGSOc) (mean age=68.0 y, StDev=10.1 y) were identified. Disease origin was ovaries in 75% of patients and peritoneum in 19%. Baseline characteristics and first-line treatment (TL1) patterns among patients with HGSOc can be found in table 1. During the observation period, 57% of patients received TL2, with 48% of these moving to TL3. The probability of undergoing TL2 was higher among stage III/IV patients and those with residual disease. In TL2, the most common treatment was platinum-based chemotherapy (32%); 26% received 'other chemotherapy', 33% of patients did not receive TL2 during this period but were still alive, and 9% died before initiating TL2.

**Conclusion** This study documents real-life treatment patterns across lines of treatment among patients with OC and HGSOc during the first years of disease from the 3 biggest university hospitals in Finland. These results can provide useful baseline information about the rapidly evolving treatment landscape in OC in recent years.

### 2022-RA-706-ESGO PROGNOSTIC IMPACT OF VAGUS NERVE ACTIVITY AT INITIAL MANAGEMENT OF OVARIAN CANCER

<sup>1,2</sup>Francois Cherifi, <sup>2</sup>Sophie Lefevre Arbogast, <sup>3</sup>Jonaz Font, <sup>2</sup>Justine Lequesne, <sup>4</sup>Stephanie Becourt, <sup>4</sup>Cyril Abdeddaim, <sup>5</sup>Yori Gidron, <sup>2</sup>Florence Joly. <sup>1</sup>Medical Oncology department, Centre François Baclesse, Caen, France; <sup>2</sup>Clinical Research department, Centre François Baclesse, Caen, France; <sup>3</sup>Cardiology department, CHU de Caen, Caen, France; <sup>4</sup>Medical Oncology department, Centre Oscar Lambret, Lille, France; <sup>5</sup>Nursing, Haifa University, Haifa, Israel

10.1136/ijgc-2022-ESGO.550

**Introduction/Background** Finding new modifiable prognostic markers is important in ovarian cancer (OC). The autonomic nervous system plays an important role in cancer initiation and progression. Low parasympathetic nervous system activity is associated with inflammation, oxidative stress and sympathetic activation. Low vagal nerve activity, measured by low heart rate variability (HRV) predicts poor cancer prognosis. Our study examined the prognostic value of HRV in OC.

**Methodology** We conducted a bicentric retrospective study. We analyzed patients diagnosed with serous OC stage FIGO ≥ IIB, between January 2015 to August 2019, with an electrocardiogram (ECG) available around diagnosis. We used the time domain HRV parameter of the standard deviation of all normal-to-normal beat interval (SDNN) in 10 seconds ECG. Optimal SDNN cut-off was found using the Youden index criteria of time-dependent ROC curve. We carried out multivariable analysis including HRV and well-known OC prognostic factors.

**Results** We included 202 patients with a median age of 65 years, 93% had stage FIGO IIIC/IV, 56% had complete surgical resection. Median overall survival (OS) was 38.6 months [95%CI:34.4–47.4]. The median SDNN was 11.1 ms (min=1.93; max=74.5), with an optimal cut off of 10 ms to predict OS. Median OS was significantly shorter for patients with low HRV compared to high HRV (26.4 vs 45.1 months;  $p < 0.001$ ). In a multivariable analysis, HRV remained a strong independent prognostic factor with a two-fold higher risk of death among patients with low SDNN compared to those with high SDNN (HR=2.09 [1.40–3.124],  $p < 0.001$ ); other associated factors with higher risk of death were ECOG > 0, high CA125 level and incomplete resection.