

Inguino-abdominal combined approach for laterally extended pelvic resection: a step by step procedure

Giuseppe Vizzielli,¹ Emanuele Perrone,^{1,2} Alessandro Gioè,^{1,2} Giovanni Tinelli,³ Giovanni Scambia^{1,2}

► This video is too large to play in the PDF, please visit the full text version online at: <https://ijgc.bmj.com/content/29/2/443>

¹UOC di Ginecologia Oncologica, Department of Women's and Children's Health, Fondazione Policlinico Universitario A Gemelli IRCCS, Roma, Italia
²Istituto di Ostetricia e Ginecologia, Università Cattolica del Sacro Cuore, Roma, Italia
³Division of Vascular Surgery, Fondazione Policlinico Universitario A Gemelli IRCCS, Roma, Italia

Correspondence to

Emanuele Perrone, Department of Women's and Children's Health, Fondazione Policlinico Universitario A Gemelli IRCCS, Roma 00168, Italy; ema.perrone88@gmail.com

Accepted 7 December 2018

SUMMARY

This video article demonstrates an inguino-abdominal combined approach for laterally extended pelvic resection, a major surgical procedure for locally advanced primary or recurrent gynecological cancer infiltrating the pelvic sidewall, for which palliative therapy is the only alternative.¹


After local institutional review board approval (protocol No CICOG 02/03/62), we made a step by step surgical video of an inguino-abdominal combined approach for laterally extended pelvic resection [Video 1](#), defined as an en bloc resection of a pelvic tumor with pelvic sidewall structures, including the iliopsoas and/or obturator internus muscles.^{2,3}

The patient, a 48-year-old woman, diagnosed with single pelvic platinum resistant recurrence after five lines of chemotherapy for serous ovarian cancer G3, International Federation of Gynecology and Obstetrics (FIGO) stage IIIC, BRCA wild type. The preoperative positron emission tomography/computed tomography scan detected uptake on the right side at the level of the external iliac region and obturator fossa: the tumor surrounded the right external iliac vessels by more than 50% of their circumferences, with possible involvement of the vascular wall and venous vascular compression (Tinelli's score=4).⁴ The tumor extended towards the

obturator fossa, with possible involvement of the inguinal canal. Due to an uncertain pathological response, the size of the recurrence, and its close contiguity with the ureter and bowel, we decided to avoid radiation therapy as it could result in a ureteral or intestinal fistula. We performed a laterally extended pelvic resection, as shown step by step in the video.

The procedure was conducted until complete removal of recurrence (R0). Estimated blood loss was 1000 mL and total operative time was 240 min. The patient was discharged after 15 days; we reported a urinary infection, a likely postoperative complication. The pathology report described a lymphodal relapse of ovarian cancer (diameter=6 cm) with infiltration of surrounding tissue and in the sano margins. Six months after surgery, the patient is alive without evidence of relapse.

The borders of pelvic surgical anatomy are continually extending, requiring surgeons to use a personalized approach and to continually update their anatomic knowledge. In this context, laterally extended pelvic resection could be a feasible surgical procedure, representing a salvage treatment in recurrent or persistent primary gynecological malignancies infiltrating the pelvic sidewall, when other approaches have failed. However, additional clinical trials are needed to confirm these results.³



Video 1



© IGCS and ESGO 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Vizzielli G, Perrone E, Gioè A, *et al.* *Int J Gynecol Cancer* 2019;**29**:443–444.

Video Article

Contributors Conception: PE, VG; Design: PE, VG; Supervision: SG; Materials and data collection: PE, GA; Analysis/interpretation: VG, TG; Literature review: PE, VG; Writer: PE; Critical review: PE, VG, SG.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Ethics approval The study was approved by the local institutional review board.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Vizzielli G, Chiantera V, Tinelli G, *et al.* Out-of-the-box pelvic surgery including iliopsoas resection for recurrent gynecological malignancies: does that make sense? a single-institution case-series. *Eur J Surg Oncol* 2017;43:710–6.
2. Cibula D. Principles of gynecologic oncology surgery. In: *Pelvic exenteration for gynecological cancers*. Philadelphia: Elsevier, 2018.
3. Vizzielli G, Naik R, Dostalek L, *et al.* Laterally extended pelvic resection for gynaecological malignancies: a multicentric experience with out-of-the-box surgery. *Ann Surg Oncol* 2018. [Epub ahead of print 11 Dec 2018].
4. Tinelli G, Cappuccio S, Parente E, *et al.* Resectability and vascular management of retroperitoneal gynecological malignancies: a large single-institution case-series. *Anticancer Res* 2017;37:6899–906.