

Legends

Figure 2 (supplement). Actuarial recurrence free survival of 161 patients with early cervical cancer and IR factors treated by a radical hysterectomy without adjuvant radiotherapy.

Figure 3 (supplement). Actuarial disease specific survival of 161 patients with early cervical cancer and IR factors treated by a radical hysterectomy without adjuvant radiotherapy.

Table 2 (supplement): Recurrence pattern and frequency for Sedlis risk groups 1,2,3,4 (see Patient and Methods section for legends)

Table 5 (supplement). Recurrence pattern for patients with stage IB1 and IB2.

Supplemental figures




Figure 2 (supplement).




Figure 3 (supplement).

	Sedlis 1 (n=25)	Sedlis 2 (n=62)	Sedlis 3 (n=5)	Sedlis 4 (n=69)
• Loco-regional	2 (8%)	4 (6.5%)	0	3 (4.3%)
• Distant/Combined	3 (12%)	6 (9.7%)	1 (20%)	6 (8.6%)
Total	5 (20%)	10 (16.2%)	1 (20%)	9 (12.9%)

Table 2 (supplement).

Recurrence localization	IB1 (n=87)	IB2 (n=74)
<ul style="list-style-type: none"> • Loco-regional 	4 (4.6%)	5 (6.8%)
<ul style="list-style-type: none"> • Distant/Combined 	10 (11.5%)	6 (8.1%)
Total	14 (16.1%)	11 (14.9%)

Table 5 (supplement).