

Supplementary Table 1. Comparison of the clinicopathological features according to the adjuvant treatment groups.

Stage IIIA	RT	CT	CRT	p
	n= 77 n (%)	n= 65 n (%)	n= 157 n (%)	
Age (years)				
<60	38 (49.3)	29 (44.6)	92 (58.6)	0.121
≥60	39 (50.7)	36 (55.4)	65 (41.4)	
Grade				
1	12 (15.6)	8 (12.3)	24 (15.3)	0.823
2 or 3	65 (84.4)	57 (87.7)	133 (84.7)	
MMI				
<50%	13 (16.9)	22 (33.9)	64 (40.8)	0.094
≥50%	40 (51.9)	42 (64.6)	80 (50.9)	
Not known	24 (31.2)	1 (1.5)	13 (8.3)	
LVSI				
Negative	14 (18.2)	25 (38.5)	63 (40.1)	0.086
Positive	39 (50.6)	38 (58.5)	81 (51.6)	
Not known	24 (31.2)	2 (3.0)	13 (8.3)	
Lymphadenectomy				
None	33 (42.8)	6 (9.2)	46 (29.3)	<0.0001
Pelvic	17 (22.1)	6 (9.2)	43 (27.4)	
Pelvic+Para-aortic	27 (35.1)	53 (81.5)	68 (43.3)	
	Median (range)	Median (range)	Median (range)	
LNR	0.039 (0.002-1.0)	0.028 (0.007-1.0)	0.064 (0.003-1.0)	0.187
Stage IIIC	RT	CT	CRT	
	n= 99 n (%)	n= 170 n (%)	n= 381 n (%)	
Age (years)				
<60	51 (51.5)	89 (52.3)	194 (50.9)	0.281
≥60	48 (48.5)	81 (47.7)	187 (49.1)	
Grade				
1	27 (27.3)	33 (19.4)	53 (13.9)	0.005
2 or 3	72 (72.7)	137 (80.6)	328 (86.1)	
MMI				
<50%	29 (29.3)	38 (22.4)	83 (21.8)	0.181
≥50%	65 (65.6)	132 (77.6)	294 (77.2)	
Not known	5 (5.1)	0	4 (1.0)	
LVSI				
Negative	25 (25.2)	30 (17.6)	43 (11.3)	<0.0001
Positive	67 (67.7)	135 (79.4)	324 (85.0)	
Not known	7 (7.1)	5 (2.9)	14 (3.7)	
Lymphadenectomy				
Pelvic	26 (26.3)	17 (10)	101 (26.6)	<0.0001
Pelvic+Para-aortic	73 (73.7)	153 (90)	280 (73.4)	
Substage				
IIIC1	63 (63.6)	94 (55.3)	211 (55.5)	0.171
IIIC2	36 (36.4)	76 (44.7)	170 (44.5)	
LNR	Median (range)	Median (range)	Median (range)	
	0.013 (0.003-0.6)	0.034 (0.006-1.0)	0.15 (0.006-1.0)	0.02

Abbreviations: CRT: Chemoradiation. CT: Chemotherapy. LNR: Lymph node ratio. LVSI: Lymphovascular space invasion. MMI: Myometrial invasion. RT: Radiotherapy.