(67.7%) underwent systematic LND. Patients in the SLN cohort had a median age of 66 years, compared to 68 years in the LND cohort ($p>0.05$). Median follow-up time was 23 months (range, 1–96) in the SLN cohort and 66 months (range, 4–265) in the LND cohort ($p<0.001$). In patients with stage I/II disease ($n=160$, 60.1%), the 2-year OS was 96.6% (SE ± 3.4) in the SLN cohort and 89.6% (SE ± 2.2) in the LND cohort ($p=0.8$). In patients with stage III disease ($n=77$), the 2-year OS was 73.6% (SE ± 10.2) in the SLN cohort and 77.3% (SE ± 5.8) in the LND cohort ($p=0.8$).

Conclusions SLN mapping alone and systematic pelvic and paraaortic nodal dissection (LND) led to similar survival outcomes in patients with stage I–III USC. In our practice, the SLN mapping algorithm has replaced systematic LND as the primary staging modality in the setting of apparent uterine-confined endometrial serous cancer.

IGCS19-0156

41 SENTINEL NODE MAPPING VS. LYMPHADENECTOMY IN ENDOMETRIAL CANCER: A SYSTEMATIC REVIEW AND META-ANALYSIS

G Bogani, A Ditto, F Murgia, V Chiappa, U Leone, F Raspagliesi*. *Fondazione IRCCS Istituto Nazionale dei Tumori di Milano-, Gynecologic Oncology, Milano, Italy*

10.1136/ijgc-2019-IGCS.41

Objectives Sentinel node mapping is increasingly being utilized for endometrial cancer staging. However, only limited evidence supporting the adoption of sentinel node mapping instead of conventional lymphadenectomy is still available. Here, we aimed to review the current evidence comparing sentinel node mapping and lymphadenectomy in endometrial cancer staging.

Methods This systematic review was registered in the International Prospective Register of Systematic Reviews. Der-Simonian and Laird random-effects models were used to pool log transformed event rates and estimated 95%CI for dichotomous outcomes between the two interventions for each study and we pooled the effect size using the same models.

Results Overall, 3,536 patients were included: 1,249 (35.3%) and 2,287 (64.7%), undergoing sentinel node mapping and lymphadenectomy, respectively. Pooled data suggested that positive pelvic nodes were detected in 184 out of 1,249 (14.7%) patients having sentinel node mapping and 228 out of 2,287 (9.9%) patients having lymphadenectomy (OR:2.03; (95% CI:1.30 to 3.18); $p=0.002$). No difference in detection of positive nodes located in the paraaortic was observed (OR:0.93 (95%CI:0.39 to 2.18); $p=0.86$). Overall recurrence rate was 4.3% and 7.3% after sentinel node mapping and lymphadenectomy, respectively (OR:0.90 (95%CI:0.58 to 1.38); $p=0.63$). Similarly, nodal recurrences were statistically similar between groups (1.2% vs. 1.7%; OR: 1.51 (95%CI:0.70 to 3.29); $p=0.29$).

Conclusions In conclusion, our meta-analysis underlines that sentinel node mapping is non-inferior to standard lymphadenectomy in term of detection of paraaortic nodal involvement and recurrence rates (any site and nodal recurrence); while, focusing on the ability to detect positive pelvic nodes,

sentinel node mapping could be consider superior to lymphadenectomy.

Regional Plenary

Latin America Regional Plenary

IGCS19-0394

42 SOCIAL AND ECONOMIC IMPACT OF CERVICAL CANCER IN BRAZIL: ANALYSIS OF EVA/EVITA COHORT STUDY (LACOG 0215)

¹A Nogueira-Rodrigues, ²A Melo*, ³A Calabrich, ⁴E Cronenberger, ⁵K Torres, ⁶F Damian, ⁷R Cossetti, ⁸C Azevedo, ⁹A Fonseca, ¹⁰N Yeni, ¹¹J Nunes, ¹²A Lopes, ¹³F Thomé, ¹⁴R Leal, ¹⁵G Borges, ¹⁶PR Nunes Filho, ¹⁶F Zaffaroni, ¹⁶L Hertzog, ¹⁶R Gomes, ¹⁶G Werutsky, ¹⁷F Maluf. ¹Universidade Federal de Minas Gerais, Medical Oncology, Belo Horizonte, Brazil; ²Brazilian National Cancer Institute – INCA, Medical Oncology, Rio de Janeiro, Brazil; ³Clínica Assistência Multidisciplinar em Oncologia, Medical Oncology, Salvador, Brazil; ⁴Centro Regional Integrado de Oncologia, Medical Oncology, Fortaleza, Brazil; ⁵Fundação Centro de Controle de Oncologia do Estado do Amazonas, Medical Oncology, Manaus, Brazil; ⁶Centro de Pesquisa em Oncologia, Medical Oncology, Porto Alegre, Brazil; ⁷Hospital Aldenora Bello, Medical Oncology, São Luís, Brazil; ⁸Instituto de Medicina Integral Prof Fernando Figueira, Medical Oncology, Recife, Brazil; ⁹Hospital Geral de Roraima, Medical Oncology, Boa Vista, Brazil; ¹⁰Centro de Pesquisas Oncológicas, Medical Oncology, Florianópolis, Brazil; ¹¹Hospital Erasto Gaertner, Medical Oncology, Curitiba, Brazil; ¹²Instituto Brasileiro de Controle do Câncer, Medical Oncology, São Paulo, Brazil; ¹³Hospital São Vicente de Paulo, Medical Oncology, Passo Fundo, Brazil; ¹⁴Hospital Universitário Walter Cantídio, Medical Oncology, Fortaleza, Brazil; ¹⁵Clínica Neoplasias Litoral, Medical Oncology, Itajaí, Brazil; ¹⁶Latin American Cooperative Oncology Group, Clinical Research, Porto Alegre, Brazil; ¹⁷Hospital Beneficência Portuguesa de São Paulo, Medical Oncology, São Paulo, Brazil

10.1136/ijgc-2019-IGCS.42

Objectives Compare marital status, household income and labor activity at diagnosis and at 1-year follow-up among cervical cancer (CC) patients in Brazil.

Methods LACOG 0215 EVITA is a prospective observational study, within 16 Brazilian sites, including 18 years-old, newly diagnosed, stage I to IV invasive CC patients. We present data collected at baseline and at 1-year follow-up regarding demography, histology, stage, marital status, household income and labor activity. Variables were compared using Chi-square test.

Results 593 patients were included in this analysis. 75 (12.6%) were diagnosed in stage I, 452 (76.2%) II–III and 66 (11.2%) IV. Mean age at diagnosis was 57.0, 56.4 and 50.5 years respectively. Squamous cell carcinoma was the most frequent histology. 34.2% of stage I, 30.6% of stage II–III and 24.6% of stage IV patients had 9 or more years of education. About 95% patients had a previous pregnancy history. At 1-year follow-up there was a decrease in the proportion of married patients in stage II–III patients only (53.3% to 47.5%; p -value <0.001). There was a reduction in the proportion of women engaged in labor activity in stage II–III (34.1% to 16.1%; p -value, <0.001) but not in stage I and IV. No difference was found in household income 1 year after diagnosis.

Conclusions CC may impact patients personal and work life long after diagnosis, even in those diagnosed with curable disease. In this cohort, stage II–III CC face more risk of divorce and job losses. Social support following diagnosis and treatment is essential for CC patients.