Methods Design: Cohort-study set within recruitment to the GCaPPS-trial (ISRCTN73338115).

AJ women/men >18-years, from the North-London AJ-population were recruited through self-referral. AJ-women/men underwent pre-test counselling for *BRCA*-testing through recruitment clinics (clusters). Consenting individuals provided blood-sample for *BRCA*-testing. Socio-demographic/family-history/knowledge/psychological well-being data along-with benefits/risks/cultural-influences (18-item-questionnaire measuring 'attitude') were collected.

4-item likert-scales analysed initial 'interest' and 'intention-to-test' pre-counselling.

Uni-&-multivariable logistic-regression-models evaluated factors affecting uptake/interest/intention-to undergo *BRCA*-testing. Statistical inference was based on cluster robust standard-errors and joint Wald-tests for significance. Item-Response-Theory and graded-response-models modelled responses to 18-item questionnaire.

Main Outcome Measures: Interest, intention, uptake, attitude towards *BRCA*-testing

Results 935 AJ women (67%) and men (33%) underwent pre-test genetic-counselling (mean-age=53.8(S.D=15.02) years). Pre-counselling 96% expressed interest but 60% had clear intention-to undergo *BRCA*-testing. Subsequently 88% opted for *BRCA*-testing. *BRCA*-related knowledge ($p=0.013$) and degree-level education($p=0.01$) were positively and negatively (respectively) associated with intention-to-test. Being married/cohabiting had four-fold higher-odds for *BRCA*-testing uptake ($p=0.009$). Perceived benefits were associated with higher pre-counselling odds for interest and intention-to undergo *BRCA*-testing. Reduced uncertainty/reassurance were the most important factors contributing to decision-making. Increased importance/concern towards risks/limitations (confidentiality/insurance/emotional-impact/inability to prevent cancer/marriage-ability/ethnic-focus/stigmatization) were significantly associated with lower-odds of uptake-of *BRCA*-testing, and discriminated between acceptors and decliners. Having children had stronger ($p=0.005$) while male-gender/degree-level-education ($p=0.001$) had weaker, attitudes towards *BRCA*-testing.

Conclusions *BRCA* testing in the AJ population has high acceptability. Pre-test counselling increases awareness of disadvantages/limitations of *BRCA*-testing, influencing the final cost-benefit perception and decision-making on undergoing testing.

IGCS19-0493

82

IDENTIFICATION OF NEOANTIGENS FROM NON-PRIMARY TUMOR TISSUE IN PATIENTS WITH RECURRENT OVARIAN CANCER BY SEQUENCING AND SUBSEQUENT HLA LIGAND MASS SPECTROMETRY

¹R O'Cearbhaill*, ²M Klatt, ²SS Mun, ²T Korontsvit, ³N Socci, ²S Guzman, ⁴M Mattar, ⁵O Zivanovic, ²T Dao. ¹Memorial Sloan Kettering Cancer Center and Weill Cornell Medical College, Medicine, New York, USA; ²Sloan Kettering Institute, Molecular Pharmacology, New York, USA; ³Memorial Sloan Kettering Cancer Center, Bioinformatics Core, New York, USA; ⁴Sloan Kettering Institute, Clinical Research, New York, USA; ⁵Memorial Sloan Kettering Cancer Center and Weill Cornell Medical College, Gynecologic Surgical Oncology, New York, USA

10.1136/ijgc-2019-IGCS.82

Objectives Adoptive transfer of autologous Tcells specifically targeting neoepitopes derived from tumor mutations has

achieved responses in melanoma and other cancers. We sought to identify ovarian cancer neoepitopes.

Methods We collected tumor, blood and ascites specimens from ovarian cancer patients. We combined mass spectrometry analysis of MHC class1 peptidomes with parallel sequencing of whole exome and RNA of cancer cells, to identify neoepitopes. Missense mutations were identified by whole exome sequencing and epitopes confirmed by mass spectrometry from ascites. We selected peptides predicted to have high binding affinity to HLA-A*02:01, for *in vitro* Tcell priming. We first validated the immunogenicity of these peptides for priming Tcells with synthesized peptides in healthy HLA-A*02:01+ donors.

Results 32 missense mutations were identified in patient's ascites. 7 epitopes were confirmed by mass spectroscopy; 5 were predicted to have high binding affinity to HLA-A*02:01. 3/5 epitopes induced peptide-specific Tcell responses not cross-reactive with native sequences. However, these 3 peptides failed to induce autologous peptide-specific Tcell response in TILs to patient's ascites, nor to autologous tumor cells, suggesting immunosuppression. Interestingly, when autologous tumor cells were pre-treated with IFN-gamma, Tcell responses against tumor cells were observed.

Conclusions We demonstrated effective T cell responses against tumor neoantigens, is not only dependent on stimulation of Tcells, but also on efficient epitope processing and presentation on cell surface. While efforts have been focused on activation of Tcell responses, such as immune checkpoint blockade, our study suggests that strategies to modulate tumor cells to efficiently present neoepitopes should be considered for successful immunotherapy against neoantigens.

IGCS19-0587

83

IMPACT OF INTRODUCING EXTENSIVE SURGICAL PROCEDURES TO CYTOREDUCTIVE SURGERY (CRS) IN ADVANCED EPITHELIAL OVARIAN CANCER (EOC) ON RATE OF COMPLETE DEBULKING (CD) AND SURVIVAL

¹M Seoud*, ¹A El Housseini, ¹I Jaafar, ¹K Hamed, ¹R Abdallah, ¹A Khalil, ²F Jamali. ¹American University of Beirut Medical Center, Obstetrics and Gynecology, Beirut, Lebanon; ²American University of Beirut Medical Center, General Surgery, Beirut, Lebanon

10.1136/ijgc-2019-IGCS.83

Objectives To asses the impact of extensive procedures (Diaphragmatic resection, large and small bowel resection, complete peritonectomy, complete omentectomy, paraaortic lymph node dissection, splenectomy, resection of all visible disease) in CRS on the rate of CD, progression free (PFS), and overall survival (OS) in patients with advanced EOC.

Methods Patients undergoing CRS between 2002–2018 were divided into group#1(2002–2008), and group#2 (2009–2018). Demographic and operative information were retrospectively reviewed.

Results 204 patient with EOC (IIIC-IV) underwent CRS between 2002–2018. The number of CRS as well as performing extensive procedures (figure1A) has significantly increased over the whole period (72 in group1 vs 132 in group2: p -value<0.001), and this coincided with a significant increase in the CD rate from 50% to 80% (p -value<0.001) (figure 1A). In stage IIIC, median PFS was significantly higher in group#2 versus group#1 (22 months, 95%CI 16.80–27.19 and 23