Laparotomy approach to sentinel lymph node detection in ovarian cancer using a near-infrared fluorescent system camera with indocyanine green dye

Luigi Carlo Turco ^{1,2}, Virginia Vargiu,³ Camilla Nero,³ Anna Fagotti,^{3,4} Giovanni Scambia,^{4,5} Francesco Cosentino²

¹Gynecology and Breast care unit, Mater Olbia spa, Olbia, Italy

²Gynecologic Oncology, Gemelli Molise spa, Università Cattolica del Sacro Cuore, Campobasso, Italy

³Department of Women's and Children's Health, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

⁴Università Cattolica del Sacro Cuore Facoltà di Medicina e Chirurgia, Roma, Lazio, Italy

⁵Department of Women's and Children's Health, Fondazione Policlinico Universitario A. Gemelli IRCCS, Roma, Italy

Correspondence to

Dr Luigi Carlo Turco,
Gynecology and Breast care
unit, Mater Olbia spa, 07026
Olbia, Italy; luigicarlo.turco@
gmail.com

Accepted 22 January 2020

Published Online First

19 February 2020

© IGCS and ESGO 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Turco LC, Vargiu V, Nero C, et al. *Int J Gynecol Cancer* 2020;**30**:712–713.

SUMMARY

Current guidelines for complete surgical staging in early-stage ovarian cancer recommend systematic lumbo-aortic and pelvic lymphadenectomy,¹ despite their controversial therapeutic value with only 14% of patients upstaged.² Sentinel lymph node (SLN) detection with indocyanine green in ovarian cancer is a feasible technique and could provide useful information on nodal status,³ avoiding future lymphadenectomy in the majority of patients. SLN biopsy is not yet the standard approach, and is currently under investigation.³ Laparoscopy is the most tested approach,³ but it is not advisable in voluminous ovarian masses.⁴

The patient in this video is a 70-year-old woman with a 20 cm ovarian mass suspected for cancer, who was enrolled in our prospective trial SELLY (NCT03452982) (Video 1).³ After mobilization of the ovarian mass, 2 mL of 1.25 mg/mL indocyanine green solution (Verdyne 5 mg/mL injection, 25 mg/50 mg, powder for solution for injection, Diagnostic Green GmbH Aschheim-Dornach Germany) was injected with a 20 gauge spinal needle into the perivascular connective tissue of the infundibulo-pelvic and the utero-ovarian ligament of the affected ovary. The ovarian mass was removed after 15 min and subjected to frozen sectioning. After a diagnosis of cancer, the retro-peritoneum was exposed on both sides along the Toldt fascia up to the left renal

vein and inspected using the NIR mode of the Olympus laparoscopic camera (VISERA-ELITE II-Olympus Winter & Ibe GmbH, Hamburg, Germany) with overhead lights switched-off. The first fluorescent lymph node of the lymphatic chain, identified as an inter-cavo-aortic SLN, was removed and examined separately with ultra-staging at final pathology. No pelvic lymph nodes were identified because of diffuse tissue fluorescence. Systematic lymphadenectomy was then performed.

At final pathology, the inter-cavo-aortic SLN was positive for isolated tumor cells, while two right pelvic nodes were metastatic.

In conclusion, a laparotomy approach for SLN detection in voluminous ovarian cancer is feasible. Further information about the lymphatic routes of the ovary and the accuracy of SLN biopsy in ovarian cancer will be forthcoming after completion of the SELLY trial.³

Twitter Anna Fagotti @annafagottimd

Acknowledgements We thank Dr Maria De Ninno, pathology department of Gemelli Molise SpA, for the kind use of the pathology image.

Contributors LCT: responsible surgeon, manuscript preparation, patient recruitment; VV: conception, manuscript and video preparation, video editing; CN: video narration; AF: conception, interpretation; GS: conception, interpretation; FC: surgeon, conception, interpretation.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon request.

ORCID iD

Luigi Carlo Turco <http://orcid.org/0000-0002-3000-178X>

REFERENCES

- Colombo N, Sessa C, du Bois A, et al. ESMO–ESGO ovarian cancer consensus conference working group.

Video 1. Inter-cavo-aortic sentinel lymph node marked by indocyanine green dye.

- ESMO-ESGO consensus conference recommendations on ovarian cancer: pathology and molecular biology, early and advanced stages, borderline tumours and recurrent disease. *Ann Oncol* 2019;30:672–705.
- 2 Kim HS, Ju W, Jee BC, *et al.* Systematic lymphadenectomy for survival in epithelial ovarian cancer: a meta-analysis. *Int J Gynecol Cancer* 2010;20:520–8.
 - 3 Uccella S, Nero C, Vizza E, *et al.* Sentinel-node biopsy in early-stage ovarian cancer: preliminary results of a prospective multicenter study (SELY). *Am J Obstet Gynecol* 2019.
 - 4 Casarin J, Laganà AS, Uccella S, *et al.* Surgical treatment of large adnexal masses: a retrospective analysis of 330 consecutive cases. *Minim Invasiv Ther* 2019;55:1–9.