Time to accept a new old standard of care in cervical cancer

Roni Nitecki ,¹ Alexander Melamed^{2,3}

For numbered affiliations see end of article.

Correspondence to

Dr Alexander Melamed,
Department of Obstetrics and
Gynecology and Herbert Irving
Comprehensive Cancer Center,
Columbia University Vagelos
College of Physicians and
Surgeons, New York NY 10032,
USA; am5195@cumc.columbia.
edu

Received 14 July 2020

Accepted 16 July 2020

Published Online First

12 August 2020

The Laparoscopic Approach to Cervical Cancer (LACC) trial disrupted the long-standing consensus that minimally invasive and open surgery are equally effective approaches to radical hysterectomy for cervical cancer.¹ The study's main finding, namely that radical hysterectomy by a minimally invasive route leads to higher rates of recurrence than laparotomy, has been cited 313 times in less than 2 years, underscoring the vibrant discourse that has ensued. While some commentators continue to favor the use of minimally invasive surgery in clinical practice while additional randomized trials are in progress, national and international guidelines have coalesced around the notion that laparotomy is the evidence-based approach to radical hysterectomy for early-stage cervical cancer.^{2–5} In addition to LACC, these guidelines have cited a growing literature of adequately powered, well-designed observational studies that, overall, have had concordant results.^{6–15}

In this issue of the *International Journal of Gynecological Cancer*, Chiva and colleagues¹⁶ present additional evidence of the deleterious effect of minimally invasive radical hysterectomy, and suggest possible mechanisms underlying this association. The SUCCOR study is a large, retrospective, observational cohort study that evaluates associations between surgical approach and survival outcomes among women with International Federation of Gynecology and Obstetrics (FIGO) 2009 stage 1B1 cervical cancer within 126 institutions in 29 European countries. This study has several important strengths that distinguish it from related observational studies. First, the inclusion of 693 patients provided enough statistical power to detect clinically meaningful differences in survival outcomes. Moreover, because women who received minimally invasive surgery were compared with others who underwent laparotomy during the same time period, the SUCCOR study avoids biases introduced by the use of historic controls, a practice that plagues many observational studies of minimally invasive radical hysterectomy. Chiva and colleagues also utilized well-defined inclusion and exclusion criteria and employed propensity score methods to adjust for variables that may confound the relationship between surgical approach and oncologic outcomes. Finally, their efforts to assess how manipulator use

and vaginal closure may modify the magnitude of the associations between minimally invasive hysterectomy and adverse survival outcomes is an important step in elucidating the factors that may mediate these effects. While some recent well-designed observational studies have not found an association between minimally invasive radical hysterectomy and recurrence or death,^{17–18} the majority of such studies^{6–15} have found evidence of harm. The SUCCOR study delivers another significant blow to a surgical paradigm that is increasingly difficult to justify in routine clinical practice.

While SUCCOR is a high-quality observational study, there are several important limitations. The possibility of residual confounding limits our confidence that the estimated hazard ratios are unbiased measures of the causal effect of minimally invasive radical hysterectomy. However, since patients who undergo minimally invasive surgery tend to have favorable prognostic features compared with those who receive open surgery, residual confounding is likely to lead to an underestimation of the harms of minimally invasive surgery, and therefore unlikely to alter the study's qualitative conclusions. Another important caveat is that while Chiva and colleagues did not find statistically significant differences in outcomes between surgical approaches among women with tumors smaller than 2 cm, the absence of statistical significance should not be interpreted as evidence of the absence of harm in this population. The hazard ratio estimates and associated confidence intervals in this subgroup are not reassuring, and we caution against concluding, based on these data, that a minimally invasive approach is safe for patients with small tumors. Similarly, while the data, which suggest that avoidance of uterine manipulators and vaginal closure before colpotomy might reduce harm from minimally invasive surgery, are provocative, such techniques must be tested in controlled trials before any version of a minimally invasive radical hysterectomy can be considered for routine clinical practice.

The benefits of minimally invasive surgery are well known, yet they cannot justify an increased risk of recurrence or death. Until a phase III randomized trial demonstrates acceptable survival outcomes with minimally invasive surgery, radical hysterectomy for

© IGCS and ESGO 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Nitecki R, Melamed A.
Int J Gynecol Cancer
2020;**30**:1278–1279.

cervical cancer should be performed by laparotomy outside of research studies and unusual clinical circumstances.

Author affiliations

¹Gynecologic Oncology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA

²Department of Obstetrics and Gynecology and Herbert Irving Comprehensive Cancer Center, Columbia University Vagelos College of Physicians and Surgeons, New York, New York, USA

³NewYork-Presbyterian/Columbia University Medical Center, New York, New York, USA

Contributors Dr Nitecki and Dr Melamed coauthored the manuscript and contributed equally to the ideas expressed. Dr Nitecki wrote the first draft of the manuscript, Dr Melamed contributed to revisions, and both authors approved the final draft.

Funding This work was supported by grants from The National Institutes of Health National Cancer Institute (RN: P30 48CA016672 and 5T32 CA101642) and National Center for Advancing Translational Sciences (AM: KL2TR001874).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

ORCID iD

Roni Nitecki <http://orcid.org/0000-0002-6176-9441>

REFERENCES

- Ramirez PT, Frumovitz M, Pareja R, *et al.* Minimally invasive versus abdominal radical hysterectomy for cervical cancer. *N Engl J Med* 2018;379:1895–904.
- National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology: cervical cancer version 1.2020-January 14, 2020. Available: https://www.nccn.org/professionals/physician_gls/pdf/cervical_blocks.pdf [Accessed 1 Apr 2020].
- European Society for Medical Oncology (ESMO) Guidelines Committee. e-Update-Cervical cancer treatment recommendations, 2020. Available: <https://www.esmo.org/guidelines/gynaecological-cancers/cervical-cancer/eupdate-cervical-cancer-treatment-recommendations> [Accessed 9 Jul 2020].
- Querleu D, Cibula D, Concin N, *et al.* Laparoscopic radical hysterectomy: a European Society of Gynaecological Oncology (ESGO) statement. *Int J Gynecol Cancer* 2020;30.
- FIGO Gynecologic Oncology Committee. FIGO statement on minimally invasive surgery in cervical cancer. *Int J Gynaecol Obstet* 2020;149.
- Nitecki R, Ramirez PT, Frumovitz M, *et al.* Survival after minimally invasive vs open radical hysterectomy for early-stage cervical cancer. *JAMA Oncol* 2020;6.
- Wallin E, Flöter Rådestad A, Falconer H. Introduction of robot-assisted radical hysterectomy for early stage cervical cancer: impact on complications, costs and oncologic outcome. *Acta Obstet Gynecol Scand* 2017;96:536–42.
- Cusimano MC, Baxter NN, Gien LT, *et al.* Impact of surgical approach on oncologic outcomes in women undergoing radical hysterectomy for cervical cancer. *Am J Obstet Gynecol* 2019;221:1–24.
- Doo DW, Kirkland CT, Griswold LH, *et al.* Comparative outcomes between robotic and abdominal radical hysterectomy for IB1 cervical cancer: results from a single high volume institution. *Gynecol Oncol* 2019;153:242–7.
- Kim SI, Cho JH, Seol A, *et al.* Comparison of survival outcomes between minimally invasive surgery and conventional open surgery for radical hysterectomy as primary treatment in patients with stage IB1-IIA2 cervical cancer. *Gynecol Oncol* 2019;153:3–12.
- Paik ES, Lim MC, Kim M-H, *et al.* Comparison of laparoscopic and abdominal radical hysterectomy in early stage cervical cancer patients without adjuvant treatment: ancillary analysis of a Korean Gynecologic Oncology Group study (KGOG 1028). *Gynecol Oncol* 2019;154:547–53.
- Yuan Z, Cao D, Yang J, *et al.* Laparoscopic vs. open abdominal radical hysterectomy for cervical cancer: a single-institution, propensity score matching study in China. *Front Oncol* 2019;9:1–7.
- Chen C, Liu P, Ni Y, *et al.* Laparoscopic versus abdominal radical hysterectomy for stage IB1 cervical cancer patients with tumor size ≤ 2 cm: a case-matched control study. *Int J Clin Oncol* 2020.
- Uppal S, Gehrig PA, Peng K, *et al.* Recurrence rates in patients with cervical cancer treated with abdominal versus minimally invasive radical hysterectomy: a multi-institutional retrospective review study. *J Clin Oncol* 2020;38:1030–40.
- Melamed A, Margul DJ, Chen L, *et al.* Survival after minimally invasive radical hysterectomy for early-stage cervical cancer. *N Engl J Med* 2018;379:1905–14.
- Chiva L, Zanagnolo V, Querleu D, *et al.* SUCCOR study: an international European cohort observational study comparing minimally invasive surgery versus open abdominal radical hysterectomy in patients with stage IB1 cervical cancer. *Int J Gynecol Cancer* 2020;30:1269–77.
- Alfonzo E, Wallin E, Ekdahl L, *et al.* No survival difference between robotic and open radical hysterectomy for women with early-stage cervical cancer: results from a nationwide population-based cohort study. *Eur J Cancer* 2019;116:169–77.
- Wenzel HHB, Smolders RGV, Beltman JJ, *et al.* Survival of patients with early-stage cervical cancer after abdominal or laparoscopic radical hysterectomy: a nationwide cohort study and literature review. *Eur J Cancer* 2020;133:14–21.