Abstracts

PR. Median PFS was 4.1 months (95%CI: 2.1–11.3M), and median OS was 33.7M (95%CI: 13.5–33.7M, figure 1).

Conclusion The tumor dormancy was probably maintained by long administration with mild toxicities. These cases demonstrate the use of mCPA-BEV with minimal toxicity and expected anti-cancer activity and indicate that this regimen could be considered for the second-line chemotherapy in advanced recurrent cervical cancer.

IGCS20_1380

355 RADICAL VS. SIMPLE HYSTERECTOMY: A RETROSPECTIVE STUDY ON THE SURVIVAL OUTCOMES OF CERVICAL CANCER PATIENTS

R Rivera*, L Cole. *JRRMMC, Philippines*

10.1136/ijgc-2020-IGCS.305

Cervical cancer remains to be the most common gynecologic malignancy among Filipino women despite being a preventable disease. Radical hysterectomy with pelvic lymphadenectomy is considered the standard surgical treatment of choice for patients with cervical cancer confined to the cervix up to the upper vagina. However, recent studies show that a less radical approach can be offered to these patients with comparable outcomes to radical hysterectomy, but with lesser perioperative and post-operative morbidity. The purpose of this study was to compare the outcomes in terms of recurrence and survival

among cervical cancer patients who underwent simple hysterectomy and radical hysterectomy seen in a tertiary government hospital. Records of all cervical cancer patients who underwent radical hysterectomy and simple hysterectomy for the past ten years were reviewed. The incidence of cervical cancer patients who underwent simple hysterectomy from 2009–2018 is 0.37 per 100 person years or 0.592:16, lower than 1:16 ratio from 1964–1974, as reported by Manalo and Soto. Only 9 out of 42 patients who underwent simple hysterectomy had cervical cancer screening within 1 year prior to surgery. The most common indication for surgery was myoma uteri. Those who underwent radical hysterectomy had better recurrence free survival and overall survival than those who had simple hysterectomy, but among low risk patients, those with 2 cm or less tumor size with no other risk factors, there was no significant difference in survival outcomes between the two groups.

IGCS20_1381

356 PAPILLARY SQUAMOUS CARCINOMA OF THE CERVIX WITH METACHRONOUS CLEAR CELL RENAL CELL CARCINOMA

R Medalla*, J Luna. *Philippine General Hospital, Philippines*

10.1136/ijgc-2020-IGCS.306

Multiple primary tumors can be classified as synchronous or metachronous. Cases have been reported, with a prevalence,

Abstract 356 Figure 1 A. Low Power Objective (100x magnification) of Papillary Squamous Cell Carcinoma with tumor cells arranged in nests and papillary configuration. B. High Power Objective (400x). These tumor cells exhibit moderate pleomorphism, with enlarged, round to oval, nuclei, some with prominent nucleoli, and abundant eosinophilic cytoplasm and distinct cell borders

Abstract 356 Figure 2 Intraoperative pictures. A. The left kidney was converted to a solid, necrotic mass measuring 12.0 × 10.0 × 5.0 cm. B. Cut section of the resected left kidney