patients with EC and healthy female subjects. MiRNA was isolated and qPCR was used to detect expression levels of miRNAs.

Results A total of 76 women were included; 36 EC patients, 40 healthy controls. A distinct panel of miR-200a, miR-200b, miR-200c, miR-205 and miR-182 showed AUC of 0.958, sensitivity 92%, specificity 89%, positive predictive value of 89% and negative predictive value of 91% in diagnosing EC. MiR-182 expression levels were significantly related to high-grade endometrioid tumours compared to low grade.

Conclusion We demonstrated high diagnostic accuracy of miRNA for detecting EC. In addition, miRNA contributed to improved distinguishing between high-grade and low-grade endometrioid tumours. Validation of miRNA expression levels in urine will be performed in order to further optimise a non-invasive diagnostic tool.

IGCS20_1465

428 PROSPECTIVE EVALUATION OF AN ERAS PATHWAY AT A GYNAECOLOGICAL ONCOLOGY UNIT IN OSLO, NORWAY

^{1,2}K Lindemann*, ³B Eijólfssdóttir, ⁴S Heimisdóttir Danbolt, ¹YY Wang, ⁵AG Heli-Haugestøl, ⁵H Thorsrud, ²TK Nguyen, ⁶O Mjälund, ¹GA Navestad, ¹S Hermanrud, ⁷K E Juul-Hansen, ^{8,9}A Kleppe, ¹T Skeie-Jensen, ^{2,7}U Kongsgaard. ¹Department of gynecologic oncology, Oslo University Hospital; ²Institute of clinical medicine, University of Oslo; ³Department of gynecologic oncology, Oslo University Hospital; ⁴Institute of health and society, University of Oslo; ⁵Department of Clinical Service, Oslo University Hospital; ⁶Department of gastroenterological surgery, Oslo University Hospital; ⁷Department of anesthesiology, Oslo University Hospital; ⁸Institute for Cancer Genetics and Informatics, Oslo University Hospital; ⁹Department of Informatics, University of Oslo, Oslo, Norway

10.1136/ijgc-2020-IGCS.371

Introduction We prospectively evaluated an ERAS in a large gynaecologic oncology tertiary centre serving the South East Health region in Norway.

Methods Patients undergoing laparotomy for (suspected) ovarian cancer at the Oslo University Hospital were prospectively included in a pre-implementation and post-implementation cohort. Baseline characteristics, adherence to the pathway and clinical outcomes were assessed.

Results Of the 439 patients, 235 (54%) underwent surgery for advanced ovarian cancer and 204 (46%) for a suspicious ovarian mass. Median fasting times for solids (13 vs 16 h, $p<0.001$) and fluids (3.7 vs 11.9 h, $p<0.001$) were significantly reduced. Perioperative fluid administration varied less ($p<0.001$) and was reduced (median 11.5 vs 15.8 ml/kg/h, $p<0.001$). Epidural analgesia was the mainstay of analgesia in both cohorts, but with ERAS more patients received continuous vasopressor intraoperatively (87% vs 70%, $p=0.003$). More patients received dual PONV prophylaxis (85% vs 64%, $p<0.001$). Length of stay remained unchanged for patients with advanced disease with a median of 5 days both before and after the implementation ($p=0.94$), but patients undergoing surgery for an ovarian mass stayed shorter in hospital after the implementation of ERAS ($p=0.026$). For all patients, more patients were discharged directly home as opposed to transferred to local referring hospital after ERAS implementation (70% vs 51%, $p<0.001$). There was no difference in re-admission rates or postoperative 30 d morbidity.

Conclusion Introduction of an ERAS pathway resulted in less variance in practice and increased adherence to current standards in perioperative management. Patients were more often able to be discharged directly home without an increase in admission rates.

IGCS20_1466

429 IDENTIFICATION OF CLINICOPATHOLOGICAL PROGNOSTIC FACTORS IN RECURRENT ENDOMETRIAL CANCER

Y Koike*, M Takenaka, J Suzuki, Y Shoburu, K Tomita, A Ikenaga, M Kamii, S Hirose, H Takano, K Yamada, S Niimi, A Okamoto. *The Jikei University School of Medicine, Japan*

10.1136/ijgc-2020-IGCS.372

Objectives Although, endometrial cancer (EC) is commonly diagnosed at an early stage and has a favorable prognosis, recurrent disease usually shows poor prognosis. However, clinicopathological prognostic factors for recurrent EC have been still unclear. The current study aimed to identify clinicopathological prognostic factors, especially in long term survivors of recurrent EC.

Methods We collected 2044 patients who underwent surgery including hysterectomy and were pathologically diagnosed as EC in our hospitals between 2001 and 2018. Clinicopathological information and survival data were retrospectively obtained by the review of medical record. Among 2044, 131 patients were included in this retrospective analysis. In recurrent cases, we analyzed the prognostic factors for long-term survivors of recurrent EC. Then we defined long-term and short-term survivors, as patients who survived ≥ 3 years ($n=53$) and died within 3 years ($n=78$) after first recurrence, respectively. The correlation between prognosis and clinicopathological factors was statistically analyzed.

Results In the multivariate analysis of long-term survivors, we identified that TTP ≥ 1 year after surgery ($P<0.01$), surgical reduction of the first recurrence ($P=0.03$), non-peritoneal metastasis ($P=0.045$), were significantly associated with the long-term survivors of recurrent EC.

Conclusion TTP ≥ 1 year after surgery, non-peritoneal metastasis, surgical reduction of first recurrence were good prognostic factors of long-term survivors in recurrent EC.

IGCS20_1467

430 MACHINE LEARNING MODELS TO PREDICT SURVIVAL OUTCOMES AFTER RADICAL HYSTERECTOMY ACCORDING TO SURGICAL APPROACH IN PATIENTS WITH FIGO STAGE IB CERVICAL CANCER

¹S Kim*, ²C Choi, ¹M Lee, ³DH Suh, ¹J Kim, ³YB Kim. ¹Seoul National University College of Medicine, South Korea; ²Samsung Medical Center, South Korea; ³Seoul National University Bundang Hospital, South Korea

10.1136/ijgc-2020-IGCS.373

Objective To develop preoperative machine learning models predicting survival outcomes according to the surgical approach in early-stage cervical cancer.

Methods We retrospectively identified patients with FIGO stage IB cervical cancer who underwent either primary open RH or laparoscopic RH at three tertiary institutional hospitals between 2000 and 2018. Patients' clinicopathologic, image, and survival data were collected. The whole dataset was separated into training and test sets with a 4:1 ratio. Combining both statistical and deep neural network models, we constructed hybrid ensemble predictive models for 5-year PFS and OS rates. Only the variables that could be obtained before surgery were used. Model development was conducted in the training set with ten-fold cross-validation, and the developed models were validated in the test set.

Results In total, 1,141 patients were included; 578 and 563 received open RH and laparoscopic RH, respectively. The median length of observation was 57.6 months during which 157 patients (13.8%) experienced disease recurrence and 86 patients (7.5%) died. In terms of preoperative prediction, while the logistic regression model showed AUCs of 0.68 and 0.71 for 5-year PFS and OS rates, respectively, the ensemble model showed better performance: AUCs, 0.71 and 0.78. These models commonly included the surgical approach as the main prognostic factor.

Conclusion We developed preoperative models predicting survival outcomes according to the surgical approach in early-stage cervical cancer. These models will be useful for making decisions in choosing open RH or laparoscopic RH in individualized counseling practices.

IGCS20_1468

431 DISTANT METASTATIC SPREAD AND CLONAL EVOLUTION OF HIGH-GRADE ENDOMETRIAL CANCERS

¹K Miller*, ²P Selenica, ²J Reis-Filho, ¹NR Abu-Rustum, ³V Makker, ³CA Aghajanian, ²CA Iacobuzio-Donahue, ²R Murali, ²B Weigelt. ¹Department of Surgery, Memorial Sloan Kettering Cancer Center, USA; ²Department of Pathology, Memorial Sloan Kettering Cancer Center, USA; ³Department of Medicine, Memorial Sloan Kettering Cancer Center, USA

10.1136/ijgc-2020-IGCS.374

Introduction The incidence and mortality of endometrial cancer (EC) is rising worldwide. The biological underpinnings of EC progression are poorly understood. We sought to characterize the genetic alterations and clonal evolution of two primary high-grade ECs and their matched multiple distant metastases.

Methods Research autopsies were performed on two women: one with treatment-naïve, widely metastatic undifferentiated carcinoma at diagnosis (case 1); and one with serous EC and heavily treated metachronous disease (case 2). Whole-exome sequencing of primary tumors, metastases (n=8 and n=7), and normal tissues was performed and analyzed using validated bioinformatics methods.

Results In case 1, truncal hotspot PIK3CA p.Q546K and PTEN p.R130G mutations, chromosome 8 and 16p losses, and mutational aging signature 1 were present in the primary and all metastases. Low levels of genetic heterogeneity between all samples were observed (70% of mutations

shared). In case 2, we found high levels of genetic heterogeneity and acquisition of mutations during progression. A truncal PIK3CA p.H1047L hotspot mutation was present in the primary tumor and all metastases, with a primary subclonal TP53 frameshift mutation becoming clonal in a subset of metastases (n=3). Evidence of clonal progression between metastatic sites was observed as well as a combination of aging and homologous recombination-deficiency mutational signatures.

Conclusion Genetic alterations identified in case 1 were likely early events in the clonal expansion of the primary tumor, reflective of aggressive disease and absence of treatment. In case 2, evidence of clonal diversity and progression was observed, potentially representing clonal selection due to therapeutic effect.

IGCS20_1471

433 DETERMINING THE HAEMATOLOGICAL MORBIDITY ASSOCIATED WITH CYTOREDUCTIVE SURGERY

¹E Corry*, ²H Bartels, ¹J Mulhall, ¹J Grundy, ¹J Mulsow, ¹D Brennan. ¹Mater Misericordiae University Hospital, Ireland; ²National Maternity Hospital, Ireland

10.1136/ijgc-2020-IGCS.375

Introduction Cytoreductive surgery (CRS) is a complex procedure performed in the setting of epithelial ovarian cancer (EOC), pseudomyxoma peritonei (PMP) and colorectal metastases (CRCm). Due to the complexity of the surgery with significant fluid volume exchange calculating estimated blood loss (EBL) is challenging. To determine the hematological morbidity associated with CRS we reviewed the EBL and transfusion rates in a National Peritoneal Cancer Centre.

Method The surgical oncology anonymized database provided data on demographics, surgical oncological intervention, timing, EBL, laboratory measurements, and transfusion requirements. The EBL recorded from the operative notes calculated in the operating room by the gravimetric method (suction volume and swab weights).

Results A total of 120 patients who underwent CRS±HIPEC, with a median age of 57 were included. The median EBL for the entire cohort (n=120) was 981 ml (range 50–6500). The median EBL for metastatic colorectal CRS (n=38) was 830 ml (range 50–3800 ml), for ovarian cancer (n= 51) was 900 ml (range 150–4600 ml) and for PMP (n=31) was 1300 (range 100–6500). The average number of red cells (RCC) transfused for the entire cohort was 1.14 (SEM 0.147), and was similar for ovarian cancer (1.28 (SEM 0.251) and PMP (1.26 (SEM 0.314)) but was significantly lower in the colorectal metastases group (0.87 (SEM 0.189)).

Conclusion CRS for EOC, PMP, or CRCm is associated with significant hematological morbidity, which should be taken into account for pre-operative optimization. The variation in EBL reflects the heterogeneous nature of these complex procedures, with more extensive surgery often warranted in PMP and EOC.