IGCS20_1472

434 CORRELATION OF ENDOMETRIAL TUMOR DNA MISMATCH REPAIR STATUS AND MELF-PATTERN INVASION

C DiNobile*, E Kertowidjojo, X Wen, C Tornos, G Gossner, W Burke, M Pearl. *Stony Brook University Hospital, USA*

10.1136/ijgc-2020-IGCS.376

Introduction Endometrial adenocarcinoma is the most common gynecologic malignancy in the United States. Between 10–30% of endometrial adenocarcinomas exhibit microsatellite instability (MSI), a type of genetic hypermutability that results from impaired DNA mismatch repair (MMR). On pathologic review, MSI can be indirectly identified by immunohistochemical (IHC) staining for deficient MMR protein expression. Another component of pathologic review is to assess for microcystic, elongated, and fragmented (MELF) pattern of invasion, characterized by a fibromyxoid stromal reaction and the formation of microcysts. To date, there are no studies assessing the correlation of MMR status and MELF-pattern invasion in endometrial adenocarcinoma.

Methods We performed an IRB-approved, retrospective review of medical records and pathology slides of surgical cases of endometrioid endometrial adenocarcinoma between January 2016 and January 2020.

Results Our results did not demonstrate a correlation between MMR mutation status and the presence of MELF-pattern invasion. The presence of MMR mutation was associated with age, stage of disease, and a history of stroke. The presence of

MELF-pattern invasion was not associated with various clinical factors or comorbidities.

Conclusions Our study did not demonstrate a relationship between MMR status and presence of MELF pattern invasion. We also did not re-demonstrate prior findings that MELF-pattern invasion is associated with higher rates of lymph node metastases or lymphovascular space invasion (LVSI). Future directions include investigation into the relationship among MMR and age, stage of disease, and history of stroke, as this could potentially impact treatment planning.

IGCS20_1473

435 PALLIATIVE ELECTROCHEMOTHERAPY IN VULVAR CANCER: A DOUBLE INSTITUTION STUDY AND REVIEW OF THE LITERATURE

¹G Corrado*, ²G Cutillo, ¹S Fragomeni, ²V Bruno, ³L Tagliaferri, ²E Mancini, ²C Certelli, ²E Vizza, ¹G Scambia, ¹G Garganese. ¹Gynaecologic Oncology Unit, Fondazione Policlinico Universitario A. Gemelli – IRCCS, Italy; ²Gynaecologic Oncology Unit, IRCCS 'Regina Elena' National Cancer Institute, Italy; ³Radioterapia Oncologica ed Ematologia – Gemelli ART (Advanced Radiation Therapy), Interventional Oncology Center (IOC), Fondazione Policlinico Universitario A. Gemelli – IRCCS, Italy

10.1136/ijgc-2020-IGCS.377

Introduction The aim of this study is to revise the experience of two reference centers for vulvar cancer treatment during the last 2 years, evaluating the ECT procedure in terms of clinical outcome and side effects profile.

Abstract 435 Figure 1