metastasis in primary low-grade ovarian cancer. Further prospective trials evaluating LVSI and Ki-67 as a predictor of lymph-node metastasis should be planned.

IGCS20_1508

464

COMPARISON OF DIFFERENT METHODS TO DETERMINE MYOMETRIAL INVASION IN ENDOMETRIAL CANCER – A NATIONWIDE SWEGCG STUDY

¹J Björg*, ²M Björberg, ³C Borgfeldt, ⁴P Dahm-Kähler, ⁵A Flöter-Rådestad, ⁶K Hellman, ⁷E Hjerpe, ⁸E Holmberg, ⁹P Kjölhede, ¹⁰J Marcinkiewicz, ¹¹P Rosenberg, ¹²B Tholander, ¹³E Åvall-Lundquist, ¹⁴K Ståhlberg, ¹⁵T Högberg. ¹Department of Women's and Children's health, Uppsala University, Sweden; ²Department of Hematology, Oncology and Radiation Physics, Skåne University Hospital, and Department of Clinical Sciences, Lund University, Sweden; ³Department of Obstetrics and Gynecology, Skåne University Hospital and Lund University, Sweden; ⁴Department of Obstetrics and Gynecology, Sahlgrenska University Hospital, Sweden; ⁵Department of Women's and Children's Health, Division of Obstetrics and Gynecology, Karolinska Institutet, Karolinska University Hospital, Sweden; ⁶Department of Gynecologic Cancer, Theme Cancer, Karolinska University Hospital, Sweden; ⁷Department of Obstetrics and Gynecology, Visby Hospital, Sweden; ⁸Region Västra Götaland, Regional Cancer Centre West, Sweden; ⁹Department of Obstetrics and Gynecology in Linköping, and Department of Biomedical and Clinical Sciences, Sweden; ¹⁰Department of Obstetrics and Gynecology, Halland Hospital, Sweden; ¹¹Department of Oncology, and Department of Biomedical and Clinical Sciences, Sweden; ¹²Department of Oncology, Uppsala University Hospital, Sweden; ¹³Department of Oncology, and Department of Biomedical and Clinical Sciences, Linköping University, Sweden; ¹⁴Department of Cancer Epidemiology, Lund University, Sweden

10.1136/ijgc-2020-IGCS.402

Background Deep myometrial invasion (MI) ($\geq 50\%$) is a prognostic factor for lymph node metastases and poorer survival in endometrial cancer. There is no consensus regarding which pre/peroperative diagnostic method should be preferred.

Aim To explore the pattern of different diagnostic methods for MI assessment in Sweden and to evaluate differences between MRI, vaginal ultrasound, frozen section and gross examination in clinical practice.

Methods Women with endometrial cancer registered in the Swedish Quality Registry for Gynecologic Cancer (SQRGC) between January 2010 and December 2019 were eligible. Inclusion criteria were FIGO stage I-III and available information on histology and on assessment of MI. Data on age, histology, FIGO stage, degree of MI, histology results, method for MI assessment and hospital level were collected from the SQRGC. The final assessment by the pathologist on specimens from hysterectomy was golden standard.

Results The study population included 1,950 women, 33% (n=649) had a MI $\geq 50\%$. The methods used for MI assessment were vaginal ultrasound in 54%, MRI in 22%, gross examination in 13% and frozen section in 11% of cases. Age, histology or FIGO stadium did not differ between the methods. The sensitivity, specificity and accuracy of vaginal ultrasound was 61.2%, 83.3% and 0.75% respectively, and for MRI 74.2%, 72.7% and 0.73%. The highest accuracy was for frozen section; 95.0%.

Conclusion The assessment of deep myometrial invasion in endometrial cancer is most often performed with vaginal ultrasound in Sweden. The sensitivity of this method is lower in clinical practice than for MRI and perioperative methods.

IGCS20_1511

465

DELAYS IN TREATMENT IN GYNAECOLOGY ONCOLOGY PATIENTS IN QATAR SEEKING MANAGEMENT OVERSEAS

S Brich*, T Alsayed, A Ansari, H ElMalik, J Herod. Hamad medical corporation, Qatar

10.1136/ijgc-2020-IGCS.403

Introduction The gynaecological oncology service in Doha treats all women living in or visiting Qatar. Despite the quality and affordability of the service many women travel overseas for their treatment following diagnosis or present following previous treatment overseas requesting further management. Although they must perceive potential advantages which encourage them to do so, there are difficulties which could arise including delay in treatment of a malignancy that could affect their outcomes. We wished to understand the impact of travel overseas on the waiting time for treatment.

Methods All patients seen over a period of 3 yrs who had travelled overseas were identified. Records were reviewed to identify what impact the decision to travel abroad had made on the timing of their treatment. According to Qatari cancer treatment standards, treatment should be within 14 days of a decision made by MDT. We considered that a delay in treatment would reasonably be defined as an interval of >4 weeks.

Results 18% of patients (n=153/850) with a recorded care plan by the MDT sought medical treatment overseas between 4/2015 and 3/2018. Patients had 25 different nationalities; Qatari nationals represented the majority (40.5%). Patients travelled to 28 different destinations. Most travelled to the U.S.A(15.7%), Philippines(15%), the UK(10.5%) and Thailand (9.2%).

23.5% of patients had a delay in treatment; 9,2% had an unknown treatment timing plan. Most had delays of <6 weeks; 10% had significant delays of many weeks, months and even >1 year.

Conclusion The decision to travel overseas in our patients resulted in delays of treatment for roughly 1/4 of patients. In 10% these delays would be expected to have an adverse effect on outcomes.

IGCS20_1512

SIMVASTATIN MODIFIES THE INTERNALIZATION, ENDOCYTIC TRAFFICKING, AND THE CONTENT OF OVARIAN CANCER CELL-DERIVED EXTRACELLULAR MICROVESICLES WHICH ARE RESPONSIBLE OF INDUCING MIGRATION AND INVASION IN VITRO

¹P Mancilla, ¹MF Liberona, ¹S Kato, ²J Barra, ²A Gonzalez, ¹M Cuello*. ¹Department of Gynecology, School of Medicine, Pontificia Universidad Católica de Chile CEBICEM, Santiago Chile; ²Universidad San Sebastián, Santiago Chile

10.1136/ijgc-2020-IGCS.404

Introduction High grade serous ovarian carcinoma (HGSOC) is the leading cause of death among all gynecological malignancies. Extracellular microvesicles (MVs) are secreted by most cells in the body and play a crucial role regulating cell-to-cell communication and several biological functions. Current