prognostic and predictive value of the modeled CA-125 elimination rate constant, KELIM.

Methods KELIM was estimated from treatment-related pharmacodynamic modeling of CA-125 values. Median KELIM was used to define favourable ($\geq$ median)/unfavourable ($<$ median) KELIM groups. Patients were analyzed by surgery type: primary (PDS) or interval (IDS) debulking surgery.

Results In the IDS population (N=154), patients with favourable KELIM had a higher frequency of complete surgery vs unfavourable KELIM (51.9% vs 32.4%), confirming KELIM as a chemosensitivity marker. In both PDS (N=700) and IDS populations, median PFS was longer with favourable KELIM vs unfavourable KELIM, demonstrating a prognostic value. In the PDS population, median PFS was longer in the veliparib-throughout arm relative to control irrespective of KELIM (29.6 vs. 20.9 and 18.2 vs 15.4 months in favourable and unfavourable KELIM groups, respectively; figure 1). In the IDS population, median PFS was longer with veliparib-throughout vs control for patients with favourable KELIM only (29.3 vs 20.8 months; figure 2).

Conclusion In VELIA, KELIM was prognostic for PFS and IDS outcomes. Current data suggest KELIM may be associated with veliparib benefit. Ongoing analyses will explore how baseline characteristics contribute to KELIM predictive/prognostic value.

IGCS20_1165

37 QUANTITATIVE EVALUATION OF LYMPH-VASCULAR SPACE INVASION (LVSI) IN PATIENTS AFFECTED BY ENDOMETRIAL CANCER: PROGNOSTIC AND CLINICAL IMPLICATIONS

¹S Restaino*, ¹L Tortorella, ¹E Perrone, ¹G Monterossi, ²A Gioè, ²I Capasso, ²E La Fera, ²G Dinoi, ²G Scambia, ²F Fanfani, ³G Zannoni, ⁴V Chiantera. ¹Fondazione Policlinico Universitario A. Gemelli IRCCS, UOC di Ginecologia Oncologica, Dipartimento per la Tutela della Salute della Donna e della Vita Nascente, Italy; ²Università Cattolica del Sacro Cuore, Italy; ³Department of Histopathology, Università Cattolica del S. Cuore, Italy; ⁴Department of Gynecologic Oncology, University of Palermo, Italy

10.1136/ijgc-2020-IGCS.37

Objectives Lymph-vascular space invasion (LVSI) is associated with an increased risk of recurrence. Usually, the interpretation of LVSI is just qualitative, as presence or absence. The aim of this study is to examine the quantitative analysis of LVSI.

Methods Retrospective multicentre study. It included 2300 consecutive patients who received a histologically confirmed diagnosis of endometrial cancer between January 2000 and December 2018 at the Gynaecologic Oncology Unit of two tertiary hospitals in Italy.

Results The rate of lymph node metastasis increased from the 5% in patients with no LVSI, to 15% in patients with focal LVSI and 33% in those with diffuse LVSI ($p < 0.001$). There is a progressive increase of prognostic impact of focal and diffuse LVSI on the risk of node metastasis (AOR = 2.12, 95% CI 1.19, 3.79 for focal LVSI and AOR = 3.67, 95% CI 2.30, 5.86). Distant recurrences were more frequent in patients with diffuse LVSI compared with no LVSI (24.9% vs 6.6% respectively, $p < 0.001$). We found

that adjuvant treatments, specifically adjuvant radiations, were associated with improved OS and DFS (median DFS = 19 months vs 97 months respectively for no adjuvant and radiation therapy, $p < 0.001$) in patients with diffuse LVSI.

Conclusion The presence of diffuse LVSI is an independent risk factor of both lymph node metastasis and distant recurrence in endometrial cancer patients. It is associated with decreased OS and DFS. Adjuvant radiation improved survival regardless of grading, histotype, and lymph nodal metastasis in women with diffuse LVSI.

IGCS20_1184

38 A TRAINING PROGRAM TO BUILD CAPACITY FOR CERVICAL CANCER PREVENTION IN MOZAMBIQUE

¹M Salcedo*, ²M Varon, ²M Varon, ²E Baker, ²E Baker, ³N Osman, ⁴E David, ⁵R Rangeiro, ⁵D Changule, ⁶V Andrade, ⁷CM Oliveira, ⁸A Neves, ⁹J Carns, ⁵C Lorenzoni, ²KM Schmeler, ²KM Schmeler. ¹Obstetrics and Gynecology Department, Federal University of Health Sciences of Porto Alegre/Irmandade Santa Casa de Misericórdia de Porto Alegre, Brazil; ²Department of Obstetrics and Gynecology, King Chulalongkorn Memorial Hospital, Thailand; ³Universidade Eduardo Mondlane (UEM), Mozambique; ⁴Ministerio da Saude de Moçambique (MISAU), Mozambique; ⁵Hospital Central de Maputo, Mozambique; ⁶Hospital de Câncer de Barretos, Brazil; ⁷Diagnóstico das Américas-DASA, Brazil; ⁸Hospital Geral José Macamo, Mozambique; ⁹Department of Bioengineering, Rice University, USA

10.1136/ijgc-2020-IGCS.38

Introduction Cervical cancer is the primary cause of cancer among women in Mozambique. There is a shortage of providers trained to deliver cervical cancer screening and manage pre-invasive lesions. We describe a training program to build local capacity to deliver quality cervical cancer prevention services in Mozambique.

Methods The program includes training courses led by faculty from the United States (US) and Brazil, and consists of lectures followed by hands-on training stations to practice colposcopy, cervical biopsy, ablation and loop electrosurgical excision procedure (LEEP) using innovative training models. Participants then perform the procedures in clinic with supportive supervision from the international faculty. The courses are complemented by monthly videoconference tele-mentoring sessions, held in Portuguese using the Project ECHO® (Extension for Community Healthcare Outcomes) model.

Results From 2016 to 2020, 10 courses were held in the cities of Maputo (n=8), Beira (n=1) and Nampula (n=1). There were 347 participants with an average of 34 participants per course. The courses have recently transitioned from only international faculty to include Mozambican gynecologists, including two fellows from the IGCS Global Curriculum program. A total of 15 ECHO sessions were held with ~25 participants/session, 30 patient cases discussed and 14 lectures delivered.

Conclusion This collaboration between Mozambique, Brazil and the US is building local capacity to prevent cervical cancer through training, mentoring and support of local providers. Due to the COVID-19 pandemic, the courses are being transitioned to a virtual format led by the Mozambican doctors with the international faculty joining remotely.