progression-free survival was 18 months, median overall survival was 41 months.

Conclusion Treatment with bevacizumab in daily clinical practice is safe and effective- in concordance with published data from prospective studies GOG 218 and ICON 7.

IGCS20_1090

118

EPIDEMIOLOGY OF CERVICAL CANCER IN THE REPUBLIC OF BELARUS

O Matylevich*, A Shushkevich, D Sharko, A Evmenenko. *NNAlexandrov National Cancer Center of Belarus, Belarus*

10.1136/ijgc-2020-IGCS.101

Introduction In Belarus, Cervical Cancer (CC) is the 7th most common cancer accounting for 5.8% of all new cancer cases.

Objectives The aim of the study was to estimate incidence, mortality and survival rate of newly diagnosed CC in Belarus from 2009 to 2018.

Methods The data from the Belarusian Cancer Registry were analyzed.

Results The estimated age-standardized incidence rate of CC per 100000 female population in Belarus has decreased from 12.8 in 2009 to 11.2 in 2018 ($p < 0.01$). Mortality decreased by 12.1% ($p > 0.05$).

Comparison of two five-year periods (2009–2013 and 2014–2018) showed that rate of locally advanced (III-IV stage) CC increased from 25.9% to 33.9%, as rate of stage I and II decreased by 7% and 16%, respectively.

Comparison of 5-year adjusted survival rates between 2013 and 2018 showed no change for CC stage I (92.8%, SE 0.8% and 93.3%, SE 0.9%, respectively), increased by 13.5% for stage II (57.6%, SE 1.6% and 65.4%, SE 1.8%), increased by 9.4% for stage III (40.3%, SE 2.3% and 44.1%, SE 2.1%), and increased by 67% for stage IV (9.4%, SE 1.9% and 15.7%, SE 2.2%).

Conclusions With improvement in diagnostics and introduction of the pilot projects of CC screening in some regions of the country over the last decade, CC incidence rates in Belarus have decreased by 12.5%, primarily due to decrease in incidence of early stages of CC. While there was no significant decrease in mortality, adjusted survival for advanced CC stages was improved.

IGCS20_1091

119

POSTRADIATION CUTANEOUS ANGIOSARCOMA AFTER TREATMENT OF BREAST CARCINOMA

S Sakhrî*, M Bouheni, H Bouaziz, R Chargui, K Rahal. *Slah Azaiez institut, surgicaldepartment, Tunis, Tunisia, Tunisia*

10.1136/ijgc-2020-IGCS.102

Introduction Postradiation cutaneous angiosarcoma of the breast (PRCA) is very rare malignancy with an incidence ranged from 0.09 to 0.16%. It usually occurs after adjuvant radiotherapy following the treatment of breast cancer. In article published in August 2019 found about 50 cases of PRCA, the first case was published in 1948.

Case Report A 40-year-old woman was referred for a palpable right breast mass. She denied history of breast masses or issues.

Physical examination revealed a palpable right breast mass located at upper-outer right breast quadrant.

There were no other palpable breast masses, no palpable axillary, supra-clavicular, or infra-clavicular lymph nodes. The patient underwent a right breast core biopsy of the mass the pathology of the biopsy revealed invasive ductal carcinoma. She underwent a right lumpectomy with a right axillar lymph node dissection.

There was no metastatic lymph node. the patient underwent standard external radiotherapy (50 gy to the whole breast and 16 gy as a boost to the tumor bed) and chemotherapy then hormonotherapy by tamoxifen.

She was followed with clinical and radiological surveillance. Seven years after the surgery she was presented with erythematous plaque associate with multiple purplish nodules, the mammography and the ultrasound was normal, the patient underwent a biopsy, the histological examination showed a post radiation cutaneous angiosarcoma.

Conclusion The PRCA is rare and aggressive disease, the diagnostic is clinically difficult and often missed on mammography, usually the biopsy confirm the diagnostic.

IGCS20_1093

121

SYSTEMIC IMMUNE-INFLAMMATORY INDEX (SII) AS A PROGNOSTIC FACTOR FOR CERVICAL CANCER RECURRENCE

¹A De Guzman, ¹K Baldivia, ²GM Facun*, ²C Zalameda - Castro. ¹Division of Radiation Oncology, Department of Radiology, University of the Philippines – Philippine General Hospital, Philippines; ²Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, University of the Philippines – Philippine General Hospital, Philippines

10.1136/ijgc-2020-IGCS.103

Objectives Numerous studies on inflammatory hematologic markers have been published in relation to cancer survival and progression, but only two publications have studied the Systemic Immune-Inflammatory Index (SII) among cervical cancer patients. This study aims to validate the cut-off presented by Huang et al., (2019) in his study on cervical cancer and the prognostic ability of the Systemic Immune-Inflammatory Index (SII).

Methodology Data from 140 patients diagnosed with cervical cancer recurrence and those without evidence of disease post-treatment were collected retrospectively. The SII, Neutrophil-Lymphocyte Ratio, Platelet-Lymphocyte Ratio, and Monocyte-Lymphocyte Ratio were all evaluated.

Results Based on our univariate Cox Hazard Analysis, bulky tumor (>4 cm) and FIGO stages II and III were prognosticators of worse Progression Free Survival (PFS). Regarding SII, there appears to be an increased likelihood of disease recurrence among women with SII greater than 475 when controlling for the binary classification of the SII (HR: 1.88 (0.96–3.69), $p = 0.07$). However, this association may be diluted when other variables aside from the SII are accounted for in determining the likelihood of the outcome. On multivariate analysis, only FIGO stage was seen as an independent factor for PFS.