University of the Philippines - Philippine General Hospital. A preliminary analysis comparing OVERA and other strategies for malignancy risk prediction was performed.

Results As of this report, 347 women have been enrolled in the study. Based on sonologic classifiers, high risk patients had higher predictive scores using LR1 ($\chi^2[2]$: 260.81, $p < 0.01$), LR2 ($\chi^2[2]$: 271.57, $p < 0.01$), Sassone ($\chi^2[2]$: 127.26, $p < 0.01$), Lerner ($\chi^2[2]$: 153.98, $p < 0.01$), IOTA-ADNEX ($\chi^2[2]$: 215.22, $p < 0.01$), and ROMA ($\chi^2[2]$: 144.48, $p < 0.01$). Based on categorical classifiers, CA-125 (χ^2 : 34.59, $p < 0.01$), OVERA (χ^2 : 54.25, $p < 0.01$), and ROMA (χ^2 : 85.29, $p < 0.01$) discerned well in the high-risk group, while HE4 (χ^2 : 105.10, $p < 0.01$) discerned lower risk better. OVERA as a numerical variable appears to fare well in detecting the likelihood of malignant tumors ($\chi^2[4]$: 3.38, p : 0.50) as compared to sonologic models but was not as discerning as other models using the conventional cut-off value (5.0). A higher cut-off (6.9) would confer more optimal values of specificity across age (pre-menopausal or menopausal) and histopathologic types.

Conclusion The preliminary results stress the importance of population based studies in evaluating biochemical assays for ovarian cancer risk prediction.

IGCS20_1148

162

ABDOMINAL RADICAL TRACHELECTOMY VERSUS CHEMOTHERAPY FOLLOWED BY VAGINAL RADICAL TRACHELECTOMY IN STAGE 1B2 (FIGO 2018) CERVICAL CANCER. A SYSTEMATIC REVIEW ON FERTILITY AND RECURRENCE RATES

¹K Van Kol*, ²V Tineke, ¹R Bekkers. ¹Catharina Cancer Institute, Catharina Hospital, Netherlands; ²Radboud University Medical Center, Netherlands

10.1136/ijgc-2020-IGCS.141

Introduction There is currently no standard of care for women with cervical cancer stage IB2 (FIGO 2019, diameter 2–4 cm) who wish to preserve their fertility. Generally, two approaches are offered. Option 1: neoadjuvant platinum-based chemotherapy (NACT) to reduce the tumor size to ≤ 2 cm, followed by Vaginal Radical Trachelectomy (VRT) with Pelvic Lymph Node Dissection (PLND) either before chemotherapy or at the time of VRT. Option 2: Abdominal Radical Trachelectomy (ART) with PLND.

Objective To compare rates of fertility, pregnancy, live births as well as recurrence for women with cervical cancer stage IB2 treated with either NACT followed by VRT, or ART.

Methods A systematic review was performed using the PubMed database. Articles reporting the search term 'trachelectomy' as text word or as Medical Subject Headings (MeSH) were identified.

Results Ten studies were identified with a total of 338 patients. After NACT followed by VRT 39% of the women tried to conceive, 70% of these women got pregnant, of which 63% resulted in a live birth. The overall recurrence and death rate were 10% and 2.9% respectively. After ART 40% of the women tried to conceive, 21% of these women got pregnant, which resulted in a live birth rate of 42%. Recurrence and death rate after ART were 6.9%, and 3.4% respectively.

Conclusion Women with cervical cancer stage IB2 and a wish to preserve fertility treated with NACT followed by VRT have a significantly higher chance of pregnancy than women treated with ART, with comparative oncological results.

IGCS20_1150

164

ENGOT-CX11/GOG 3047/KEYNOTE-A18: A PHASE 3, RANDOMIZED, DOUBLE-BLIND STUDY OF PEMBROLIZUMAB WITH CHEMORADIO THERAPY IN PATIENTS WITH HIGH-RISK LOCALLY ADVANCED CERVICAL CANCER

¹D Lorusso*, ²N Colombo, ³R Coleman, ⁴L Randall, ⁵L Duska, ⁶Y Xiang, ⁷K Hasegawa, ⁸A Nogueira-Rodrigues, ⁹D Cibula, ¹⁰M Raza Mirza, ¹¹B You, ¹²A Oaknin, ¹³M Christiaens, ¹⁴C Taskiran, ¹⁵J Sehouli, ¹⁶J Korach, ¹⁷C Martin, ¹⁸S Keefe, ¹⁸M Puglisi, ¹⁹S Pignata. ¹Università Cattolica del Sacro Cuore-Fondazione Policlinico Universitario, Italy; ²Istituto Europeo di Oncologia, Italy; ³University of Texas MD Anderson Cancer Center, USA; ⁴Virginia Commonwealth University Health Massey Cancer Center, USA; ⁵University of Virginia, USA; ⁶Chinese Academy of Medical Sciences and Peking Union Medical College, China; ⁷Saitama Medical University, Japan; ⁸Universidade Federal de Minas Gerais, Brazil; ⁹Department of Obstetrics and Gynecology, General Faculty Hospital in Prague, First Faculty of Medicine, Charles University, Czech Republic; ¹⁰Copenhagen University Hospital, Denmark; ¹¹Institut de Cancerologie des Hospices Civils de Lyon (IC-HCL), CITOH, Université Claude Bernard Lyon 1, Univ Lyon, France; ¹²Vall d'Hebron University Hospital, Vall d'Hebron Institute of Oncology (VHIO), Spain; ¹³Universitair Ziekenhuis Leuven, Belgium; ¹⁴American Hospital, Turkey; ¹⁵Charité-Universitätsmedizin Berlin, Germany; ¹⁶Sheba Medical Center, Israel; ¹⁷Medizinische Universität Innsbruck, Austria; ¹⁸Merck and Co., Inc., USA; ¹⁹Istituto Nazionale Tumori IRCCS Fondazione G Pascale, Italy

10.1136/ijgc-2020-IGCS.142

Background High-risk locally advanced cervical cancer (CC) has a poor prognosis, and >50% of patients recur in 2 years. Concurrent chemoradiotherapy (CRT) may enhance the immunostimulatory activity of the PD-1 inhibitor pembrolizumab. After KEYNOTE-158, in which pembrolizumab demonstrated durable antitumor activity, pembrolizumab monotherapy was approved for patients with PD-L1-positive recurrent or metastatic CC who progressed during or after chemotherapy. ENGOT-cx11/KEYNOTE-A18 (NCT04221945) is a phase 3, randomized, placebo-controlled study evaluating pembrolizumab with concurrent CRT in locally advanced CC.

Trial design Approximately 980 patients with high-risk, locally advanced, histologically confirmed CC who have not received systemic therapy, immunotherapy, definitive surgery, or radiation will be randomized 1:1 to receive 5 cycles of pembrolizumab 200 mg Q3W + CRT (5 cycles [with optional 6th cycle] of cisplatin 40 mg/m² Q1W + external beam radiotherapy [EBRT] followed by brachytherapy) followed by 15 cycles of pembrolizumab 400 mg Q6W or 5 cycles of placebo Q3W + CRT followed by 15 cycles of placebo Q6W. Randomization is stratified by planned EBRT type, cancer stage at screening, and planned total radiotherapy dose. Treatment will continue until patient receives ≤ 20 cycles of pembrolizumab (5 cycles 200 mg Q3W, 15 cycles 400 mg Q6W) vs placebo (~2 years) or until disease progression, unacceptable toxicity, or withdrawal. Primary endpoints are PFS per RECIST v1.1 by blinded independent central review and OS. Secondary endpoints include PFS at 2 years, OS at 3 years, complete response at 12 weeks, ORR, PFS and OS in PD-L1-positive patients, and safety. Enrollment is ongoing.