rearranging the position of the ports and bringing them closer to the tip of the femoral triangle. A 30 degree telescope helped in visualising the tip of the triangle better to remove the nodal tissue enbloc at completion of surgery.

Results Improved techniques led to easy identification of sartorius and standardization of the procedure.

Conclusion Issues and tips for improvement in surgical techniques especially in novel areas like robotic Inguino-femoral node dissection surgery are addressed.

Disclosures This surgical video was presented at IGCS Conference 2019.

453

INTRA-TUMOURAL ELECTRO-CHEMOTHERAPY (ECT) WITH BLEOMYCIN FOR PALLIATION OF CUTANEOUS RECURRENCE IN GYNAECOLOGICAL MALIGNANCY

¹Stuart Rundle, ¹Christine Ang, ²Tani Fasih, ¹Porfyrios Korompelis, ¹Nithya Ratnavelu. ¹Northern Gynaecological Oncology Centre; Queen Elizabeth Hospital; ²Queen Elizabeth Hospital; Breast and General Surgery

10.1136/ijgc-2020-ESGO.183

Introduction/Background ECT utilises pulsed electrical current to transiently increase cell membrane permeability to the cytotoxic agent, bleomycin. We present the use of ECT in patients with recurrent gynaecological malignancy previously treated with a combination of surgery or radiotherapy, experiencing symptoms from cutaneous lesions. This report presents evidence of the role for ECT in second and third line treatment.

Methodology Between July 2017 and August 2019, 6 patients with cutaneous recurrence of gynaecological malignancy (5 vulvar SCC and 1 high grade serous ovarian cancer) were treated with intra-tumoural bleomycin (9000 iu or 15000 iu, dependent on tumour volume) and pulsed-probe electroporation. Response was assessed clinically in routine follow up or following self-referral with return of symptoms. Post procedure pain scores were collated as part the quality of life evaluation.

Results Median treated tumour diameter was 6 cm (range 2 – 12 cm). Pain scores peaked between day 2 and day 7 post-procedure. The median progression free interval was 3.6 months (range 0.8 – 6.7 months).

Following ECT treatment 2 patients continued to receive supportive care. Two patients underwent repeat treatment with ECT and reported symptom improvement with each treatment. Due to further progression two patients underwent radical surgery and one patient received palliative chemotherapy.

Conclusion ECT should be considered for patients with symptomatic cutaneous recurrence of gynaecological malignancy who have previously had multi-modal treatment. It can achieve symptom control and reduce the need for radical surgery in this palliative setting.

Disclosures The Authors have no conflicts of interest to disclose.

469

NEW VARIANT OF RECONSTRUCTIVE SURGERY FOR ADVANCED VULVAR CANCER TREATMENT

¹Olha Bublikova, ²Yevgeniy Kostyuchenko, ¹Valentyn Svintsitskiy, ³Igor Motuziuk, ¹Nataly Tsip, ¹Sergiy Nespryadko, ¹Alena Samokhvalova, ³Oleg Sydoruk. ¹National Cancer Institute; ²National Cancer Institute; O.O. Bogomolets National Medical University; ³O.O. Bogomolets National Medical University

10.1136/ijgc-2020-ESGO.184

Introduction/Background Surgical treatment of advanced vulvar malignant tumors usually requires immediate reconstruction. Large defects after pelvis, vagina, vulva, groin and perineum wide excision require closure with the usage of difficult reconstructive techniques. In this case the most suitable myocutaneous flap for reconstruction is rectus abdominis muscle flap, which provides the biggest volume of tissues to cover those large defects.

Methodology Woman 67 y.o. initially presented with the combined treatment of cervical cancer stage IIB. Within 5 month was diagnosed the lymphedema in the left lower extremity. Approximately 2 years she presented the vulvar tumor measured 15 × 10 cm. A biopsy was performed the lymphangiosarcoma.

Results In the National Cancer Institute of Ukraine we investigated a new variation of large defects reconstruction using rectus abdominis muscle flap. To collect a donor flap, we perform 3 arcuate incisions: one vertical by the medial line around the umbilicus, and two oblique incisions towards upper anterior iliac spine of one of the sides. In this way we use only one half of the abdomen, and in case of any complications with the flap or relapse of the disease we will have the second donor site for possible future re-operation. There is a narrow 'bridge' of tissues we leave between the excised donor and recipient sites. It is extremely important to preserve blood supply not only to the flap, but also to the 'bridge' to avoid complications. We perform a tunnel under the 'bridge' as small as possible to preserve all inferior epigastric vessels, but enough to transfer the flap and not to squeeze the pedicle. After the surgery we have a half-inverted Y-shaped scar on of the sides of the abdomen, an inverted triangular scar at the pelvis area, circumumbilical scar and a short vertical scar on the flap to imitate pudendal cleft with central structures of vulva (urethra and vaginal tube).

Conclusion We consider our variation of this type of surgery the most safe and efficient, with the opportunity of re-operation if needed. Advanced vulvar malignancies are quite rare, so we will keep working on development and enhancement of the technique to help these patients.

Disclosures Authors declare no disclosures.

474

VULVAR RECONSTRUCTION IN PATIENTS WITH ADVANCED MALIGNANCIES

¹Olha Bublikova, ²Yevgeniy Kostyuchenko, ³Igor Motuziuk, ¹Valentyn Svintsitskiy, ¹Sergiy Nespryadko, ¹Nataly Tsip, ³Oleg Sydoruk, ¹Alena Samokhvalova. ¹National Cancer Institute; ²National Cancer Institute; O.O. Bogomolets National Medical University; ³O.O. Bogomolets National Medical University

10.1136/ijgc-2020-ESGO.185

Introduction/Background Vulvar malignant tumors are the rarest oncological disease of women's reproductive system in Ukraine. In 2018 there were registered only 477 new cases of vulvar cancer among all Ukrainian women. It is approximately three cases per 100 thousand among female population. However there are less common histological types, such as leiomyosarcoma, melanoma and lymphangiosarcoma, which make up approximately 1% of all cases. The main goal in surgical treatment is to obtain gross surgical margins, which sometimes require the usage of reconstructive techniques.

Methodology