Conclusion With the overall detection rate of 87.2% (63.5% bilateral and 24.0% - unilateral) we find SLB ICG procedure feasible. Most frequently SLs were detected in the external iliac region. Age and extragenital pathology are the statistically significant factors associated with the failure of SLB procedure.

309

ROLE OF THREE-DIMENSIONAL TRANSVAGINAL ULTRASOUND AND DIFFUSION-WEIGHTED MAGNETIC RESONANCE IMAGING FOR ASSESSMENT OF MYOMETRIAL INVASION IN PATIENTS WITH LOW-RISK ENDOMETRIAL CANCER

¹Núria Carreras Diéguez, ¹Isabel Matas, ²Cristian de Guirior, ²Meritxell Munmany, ¹Pere Fusté, ¹Núria Agustí, ¹Ariel Glickman, ¹Berta Diaz-Feijoo, ¹Jaume Pahisa, ¹Aureli Torne. ¹Hospital Clínic de Barcelona; Gynecologic Oncology Unit; ²Hospital Clínic de Barcelona; Gynecology and Obstetrics

10.1136/ijgc-2020-ESGO.67

Introduction/Background In patients with early-stage, grade 1–2, endometrioid endometrial cancer, preoperative assessment of myometrial invasion is essential to define the need of pelvic and paraaortic lymph node dissection. Our aim was to evaluate the role of three-dimensional transvaginal ultrasound (3D-TVUS) and diffusion-weighted magnetic resonance imaging (DW-RMI) for the assessment of myometrial invasion in patients with low-risk endometrial cancer.

Methodology We performed a single center retrospective study, including patients who underwent surgery for grade 1–2 endometrioid endometrial cancer, FIGO stage I-II, in Hospital Clínic de Barcelona between 2010 and 2019. We computed sensitivity, specificity, and predictive values of 3D-TVUS and DW-RMI, as well as of intraoperative frozen section pathological study of surgical specimen, for diagnosis of deep myometrial invasion ($\geq 50\%$). Definitive pathological analysis of surgical specimen was considered gold standard for diagnosis of deep myometrial invasion.

Results One hundred and fifty-three patients were included, 120 (78.43%) patients presented myometrial invasion $< 50\%$ in postoperative analysis of surgical specimen and 33 (21.57%) patients presented deep myometrial invasion. Sensitivity and specificity of 3D-TVUS for diagnosis of deep myometrial invasion was 68.8% and 80.5% respectively, while DW-RMI showed a sensitivity and specificity of 76.2% and 84.4%. When combining both techniques (we considered that a patient had deep myometrial invasion when 3D-TVUS or DW-RMI – or both of them – showed deep myometrial invasion), sensitivity was 93.1% and specificity was 68.4%. The proportion of patients with uterine fibroids was higher in the group of patients with false negative (60%) or false positive (39.13%) result in 3D-TVUS, although these results did not reach statistical significance. Regarding the intraoperative frozen section pathological study of surgical specimen, it showed a sensitivity of 75% with specificity of 96.4% for diagnosis of deep myometrial invasion.

Conclusion The combination of 3D-TVUS and DW-RMI offers a better sensitivity, higher than intraoperative frozen section pathological study of the surgical specimen, for the diagnosis of deep myometrial invasion in patients with early-stage, grade 1–2, endometrioid endometrial cancer. Such information may be useful in selecting patients who require lymph node dissection.

Disclosures The authors of this abstract have no disclosures.

345

CLINICAL PARAMETERS PREDICTING RISK OF CONCURRENT INVASIVE CARCINOMA AND HIGH-RISK CARCINOMA IN PATIENTS WITH ENDOMETRIAL INTRAEPITHELIAL NEOPLASIA

Nazlı Orhan¹, Raziye Melike Yıldırım¹, Halise Meltem Batur¹, Utku Akgor², Nejat Ozgul¹, Murat Gultekin¹, Mehmet Coskun Salman¹. ¹Hacettepe University Faculty of Medicine; Department of Obstetrics and Gynaecology; ²Hacettepe University Faculty of Medicine; Department of Gynecological Oncology; Department of Obstetrics and Gynaecology

10.1136/ijgc-2020-ESGO.68

Introduction/Background Endometrial intraepithelial neoplasia (EIN) is a premalignant lesion, but risk of concurrent endometrial adenocarcinoma (EAC) is also high. Although most patients with EIN diagnosed with concurrent EAC will have low risk disease, some will have high-risk disease who require. Clinical characteristics may help determine such patients.

Methodology Patients with a diagnosis of EIN who were operated at Hacettepe University Faculty of Medicine, Department of Obstetrics and Gynaecology were identified. The rate of concurrent EAC and high-risk EAC were determined. Preoperative characteristics were reviewed in order to determine the predictors of concurrent malignancy.

Results A total of 252 patients constituted study group. Mean age was 46.6 years and 43.7% were postmenopausal. 44.0% had co-existing one or more medical diseases while 14.3% had diabetes, 19.5% had hypertension, and 7.5% had both. The most common surgery was total hysterectomy with or without adnexal removal performed in 93.6% of patients. Frozen section was requested for 82.5% of patients. Final pathology revealed EAC in 17.5%, but only 4.4% had high-risk disease. The accuracy of frozen section for predicting final pathology in terms of the presence or absence of EAC was 89.4%. Patient with malignancy tend to be significantly older (47.4 vs. 54.1 years, $p=0.02$) and risk of malignancy was significantly higher in postmenopausal women (9.2% vs 28.2%, $p<0.001$) and in women with hypertension (13.8% vs 32.7%, $p=0.02$). similarly, patients with high-risk disease were significantly older (48.2 vs. 58.2 years, $p=0.01$) and this risk was higher in postmenopausal women (1.4% vs. 8.2%, $p=0.01$) and women with hypertension (3.0% vs. 10.2%, $p=0.04$).

Conclusion Surgery is the mainstay of treatment in patients with EIN. During surgery, frozen section evaluation should be requested since a significant proportion of patients have concurrent EAC and frozen section is highly effective in determining these patients. Although rare, some patients may have concurrent high-risk endometrial carcinoma necessitating surgical staging. Both concurrent invasive carcinoma and high-risk disease are associated with older age, being in postmenopausal period, and having hypertension.

Disclosures No potential conflict of interest to declare.

357

ENDOMETRIAL BIOPSIES: FOR WHOM AND WHEN?

¹Bilal Esat Temiz, ¹Esra Karatas, ²Utku Akgor, ³Murat Gultekin, ³Mehmet Coskun Salman, ¹Nejat Ozgul. ¹Hacettepe University Hospital; Obstetrics and Gynecology; ²Hacettepe University Faculty of Medicine; Department of Gynecological Oncology; Department of Obstetrics and Gynaecology; ³Hacettepe University Faculty of Medicine; Department of Obstetrics and Gynaecology

10.1136/ijgc-2020-ESGO.69