a) Injection into the anterior myometrial wall. b) Injection into the posterior myometrial wall. Arrow: needle; Triangle: Radiotracer cluster.

Abstract 443 Figure 1 Radiotracer injection using the TUMIR technique

Planar lymphoscintigraphy at 2h post injection of the radiotracer (a). Volumetric reconstruction of right hemipelvis SPECT / CT data showing pelvic and paraaortic drainage (b) Coronal section of fused SPECT/CT image demonstrating bilateral pelvic drainage (c)

Abstract 443 Figure 2 Planar lymphoscintigraphy and presurgical SPECT/CT

of the patients had positive SLN (1.5% exclusively para-aortic). The percentage of false negative (FN) was 12.5%.

The intraoperative SLN detection in the RT-ICG group was 68% (56% of pelvic drainage, 33% of paraortic drainage, without cases with exclusively paraortic drainage). A bilateral drainage was found in 56% of the cases. The 11.1% of the patients had a positive SLN without FN cases.

Conclusion Detection of the SLN with RT is slightly higher than with hybrid tracer. The hybrid tracer obtains a higher percentage of SLN with bilateral pelvic drainage than RT and reduces the number of contralateral pelvic lymphadenectomies. The TUMIR technique allows detection of para-aortic SNs in more than 30% of patients, much higher than that obtained with other techniques.

Disclosures No disclosures.

483

SHOULD WE RE-STAGE AFTER POSITIVE SENTINEL NODE BIOPSY ON ENDOMETRIAL CANCER?

¹Mikel Gorostidi, ²Rubén Ruiz Sautua, ²Ibon Jaunarena, ²Paloma Cobas, ²Arantxa Lekuona. ¹Osakidetza, Hospital Universitario Donostia; Gyn Onc; ²Osakidetza, Hospital Universitario Donostia

10.1136/ijgc-2020-ESGO.75

Introduction/Background Approximately 10% of patients with intraoperative diagnosis of low risk Endometrial Cancer (EC) will suffer an upstage after the definitive histological evaluation of the piece of hysterectomy and bilateral adnexectomy. We aim to explore the results associated with the performance of pelvic and para-aortic lymphadenectomy as restaging these patients that will require a second surgery, and to compare

those with and without Sentinel Node Biopsy (SNB) in the first procedure.

Methodology Retrospective cohort study involving 27 patients diagnosed with low-risk EC (ESMO-ESGO-ESTRO criteria) with surgical restaging due to upstage in the final histological result at the Hospital Universitario Donostia from April 2013 to September 2018. Surgical and oncological results were compared between patients who underwent hysterectomy and double adnexectomy without any additional procedure (SNB-) n=17 and those who also had a pelvic and aortic SNB (SNB+) n=10. The main outcome evaluated in the study was intraoperative complications. Secondary outcomes were mean operative time, length of hospital stay, number of nodes obtained, Progression-Free Survival (PFS) and Overall Survival (OS).

Results The median duration of restaging surgery was 240 minutes (Q25 - Q75: 180 - 300) in the SNB(-) group, and 300 (Q25 - Q75: 247.5 - 330) minutes in the SNB(+) group, this difference being statistically significant (one-side t-student test, p=0.0295). With regard to intraoperative complications, there were 17.65% vs 40% respectively, all of them vascular, this difference being not significant. There were no statistical differences in length of hospital stay and number of pelvic nodes obtained. PFS and OS in both groups were the same.

Conclusion Women with EC who require lymph node restaging due to upstage and have previously undergone sentinel lymph node biopsy have greater surgical difficulty with longer duration of the procedure. The risk of complications is increased. We advise against performing a second re-staging surgery in patients sentinel node biopsy.

Disclosures No disclosures.

491

HAVE WE IMPROVED UTERINE SEROUS CARCINOMAS MANAGEMENT THE LAST TWO DECADES? A SINGLE CENTER EXPERIENCE

¹Michail Lontos, ²Anna Svana, ²Roubini Zakopoulou, ²Flora Zagouri, ³Charalampos Theofanakis, ³Nikolaos Thomakos, ²Dimitrios Haidopoulos, ⁴Alexandros Rodolakis, ⁵Maria Sotiropoulou, ⁶Meletios-Athanasios Dimopoulos, ⁶Efi Skafida, ⁶Maria Kaparelou, ⁶Konstantinos Koutsoukos. ¹General Hospital of Athens 'Alexandras'; ²National and Kapodistrian University of Athens; Department of Clinical Therapeutics; ³University of Athens; Alexandra Hospital; 1st Department of Obstetrics and Gynecology; ⁴National and Kapodistrian University of Athens; 1st Department of Obstetrics and Gynecology; ⁵Department of Pathology, Alexandra Hospital; ⁶Department of Clinical Therapeutics, National and Kapodistrian University of Athens, Alexandra Hospital

10.1136/ijgc-2020-ESGO.76

Introduction/Background Uterine Serous Carcinomas (USC) account for ten percent of endometrial cancers. Management of the disease is challenging due to its aggressiveness. Despite numerous clinical trials and advances in the molecular pathology of the disease it remains the most lethal uterine carcinoma. We undertook this study to determine changes in the management of the disease and the survival of our patients the last two decades.

Methodology Medical records of women with USC treated at Alexandra Hospital from 2000 to 2018 were retrospectively identified. Clinicopathological data, treatment and survival data were analyzed. Kaplan-Meier Survival curves were generated using Strata SE 15.1; survival differences were estimated using the long-rank test.

Results 121 patients with USC and complete clinical data were identified. Median age was 66.9 years. Pure serous carcinomas were diagnosed in 66 patients, while the remaining had mixed histologies. At diagnosis, 33 patients (28.0%) had stage Ia disease, but also 28 patients had stage IV disease (23.7%). 115 patients were treated with surgery including lymphadenectomy in 63 cases (55.8%) and omentectomy in 83 cases (73.5%). 64 patients (52.9%) were treated from 1999 to 2009. There was no statistical difference in the stage distribution, omission of lymphadenectomy or omentectomy and type of adjuvant treatment among patients treated prior or after 2010. Also, for patients with stage I-III disease, eligible for adjuvant treatment, both disease free survival and overall survival (OS) per stage did not differ significantly between the two decades. Finally, in patients with de novo stage IV disease there was no improvement in either first line Progression Free Survival or OS between the two decades (mPFS 1st vs 2nd decade: 8.6 vs 5.9 months 95% CI [1.5-15.7].[3.4-8.5] and mOS 18.4 vs 11.6 months 95% CI [1.7-34.9].[0.0-25.1].

Conclusion Despite recent clinical trials, surgical and medical management of USCs had not changed significantly the last two decades in everyday clinical practice. This is reflected in the survival of the patients.

Disclosures Authors have no conflict of interest.

492

ENDOMETRIAL CANCER: THE ROLE OF PROGNOSTIC FACTORS AND THEIR IMPACT ON RECURRENCE PATTERN

¹Giulia Parpinel, ²Luca Fuso, ¹Maria Elena Laudani, ¹Annalisa Carapezzi, ²Enrico Badellino, ²Nicoletta Petronio, ³Marisa Ribotta, ¹Martina Barboni, ²Annamaria Ferrero, ¹Paolo Zola. ¹Sant'anna Hospital; Department of Surgical Sciences; ²Mauriziano Hospital; Department of Surgical Sciences; ³Aou Città Della Salute e Della Scienza; Department of Pathological Anatomy

10.1136/ijgc-2020-ESGO.77

Introduction/Background The rate of recurrence of endometrial cancer is 11-19%. It is related to different prognostic factors which define specific risk classes in order to decide for an adjuvant treatment. The objective of this study is to evaluate how prognostic factors influence the probability and pattern of recurrence.

Methodology This multicentric observational retrospective study was conducted on 552 patients treated for endometrial cancer between February 2011-2019. The considered parameters were: age, BMI (Body Mass Index), surgery, stage, LVSI (Lymphovascular Space Involvement), myometrial infiltration, histological grade, lymph node involvement, adjuvant therapy, relapse. DFS (Disease Free Survival) and OS (Overall Survival) were stratified by the presence or absence of prognostic factors.

Results The rate of recurrence was 14,7%. Median time to recurrence was 15 months and 5-years OS was 84%. We observed an increase from 6 to 30% in the distance metastasis rate associated with positive LVSI. Monovariate analysis showed a correlation between DFS and advanced stage (O.R. II 6.7; III-IV 9.2 p=0,0001), positive lymph nodes (O.R. 3.7; p=0,04), myometrial infiltration (O.R. 3.9 p=0,0001) and LVSI (O.R. 3.5 p=0,005). Similar results were observed considering OS and grading (p=0,002). We conducted a Cox