395

MENOPAUSAL SYMPTOMS AND SEXUAL DISORDERS IN EPITHELIAL OVARIAN CANCER SURVIVORS, A GINECO VIVROVAIRE2 STUDY

¹Florence Joly, ²Anne Gompel, ³Raffaële Fauvet, ⁴Idir Licaj, ⁵Djihane Ahmed-Lecheheb, ⁶Jean-Michel Grellard, ⁶Elsa Kalbacher, ⁷Olivier Tredan, ¹François Garnier, ⁸Patricia Pautier. ¹Centre François Baclesse; Unité Inserm 1086; ²Chu Paris Centre – Hôpital Cochin, Site Cochin; ³Chu Caen Département OF Gynecology AND Obstetrics; ⁴Centre François Baclesse; ⁵Unité Inserm 1086 Anticipo; ⁶Chu Jean Minjoz; ⁷Centre Leon Berard; ⁸Gustave Roussy

10.1136/ijgc-2020-ESGO.167

Introduction/Background We have previously shown that Epithelial Ovarian Cancer (EOC) and its treatments have significant negative effects on Quality of Life (QoL) and long term fatigue. The aim of the present multicentric VIVROVAIRE2 study was to report the main menopausal symptoms of Epithelial Ovarian Cancer survivors (EOCS).

Methodology One hundred sixty-six patients of the 322 EOCS without relapse ≥ 3 years after first line of treatment accepted to participate to a gynecological consultation carried out by a gynecologist, including a questionnaire related to menopausal symptoms (MS), sexuality, clinical examination, and osteodensitometry. MS (hot flashes and/or night sweats) were described according to natural menopause (NM) or surgically induced menopause (SIM). QoL, fatigue, insomnia and mood disorders were measured with the questionnaires (FACT-G, FACIT Fatigue, ISI, and HADS).

Results Median age was 62 years [20–83], FIGO stage III/IV (48%) and $< 10\%$ BRCA1&2 mutated. Histological subtypes were: high grade serous 28%, low grade serous 22%, endometrioid G2-3 (15%) endometrioid G1 (3%), clear-cell 21%, mucinous 5%. All EOCS had surgery, 97% of patients received platinum and taxane chemotherapy, median delay from treatment was 5 years [3–24] and 59 (36%) had SIM. 14% of EOCS had osteoporosis. Half of patients reported MS either hot flashes (47%) or night sweats (32%). 72% with SIM had MS compared to 41% with NM ($p < .001$). MS were not associated with poor global QoL, fatigue, insomnia or mood disorders. At the gynecological consultation, two-thirds of EOCS reported a decrease in sexual desire notably EOCS with SIM, which showed a greater decreased libido than NM ($p < .02$).

One hundred seven patients have never been treated with Hormone Replacement Therapy (HRT) including 59 who reported MS and 48 who (40%) had SIM.

Among 85 EOCS with MS, 80 (94%) (38 SIM and 42 NM) did not benefit from HRT after cancer treatment; 76% presented no CI of HRT.

Conclusion Menopausal symptoms and sexual disorders are frequently reported by EOCS, particularly among surgically induced menopause patients. A majority of EOCS with MS may benefit from HRT to improve these symptoms.

Disclosures The authors declare that they have no conflict of interest in relation to the subject of the article.

476

EFFECTIVENESS OF PHOTODYNAMIC THERAPY IN LICHEN SCLEROSUS WITH NEOPLASTIC BACKGROUND

¹Magdalena Bizoń-Szpernalowska, ²Danuta Maśliriska, ³Włodzimierz Sawicki. ¹Chair and Department of Obstetrics, Gynecology and Gynecological Oncology; ²Medical University in Warsaw; ³Chair and Department of Obstetrics, Gynecology and Gynecological Oncology, Medical University of Warsaw

10.1136/ijgc-2020-ESGO.168

Introduction/Background The main problem connected with lichen sclerosus is inflammatory process and correlation of this vulvar dermatose with autoimmune and neoplastic diseases. Photodynamic therapy is one of the most popular worldwide discussed method of treatment of lichen sclerosus. Despite the short history, this method has a quiet good benefits.

The aim of the study was to assess the influence of neoplastic disease on effectiveness of photodynamic therapy used in lichen sclerosus treatment.

Methodology Analysis was based on 182 patients with age 31–88 from Outpatient Vulvar Clinic, which is a part of Department of Obstetrics, Gynecology and Gynecological Oncology. Material was divided into 3 groups: patients with neoplastic disease in history, women with positive familiar history of neoplastic disease and women neither medical history nor familiar history of neoplastic diseases.

Analysis was based on immunohistochemical reaction in samples of vulvar biopsy with receptors TLR, mast cells, Langerhans cells and MeCP2 - Methyl-CpG-binding protein 2, vulvoscopic assessment with special scales our own authorship before and 10 weeks after photodynamic therapy and questionnaire of patients' vulvar symptoms before and after the treatment.

Results TLR3, TLR9 and Langerhans cells take part in advanced stage of lichen sclerosus pathogenesis. MeCP2 occurs in every stage of lichen sclerosus. Localization depends on different stage of disease and can be in every layer of the skin. The role of this protein is to take part in epigenetics process.

In vulvoscopic assessment, the percentage of improvement after PDT was higher in group of women with no correlation to neoplastic diseases (55,3%) than in women with contact with neoplastic disease (47,7%).

According to subjective symptoms connected with lichen sclerosus the most frequent was itching, which stand for 76,2%; 71,7% and 66,9% in following groups. In comparison to other complains, the highest percentage of every group were patient with neoplastic disease in the past. Satisfaction with the level above 50% after photodynamic therapy referred 80,8% patients from all analysed population.

Conclusion To sum up, the presence of neoplastic disease can influence on tissue and clinical phenotype on effectiveness of photodynamic therapy used in treatment of lichen sclerosus.

Disclosures

493

HOW TO RECONSTRUCT AN OPEN ABDOMINAL WALL AFTER NECROTIZING FASCIITIS: SURGICAL MANAGEMENT IN DIFFICULT CASES

¹Seda Sahin Aker, ²Arda Özdemir, ¹Fırat Ortaç. ¹Ankara University Faculty of Medicine; Gynecologic Oncology; ²Ankara University Faculty of Medicine; Plastic and Reconstructive Surgery

10.1136/ijgc-2020-ESGO.169

Introduction/Background Necrotizing fasciitis (NF) is a rare but very fatal infection involving and causing necrosis of the subcutaneous tissue and fascia. The incidence of NF is 0.4/100000. NF has a high mortality rate so it is needed an early diagnosis and proper treatment. There are several risk factors of NF. NF presents as painful, patchy discoloration of the skin without margins and a black necrotic plaque at the wound area. Ischemia and tissue necrosis can develop and local anaesthesia can occur because of the nerve damage.

Methodology A 59-year-old Turkish woman was admitted to hospital with a complaint of a postmenopausal bleeding.

Patient underwent a probe curettage. The pathology report showed a grade 1 endometrioid type of endometrial carcinoma. A total abdominal hysterectomy and bilateral salpingo-oophorectomy were performed. Final pathology report revealed that stage 1A endometrioid type of endometrium carcinoma.

Results Third day after surgery patient had fever (38 °C), tachycardia (102 beat/min) swelling was spread to the upper abdominal wall skin, vaginal discharge. She underwent a Hartmann procedure, abscess debridement and end sigmoid colectomy procedure. Patient then underwent extensive surgical debridement after 48 hours and a vacuum sealing drainage dressing was placed to cover the open abdominal wall and a negative sucker was placed upon the anus for 5 days. The dressing was changed every 3 days. Cultures of the exudates from the wound grew *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*. Antibiotic treatment was adjusted according to the sensitivity results. After 21 days of a negative pressure wound treatment, the abdominal wall defect was 15*15 cm diameter and the wound covered with a granulation tissue. Patient underwent a split thickness skin graft operation. In this video which we want to demonstrate how to reconstruct an open abdominal wall defect with a full thickness skin graft. After removing the granulation tissue, a good vascular supported tissue had seen and the necrotic wound had removed by a curette. The split thickness skin graft had taken from left leg's superolateral healthy skin with a measure of 10*25 cm diameter and 3 mm thickness. The skin is meshed to cover the large wound area. The graft covered the whole open abdominal wall and stitched up with 4,0 polipropilen sutures.

Conclusion Necrotizing fasciitis is an uncommon condition and has serious morbidity-mortality rate. Surgical debridement is the cornerstone of the treatment. NGWT combined with a STSG can help to heal wounds with NF.

Disclosures Picture 1:72 hours after STSG surgery

Picture 3: 3 months after STSG surgery

Translational research

295

ACCUMULATION OF 53BP1 IN CIRCULATING TUMOR CELLS DURING TREATMENT WITH ERIBULIN IDENTIFIES CHEMOTHERAPY-RESPONSIVE METASTATIC BREAST CANCER PATIENTS

¹Fabienne Schochter, ¹Kim Werner, ²Cäcilia Köstler, ³Volkmar Müller, ⁴Hans Neubauer, ⁵Tanja Fehm, ¹Thomas WP Friedl, ²Bernhard Polzer, ¹Wolfgang Janni, ¹Lisa Wiesmüller. ¹Ulm University; Department of Obstetrics and Gynecology; ²Fraunhofer-Institute for Toxicology and Experimental Medicine; Division of Personalized Tumor Therapy; ³Universitätsklinikum Hamburg-Eppendorf; Klinik und Poliklinik für Gynäkologie; Department of Gynecology, University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany; ⁴Heinrich-Heine-University of Düsseldorf; Department of Obstetrics and Gynecology; Life Science Center; ⁵Universitätsklinikum Düsseldorf; Klinik für Frauenheilkunde und Geburtshilfe; Department of Obstetrics and Gynecology, University of Duesseldorf, 40225 Duesseldorf, Germany

10.1136/ijgc-2020-ESGO.170

Introduction/Background Evidence suggests that the DNA end-binding protein p53-binding protein 1 (53BP1) expression in breast cancer is associated with poor prognosis, especially in triple-negative breast cancer (TNBC). Circulating tumor cells (CTCs) provide accessible 'biopsy material' to track cell traits and functions and their alterations during treatment.

Methodology We prospectively monitored the 53BP1 status, as a parameter for intact DNA damage response, in CTCs from 63 metastatic breast cancer (MBC) patients with HER2- CTCs before, during, and at the end of chemotherapeutic treatment with Eribulin in the DETECT-IV trial. Nuclear 53BP1 staining and genomic integrity were evaluated by immunocytochemical and whole-genome-amplification-based polymerase chain reaction (PCR) analysis. We used mean 53BP1 scores in CTC samples as dividing criteria, i.e. compared patients with 53BP1 scores <50% and ≥50%. We analyzed PFS of the patients from these two groups using scores obtained with samples at different time points during the study.

Results We found a decline of mean CTC numbers from baseline to 12 weeks of treatment but a dramatic rise at the final visit due to disease progression in 10/13 of the cases (mean CTC-values at baseline: 18, 2nd visit: 2, final visit: 118). Comparative analysis of CTCs from patients with 15 triple-negative and 48 hormone receptor positive tumors revealed elevated 53BP1 levels in CTCs from patients with HR+ metastases, particularly following chemotherapeutic treatment. Kaplan-Meier analysis between nuclear 53BP1-positivity in CTCs and progression-free survival (PFS) revealed an increasing association during chemotherapy until last examination (p=0.065).

Conclusion Our data suggest that 53BP1 detection in CTCs could be a useful marker to capture dynamic changes of chemotherapeutic responsiveness in triple-negative and HR+ MBC.

Disclosures FSch received speaker honoraria and a travel grant from Roche, Novartis, Pfizer and Lilly.

VM speaker honoraria from Amgen, Astra Zeneca, Celgene, Daiichi-Sankyo, Eisai, Pfizer, MSD, Novartis, Roche, Teva, and consultancy honoraria from Genomic Health, Hexal, Roche, Pierre Fabre, Amgen, ClinSol, Novartis, MSD, Daiichi-Sankyo, Eisai, Lilly, Tesaro, and Nektar, as well as institutional research support from Novartis, Roche, Seattle Genetics, and Genentech. Otherwise, no potential conflicts of interests were disclosed by the authors.

444

GENOMIC AND FUNCTIONAL CHARACTERISATION OF INTRA-TUMOURAL HETEROGENEITY IN HIGH GRADE SEROUS OVARIAN CANCER

¹Paula Cunnea, ²Ed Curry, ³Elizabeth Christie, ²Katherine Nixon, ²Chun Hei Kwok, ²Jennifer Ploski, ²Ratri Wulandari, ³David Bowtell, ²Christina Fotopoulou. ¹Imperial College London; Division of Cancer, Department of Surgery and Cancer; ²Imperial College London; ³Peter MacCallum Cancer Centre

10.1136/ijgc-2020-ESGO.171

Introduction/Background High-grade serous ovarian cancer (HGSOC) is characterised by high degrees of genomic instability and heterogeneity, with most patients eventually acquiring resistance to platinum-based chemotherapy. Matching the best treatment options to patients remains problematic due to diverse platinum resistance mechanisms and limited effective predictive biomarkers. This study aims to understand the extent of intra-tumoural heterogeneity (ITH) in advanced stage HGSOC, at presentation and relapse, and to define the link between ITH at the genomic and phenotypic levels.

Methodology Patients (n=49) undergoing radical upfront-debulking for advanced HGSOC at Hammersmith Hospital, UK, underwent a tumour mapping of their tumour