

Supplementary Table 1. Relapse rate, site of first relapse, and death rate in patients in the NAT, RT, and CCRT groups.

Parameter	NAT group, n=85	RT group, n=283	CCRT group, n=493	<i>p</i> -value		
				NAT vs. RT	NAT vs. CCRT	RT vs. CCRT
Relapse	10 (11.8%)	41 (14.5%)	53 (10.8%)	0.52	0.78	0.13
Site of first relapse						
Pelvic	6 (7.1%)	15 (5.3%)	13 (2.6%)	0.09	0.04*	0.43
Distant	2 (2.4%)	24 (8.5%)	33 (6.7%)			
Combined	1 (1.2%)	1 (0.4%)	3 (0.6%)			
Unknown	1 (1.2%)	1 (0.4%)	4 (0.8%)			
Disease-specific Death	7 (8.2%)	34 (12.0%)	42 (8.5%)	0.33	0.93	0.12

NAT, no adjuvant treatment; RT, radiotherapy; CCRT, concurrent chemoradiotherapy

Supplementary Table 2. Adjuvant treatment, relapse rate, site of first relapse, and death rate in patients in different subgroups of Sedlis criteria.

Parameter	-LVSI, deep or middle 1/3, ≥4cm (n=461)	+LVSI, deep 1/3 (n=193)	+LVSI, middle 1/3, ≥2cm (n=199)	+LVSI, superficial 1/3, ≥5cm (n=8)
Adjuvant treatment				
NAT	55 (11.9%)	14 (7.3%)	14 (7.0%)	2 (25.0%)
RT	167 (36.2%)	55 (28.5%)	58 (29.1%)	3 (37.5%)
CCRT	239 (51.8%)	124 (64.2%)	127 (63.8%)	3 (37.5%)
Relapse	46 (10.0%)	32 (16.6%)	25 (12.6%)	1 (12.5%)
Site of first relapse				
Pelvic	17 (3.7%)	8 (4.1%)	8 (4.0%)	1 (12.5%)
Distant	24 (5.2%)	21 (10.9%)	14 (7.0%)	0
Combined	2 (0.4%)	1 (0.5%)	2 (1.0%)	0
Unknown	3 (0.7%)	2 (1.0%)	1 (0.5%)	0
Disease-specific death	32 (6.9%)	28 (14.5%)	22 (11.1%)	1 (12.5%)

LVSI, lymph vascular space invasion; NAT, no adjuvant treatment; RT, adjuvant radiotherapy; CCRT, adjuvant concurrent chemoradiotherapy