

Supplemental Table S1. Survival analysis by Cox regression (univariate analysis) of parameters that influence recurrence-free survival in FIGO IB endocervical adenocarcinoma substages

Variable	Stage IB1			Stage 1B2			Stage 1B3		
	Recurrence-free survival HR	95% CI	p	Recurrence-free survival HR	95% CI	p	Recurrence-free survival HR	95% CI	p
Age	1.03	0.37-2.81	0.95	1.48	0.70-3.13	0.30	1.57	0.60-4.14	0.35
Surgical treatment	1.34	0.62-2.93	0.45	2.18	0.38-12.65	0.38	4.58	0.00-999.99	0.65
HPV status	3.54	1.02-12.19	0.046	1.15	0.39-3.40	0.79	3.30	1.16-9.34	0.025
Silva pattern	1.59	0.79-3.19	0.18	1.63	0.80-3.32	0.17	1.07	0.31-3.72	0.90
Histologic grade	1.36	0.67-2.74	0.38	2.02	1.12-3.66	0.019	1.28	0.56-2.89	0.54
Lymph-vascular invasion	4.48	1.78-11.23	0.001	5.39	1.85-15.65	0.002	2.09	0.47-9.18	0.32
Precursor lesions	2.71	1.07-6.84	0.035	1.53	0.67-3.48	0.30	1.62	0.49-5.26	0.42
Lymph node metastases	4.60	1.48-14.29	0.008	4.24	1.83-9.84	0.001	5.13	1.76-14.97	0.003

FIGO, International Federation of Gynecology and Obstetrics; HPV, human papillomavirus