



Development of a surgical competency assessment tool for sentinel lymph node dissection by minimally invasive surgery for endometrial cancer

Kristen Moloney,¹ Monika Janda,² Michael Frumovitz ,³ Mario Leitao,⁴ Nadeem R Abu-Rustum ,⁴ Emma Rossi,⁵ James L Nicklin,^{6,7} Marie Plante,⁸ Fabrice R Lecuru,^{9,10} Alessandro Buda ,^{11,12} Andrea Mariani,¹³ Yee Leung,¹⁴ Sarah Elizabeth Ferguson,^{15,16} Rene Pareja ,^{17,18} Rainer Kimmig,¹⁹ Pearl Shuang Ye Tong,²⁰ Orla McNally,^{21,22} Naven Chetty,²³ Kaijiang Liu,²⁴ Ken Jaaback,²⁵ Julio Lau,^{26,27} Soon Yau Joseph Ng,²⁸ Henrik Falconer ,^{29,30} Jan Persson,^{31,32} Russell Land,^{1,7} Fabio Martinelli ,³³ Andrea Garrett,¹ Alon Altman,^{34,35} Adam Pendlebury,³⁶ David Cibula,^{37,38} Roberto Altamirano,^{39,40} Donal Brennan,^{41,42} Thomas Edward Ind ,^{43,44} Cornelis De Kroon,⁴⁵ Ka Yu Tse,⁴⁶ George Hanna,⁴⁷ Andreas Obermair^{48,49}

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ijgc-2020-002315>).

For numbered affiliations see end of article.

Correspondence to

Professor Andreas Obermair, Gynaecological Oncology, University of Queensland Queensland Centre for Gynaecological Cancer Research, Herston, QLD 4029, Australia; ao@surgicalperformance.com

GH and AO are joint senior authors.

Received 9 December 2020

Revised 2 February 2021

Accepted 4 February 2021

Published Online First

4 March 2021



► <http://dx.doi.org/10.1136/ijgc-2021-002607>



© IGCS and ESGO 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Moloney K, Janda M, Frumovitz M, et al. *Int J Gynecol Cancer* 2021;**31**:647–655.

HIGHLIGHTS

- There is an urgent need to standardize surgical technique in sentinel lymph node dissection for endometrial cancer, and to develop tools for assessment of surgical quality.
- A delphi study achieved consensus on the mandatory, optional and prohibited steps of a sentinel lymph node dissection in endometrial cancer.
- A validated competency assessment tool is now available for use in sentinel lymph node dissection surgical quality assurance.
- A consensus on the mandatory, optional and prohibited steps of a sentinel lymph node dissection in endometrial cancer has been developed and validated.
- A competency assessment tool is now available for use in surgical quality assurance.

ABSTRACT

Introduction Sentinel lymph node dissection is widely used in the staging of endometrial cancer. Variation in surgical techniques potentially impacts diagnostic accuracy and oncologic outcomes, and poses barriers to the comparison of outcomes across institutions or clinical trial sites. Standardization of surgical technique and surgical quality assessment tools are critical to the conduct of clinical trials. By identifying mandatory and prohibited steps of sentinel lymph node (SLN) dissection in endometrial cancer, the purpose of this study was to develop and validate a competency assessment tool for use in surgical quality assurance.

Methods A Delphi methodology was applied, included 35 expert gynecological oncology surgeons from 16 countries. Interviews identified key steps and tasks which were rated mandatory, optional, or prohibited using questionnaires. Using the surgical steps for which consensus was achieved, a competency assessment tool was developed and subjected to assessments of validity and reliability.

Results Seventy percent consensus agreement standardized the specific mandatory, optional, and prohibited steps of SLN dissection for endometrial cancer and informed the development of a competency assessment tool. Consensus agreement identified 21 mandatory and three prohibited steps to complete a SLN dissection. The competency assessment tool was used to rate surgical quality in three preselected videos,

demonstrating clear separation in the rating of the skill level displayed with mean skills summary scores differing significantly between the three videos (F score=89.4; $P<0.001$). Internal consistency of the items was high (Cronbach $\alpha=0.88$).

Conclusion Specific mandatory and prohibited steps of SLN dissection in endometrial cancer have been identified and validated based on consensus among a large number of international experts. A competency assessment tool is now available and can be used for surgeon selection in clinical trials and for ongoing, prospective quality assurance in routine clinical care.

INTRODUCTION

Surgical trials pose methodological challenges¹ because surgeon training, experience, and skills influence the delivery of surgical interventions, leading to a variability in health practices and outcomes.² Surgical quality assurance can aid adherence to pre-defined standards and outcome measures, and enable reliable comparison across multiple clinical trial sites.^{3–7}

Management guidelines for apparent uterine-confined disease prescribe total hysterectomy, bilateral salpingo-oophorectomy for removal of the primary tumor, and assessment of locoregional lymph nodes to establish the stage of disease ('staging').⁸

Original research

This information is prognostic and may guide postoperative treatment decisions.^{8–9} Historically, surgical staging entailed full or limited pelvic/paraortic lymph node dissection. This practice was informed by the results of observational, clinicopathologic studies^{10–11} then adopted by FIGO in 1988.¹² Subsequent prospective studies failed to demonstrate differences in survival outcomes.^{13–14} Contemporary surgical staging involves sentinel lymph node (SLN) dissection.¹⁵ According to the SLN concept, tumor cells metastasize to one or two lymph nodes first, before involving further lymph nodes.¹⁶ Presumed benefits of SLN dissection include increased surgical staging precision, while sparing removal of other regional lymph nodes.¹⁷ Sentinel nodes are examined histopathologically using immunohistochemical ultrastaging.¹⁸ SLN dissection obtains accurate information about lymph node status¹⁸ such that many clinicians now elect it in place of a full lymphadenectomy.¹⁹

With rapid and global adoption of SLN dissection¹⁹ comes variability of surgical technique. Local institutional guidelines have been developed to minimize variation in outcomes.²⁰ However, these algorithms are insufficient to facilitate harmonization of the detailed surgical technique across a group of surgeons. There remains a need to define the precise surgical steps required to accomplish satisfactory bilateral SLN dissection; assess a surgeon's proficiency before enrolment of patients into clinical trials; and assist with ongoing surgical quality assurance.²¹

The purpose of this study was to establish a consensus on the specific mandatory and prohibited steps of SLN dissection in endometrial cancer, as well as develop a competency assessment tool. This facilitates assessment of surgical quality in clinical trials aiding in both the selection of surgeons and prospective quality assurance.

METHODS

Study Participants

The study was approved by the institution's Human Research Ethics Committee and informed written and/or eConsent was obtained from all participants.

Participants were expert gynecological oncology surgeons from five continents currently performing SLN dissection, henceforth referred to as 'the group'. Experts were recruited using snowball sampling, that is, first contacting surgeons known to perform SLN dissection per scientific reports or presentations in peer-reviewed forums, and then asking these surgeons to nominate other experts. Participant characteristics were summarized using descriptive statistics.

Standardization of Sentinel Lymph Node Dissection

A four-round Delphi methodology was applied in order to achieve standardization of SLN dissection steps and tasks. Several rounds of questionnaires were sent out to experts with the responses then aggregated, de-identified, and shared with the group after each round. Experts adjusted their answers in subsequent rounds, based on their interpretation of the group response provided to them. Over multiple rounds of questionnaires, the Delphi method seeks to reach best response through consensus.²² Study data was collected and managed on a secure, web-based REDCap electronic database hosted at The University of Queensland.^{23–24}

Delphi Consensus Process and Hierarchical Task Analysis

Round One (Semi-Structured Interviews)

After providing written informed consent, interviewees described their opinion about the mandatory, optional, and unwarranted steps taken in performing SLN dissection for endometrial cancer.

The interviews were conducted individually and were audio recorded. Recordings were transcribed and thematically analyzed by two reviewers (KM, AO). Each reviewer independently identified important and recurring codes (eg, uterine manipulation, identifying anatomy, troubleshooting). Codes were then compared to confirm important themes. The reviewers jointly examined codes and themes and interpreted the data. Where discordance in coding was identified, themes were refined through discussions between the two reviewers. Interviews were conducted until saturation in variations of technique. Key steps and tasks of SLN dissection were identified by a process of hierarchical task analysis.

Rounds Two–Four (Consensus Process)

An initial questionnaire was devised including all of the variations identified in the interviews. Members of the group were invited to indicate their agreement or disagreement with variations. In accordance with other published work,²⁵ consensus agreement level was set at 70%.^{4–22} Variations where consensus was reached were iteratively moved into an operation guide: those with <70% agreement remained for a subsequent survey round.

In accordance with the journal's guidelines, we will provide our data for the reproducibility of this study in other centers if such is requested.

Operation Guide

A SLN dissection operation guide was created including the mandatory, optional, and prohibited/unwarranted steps that reached 70% agreement level.

Competency Assessment Tool

Development

The final competency assessment tool was limited to mandatory and prohibited steps in the intraoperative phase of SLN dissection. A score of one to four was allocated to each step – 'skillful', 'adequate', 'inconsistent', 'lacking/deficient': for troubleshooting steps 'not applicable' was also offered.

Content Validity

Three surgical videos were selected having been agreed by KM and AO to represent poor, inconsistent, or optimal technique of SLN dissection according to the ratings conferred by application of the competency assessment tool. The videos featured the 11 surgical steps of SLN dissection assessed by the competency assessment tool, but did not include tracer preparation and injection, surgical troubleshooting, or pathological assessment of tissues. Content validity was assessed by KM and AO who discussed each step of the competency assessment tool in detail, before watching those individual steps performed with various skill levels across the three surgical videos, confirming that competency assessment tool items adequately reflected the skill required.

Contrast Validity and Internal Reliability

Contrast validity²⁶ was assessed via invitation of the group members to use the competency assessment tool in rating the three

pre-selected videos, each representing distinct performance levels. Due to the occurrence of some cells with a cell size <5, Fisher's exact tests were performed to assess if the proportion of experts who rated each of the three videos as 'skillful', 'adequate', 'inconsistent', or 'lacking/deficient' differed according to the quality of the video. An average competency assessment tool score (possible range 11–44) was computed for each video. One-way ANOVA modeling determined if the overall competency assessment tool score assigned by the SLN dissection experts to each video differed significantly. The summary score was used to assess the internal consistency (Cronbach alpha) of the competency assessment tool.

RESULTS

Thirty-five international gynecological oncology surgeons and experts in SLN dissection from 16 countries participated. Some demographic data was not available for five participants, but 28 surgeons were above 40 years of age (80%) and 27 were male (77%) (Table 1).

Twenty-four surgeons had practiced gynecological oncology for more than 10 years (69%), and 21 had performed SLN dissection for more than 5 years (60%). Nineteen surgeons (54%) reported that their institution had an endometrial cancer SLN dissection standard protocol. Twenty-one surgeons (60%) performed more than 50 SLN dissections annually, excluding those performed for cancer of the vulva. Participating surgeons reported using between one and eight methods to learn SLN dissection, for example, being self-taught (46%), learning from research papers (43%) or being trained by a senior colleague (31%).

Standardization of Sentinel Node Dissection

Delphi Round One (Hierarchical Task Analysis)

Saturation in the variation of the SLN dissection technique was reached after 25 interviews. Analysis of transcripts allocated themes into four phases: preoperative (dye selection and preparation, injection); intraoperative (pelvic dissection, identification of key anatomical structures, definition and dissection of sentinel node, extraction of tissue); troubleshooting; and a postoperative (pathology) phase. Task variations were defined as management of specific surgical steps in different ways. In total, 107 task variations were identified across the interviews (Table 2).

Delphi Rounds Two–Three (Consensus Process)

The first survey (Delphi round two) featured 107 task variations and was completed by all 35 participants (Online supplemental table 1). The second survey (Delphi round three) was informed by the results of the first survey and 33 of 35 participants responded (Online supplemental table 2). Over rounds two and three, >70% consensus was achieved in 33 of the 107 (30.8%) task variations^{4,22} on mandatory, optional, and prohibited steps of SLN dissection. Of the variations that reached consensus, 21 were classified as mandatory, nine optional, and three prohibited. For example, in round two, 79% of participants agreed that 'a transperitoneal approach of injecting dye into the uterus' should be prohibited, while 75% of participants agreed that 'the internal iliac artery must be identified for sentinel node mapping' was mandatory. An operation guide consisting of the final list of steps for which consensus was obtained is provided in Table 3.

Table 1 Participating surgeons' demographic characteristics

Variables	n=35* (%)
Age (years)	
31–41	4 (11)
41–50	15 (43)
51–60	14 (40)
>61	1 (3)
Gender	
Female	8 (23)
Male	27 (77)
Continent	
Europe	10 (29)
North America	9 (26)
Australia	10 (29)
Asia	4 (11)
South America	2 (6)
Does your Institution have its own sentinel node mapping protocol in endometrial cancer?	
No	14 (40)
Yes	19 (54)
How did you learn to perform sentinel node biopsies?	
Self-taught	16 (46)
Learned from research papers	15 (43)
Standard operating procedures/ protocols	11 (31)
Taught by senior colleague/s	11 (31)
Videos	10 (29)
Trained by surgeon/s overseas	7 (20)
Formal course/s	4 (11)
Other	7 (20)
How many years gynec? (years)	
Less than 10	11 (31)
10–19	14 (40)
20–29	9 (26)
30 or more	1 (3)
How many years performing SLND†? (years)	
Less than 5	13 (37)
5–9	8 (23)
10 or more	11 (31)
Number of SLND†	
Less than 50	14 (40)
50–99	10 (29)
100 or more	6 (17)

*Not all 35 participants provided data on demographics.

†Sentinel lymph node dissection.

There was consensus that the tracer of choice must be indocyanine green (ICG) but adding other tracers is optional. There was consensus that ICG should be administered by superficial injection (1–2 mm) into the cervix. Superficial injection was defined by the group as sub-mucosal injection into the epithelium of the ectocervix.

Table 2 Hierarchical task analysis including task variations

Phase	Theme	Sub-themes	No. of sub tasks/task variations
Peri-operative	Tracer injection	Choice of tracing agent	34
		Site of injection	
		Tracer concentration	
		Total volume of injected	
		Injection technique	
	Uterine manipulation	Use of manipulator at all	9
	Timing of manipulator insertion		
	Type of uterine manipulator		
	Sequence of initial steps	Timing of entry	7
		Timing of adhesiolysis	
		Timing of staging inspection	
Operative	Preparation/opening spaces	Transperitoneal identification of channels' pelvic side wall spaces	10
	Identifying anatomy, lymphatic channels, and sentinel nodes	Anatomical structure Methods of locating nodes	24
	Excision and confirmation of mapped nodes	Defining the SLN Technique of nodal excision Mode of ex vivo SLN confirmation	7
	Specimen retrieval	Mode of containment	5
	Troubleshooting	Action plan for no nodes mapped	5
Post-operative	Specimen labeling	Anatomical site Laterality	4
	Pathology processing		2

(akin to intra-dermal injection techniques used in SLN mapping for vulvar cancer). Deep injection (1 cm) was considered optional only when combined with mandatory superficial injection. Deep injection alone was considered prohibited by consensus. There was no consensus about the dilution of ICG (between 0.5 mg/L and 1.5 mg/mL), the total volume injected, or timing of injection (before or after establishing a pneumoperitoneum). The use of a uterine manipulator was considered optional, but if used, it should be inserted after tracer injection. There was consensus that dividing the round ligament and the infundibulopelvic ligament can be performed either before or after SLN dissection. The pelvic structures and spaces that should be demonstrated for SLN dissection include external and internal iliac vessels, ureter, obliterated umbilical ligament, and the paravesical space. The direction of the SLN dissection was considered optional (starting close to the cervix or dissecting toward the cervix).

The group agreed that the sentinel node should be defined as the most proximal node irrespective of the nodal station in which the node is found. Eighteen participants felt that mapping of presacral nodes should be optional (56.3%). There was lack of consensus on a side-specific lymphadenectomy if no nodes are mapped on one side. Participants agreed that the sentinel node should be a single mapped node with or without its next station (second echelon node(s)). A majority of participants (59.4%) but less than required for consensus, agreed that not all second, third, and fourth echelon nodes should be removed. Greater than 70 percent of participants agreed that specimen extraction should be within a containment device. There was consensus that ex-vivo fluorescence should be used to prove the sentinel node; that labeling of the sentinel node

should be according to laterality and nodal station; and enhanced pathology techniques for ultrastaging of sentinel nodes should be used.

Contrast Validity and Internal Reliability

Twenty-seven (77.1%) Delphi participants were involved in rating the quality of surgery of the three preselected videos using the competency assessment tool (Figure 1). For each of the 10 initial surgical steps, there was clear separation in the rating of the skill level displayed between the three videos (Table 4).

For example, while 78% of experts rated the 'optimal technique video' as skillfully performing the dissection of the iliac vessels, only 19% and 0% of experts rated the 'inconsistent technique video' and 'poor technique video' as skillful (Fishers' exact test=56.0; $P<0.001$). For the last step ('completion of SLND in one hemipelvis before proceeding to the contralateral side'), 25 of the 27 group members rated this step as not applicable. Overall, the mean skills summary score differed significantly between the three videos from 35.6 (SD=4.7) for the 'optimal technique video', to 25.3 (SD=5.9) for the 'inconsistent technique video' and 17.7 (SD=4.1) for the 'poor technique video' (one-way ANOVA F score=89.4; $P<0.001$). Internal consistency of the items was high (Cronbach $\alpha=0.88$).

DISCUSSION

Summary of Main Results

We report the creation of a competency assessment tool, derived by consensus among a large number of international experts, outlining the mandatory, optional, and prohibited steps of a SLN dissection

Table 3 Operation guide

Surgical step	Descriptor	Consensus recommendation
Tracer	ICG	Mandatory
	Blue dye	Optional
	Radio-labeled technetium	Optional
Injection location	Ectocervix in two or four positions	Mandatory
Injection technique	Superficial injection into the ectocervix	Mandatory
	Transperitoneal injection into the uterus	Prohibited
	Hysteroscopic injection into the uterus	Prohibited
	Surgeon appreciation of resistance at tracer injection	Mandatory
Injection needle	Gauge between 20G and 25G	Mandatory
	Length sufficient to ensure easy and accurate access to the cervix	Mandatory
Uterine manipulator	If being used, insert uterine manipulator after tracer injection	Mandatory
White light inspection	Prior to SLN mapping, conduct an inspection of the pelvic areas	Mandatory
Round ligament & Infundibulopelvic ligament	Preserve	Optional
	Divide	Optional
External vessels	Identify the external iliac vessels	Mandatory
Internal iliac artery	Identify the internal iliac artery	Mandatory
Ureter	Identify the ureter	Mandatory
Obliterated umbilical ligament	Identify the obliterated umbilical ligament	Mandatory
Uterine artery	Identify the uterine artery (medial to the ureter)	Optional
Paravesical space	Open the paravesical space	Mandatory
Direction of dissection	Start sentinel lymph node mapping at the level of the uterine artery and continue dissection LATERALLY away from the uterus	Mandatory
	Start sentinel lymph node mapping at the level of the uterine artery and and continue MEDIALLY toward the uterus	Optional
	Start sentinel lymph node mapping at the level of the uterine artery and and continue toward the pre-sacral area	Optional
	Start sentinel lymph node mapping at the most highlighted node and dissect proximally (TOWARD cervix)	Optional
	Start sentinel lymph node mapping at the most highlighted node and dissect cephalad (AWAY from cervix)	Optional
Dissection technique	Use blunt or electrosurgical technique	Mandatory
	Avoid disrupting lymphatic channels during dissection	Mandatory
	Ensure isolation of node from local anatomy	Mandatory
Definition of the sentinel node	A sentinel node is defined as ...	Mandatory
	<ul style="list-style-type: none"> ▶ The most proximal node [<u>The most proximal node is defined as the node closest to the uterus, regardless of location.</u>], irrespective of the nodal station in which the node is found ▶ A single mapped node or a single node plus its next station echelon node(s). 	Mandatory
SLN dissection	SLN dissection should be completed in one hemi-pelvis before proceeding to the contralateral side	Mandatory
Troubleshooting	Troubleshooting when no nodes are mapping includes any one, or combination of, the following options:	Mandatory
	▶ Wait, undertake dissection on the contralateral side before returning to original side	
	▶ Extend retroperitoneal dissection to encompass common, pre-sacral and/or paraaortic areas	
	▶ Re-inject ICG	
	▶ Undertake a side-specific lymphadenectomy	
Specimen extraction	Removal of nodes without using a containment device	Prohibited
Proof of sentinel node	Use ex-vivo green fluorescence to prove the sentinel node	Mandatory
Specimen labeling	Label specimens according to laterality (right/left) AND nodal station (obturator/external iliac/internal iliac/presacral/common iliac/aortic/caval)	Mandatory
Ultrastaging	Use enhanced pathology techniques, such as immunohistochemistry, for ultrastaging of sentinel nodes	Mandatory

Final consensus on mandatory and prohibited steps of sentinel lymph node dissection (SLND) by minimally invasive surgery in endometrial cancer. ICG, indocyanine green.

Phase	Assessment	Surgeon Technique is...				
		Skillful	Adequate	Inconsistent	Lacking/Deficient	
Surgical Steps	Prior to SLN mapping, a full inspection of the pelvic areas with white light is performed to exclude extrauterine disease					
	Identification of external iliac vessels					
	Identification of internal iliac artery					
	Demonstrates dissection of ureter					
	Development of paravesical space					
	Identification of obliterated umbilical ligament					
	Dissection technique avoids disrupting lymphatic channels and isolates nodal tissue from the local anatomy					
Sentinel Node Mapping	Demonstrate use of ex-vivo green fluorescence to prove sentinel node					
	Demonstrate specimen extraction involving a containment device (endocatch bag, finger of sterile glove or cup forceps). Removal of nodes through port without protection is prohibited					
	SLN mapping is completed in one hemi-pelvis before proceeding to the contralateral side					
Phase	Assessment	Not Applicable	Skillful	Adequate	Inconsistent	Lacking/Deficient
Troubleshooting	Troubleshooting includes: Wait and undertake dissection on the contralateral side before returning to original side AND/OR Extend retroperitoneal dissection to encompass common, pre-sacral and/or paraaortic areas AND/OR Re-inject ICG AND/OR Undertake a side-specific lymphadenectomy					

Figure 1 SLND competency assessment tool.

procedure for endometrial cancer. The competency assessment tool is validated by gynecological oncology surgeons and can be used by trial governance committees as a decision aide for surgeon selection and for ongoing quality assurance in surgical clinical trials.

Results in the Context of Published Literature

While local health service protocols²⁰ suggest specific steps for a SLN dissection, the present publication summarizes an operating consensus based on the opinion of a considerable number of international experts in SLN dissection. Consensus was achieved about definition of the sentinel node (the node closest to the uterus) regardless of whether it is located at the lateral pelvic wall, the

aortic/caval, or the presacral area. There was also agreement that the number of sentinel nodes removed should be kept to a minimum. There was no consensus on the mandatory need for completion lymphadenectomy on the ipsilateral side of a pelvis that fails to map. This most likely reflects the possibility of patient and uterine factors indicating against full lymph node dissection. Greater than 70 percent consensus was reached on the need to extract nodes through a containment system, the need for ex-vivo green fluorescence to prove the sentinel node, on specimen labeling, and on pathologic ultrastaging.

The competence assessment tool development undertaken in this study follows similar efforts in other surgical specialties. In general surgery, a recent systematic review reporting on quality assurance in randomized controlled trials of laparoscopic colorectal surgery identified three distinct categories of surgical quality assurance measures: trial entry criteria for surgeons and centers; standardization of surgical techniques; and continuous monitoring of surgeons and/or units.²⁵ A competence assessment tool was developed, validated, and implemented to assess technical surgical performance in the context of a summative assessment process for the National Training Program in Laparoscopic Colorectal Surgery (Lapco).²⁷ Subsequently, COLOR III⁴ investigators have described the standardization of surgical interventions followed by the development and assessment of objective surgical quality assurance tools for use in colorectal trials.

Strengths and Weaknesses

The strengths of our study include the large number of international experts who identified the mandatory, optional, and prohibited steps of SLN dissection based on consensus. In addition, the competence assessment tool was able to demonstrate contrast validity and internal reliability. Our expert participants reported a range of experience in SLN dissection despite recruitment using snowball sampling. We did not use more objective measurements of expertise such as a requirement for study participants to submit their own outcomes data, or videos of their individual technique. Another weakness of our study relates to the impact of our findings on meaningful clinical outcomes such as sensitivity for detection of metastases, or false negative rates. These parameters were not the focus of our study, but we acknowledge the absolute necessity for

Table 4 Assessment of contrast validity

	Poor video	Inconsistent video	Optimal video	Fishers' exact test
White light inspection	1 (4)	6 (22)	22 (81)	47.1; P<0.00
External vessels	0 (0)	5 (19)	21 (78)	56.0; P<0.001
Internal iliac artery	0 (0)	2 (7)	22 (82)	75.3; P<0.001
Ureter	0 (0)	6 (22)	20 (74)	70.6; P<0.001
Paravesical space	0 (0)	4 (15)	19 (70)	58.9; P<0.001
Obliterated umbilical ligament	0 (0)	2 (7.4)	19 (70)	60.3; P<0.001
Dissection technique	1 (4)	2 (7.4)	16 (59)	36.9; P<0.001
Proof of sentinel node	6 (22)	2 (7.4)	20 (74)	33.6; P<0.001
Specimen extraction	0 (0)	9 (33)	21 (78)	84.2; P<0.001
SLN mapping	1 (3.7)	5 (19)	10 (37)	15.9; P=0.03

n (%) of reviewers who rated the performance as skillful.
SLN, sentinel lymph node.

individual surgeons to monitor their performance including prospective audit of sensitivity, false negatives, recurrence patterns, and rates. This predictive clinical validity of the SLN dissection technique can only be determined with accumulation of clinical outcomes after using the competence assessment tool in SLN dissection clinical trials and educational programs, as has been demonstrated for a colorectal competence assessment tool in the Lapco program.²⁸

Implications for Practice and Future Research

Despite the benefits of SLN dissection, including shorter operating times compared with a lymphadenectomy, it remains unknown in which ways SLN dissection impacts clinically relevant outcomes, such as the need for postoperative radiation treatment or chemotherapy, recovery from surgery and quality of life, the incidence of adverse events, disease recurrence, and survival.²¹ Additionally, while new surgical procedures may appear promising, there remains a need to evaluate novel surgical procedures for safety and effectiveness.²⁹ Such surgical trials rely on the standardized delivery of the intervention (with minimal variation) to allow a meaningful and reliable comparison between intervention and control groups across multiple surgeons or trial sites. In the context of SLN dissection, variability in technique, and failure to identify sentinel nodes could translate into the need for frozen section assessment of the uterus, acceptance of unknown nodal status, or may increase the risk of an “empty package”,³⁰ all of which may confound the results of SLN dissection efficacy trials. Depending on local protocols, some patients may even require re-staging, a full ipsilateral lymphadenectomy,²⁰ or might warrant adjuvant chemotherapy or radiation treatment based on uterine risk factors. These scenarios may have significant impact on short- and long-term patient and trial outcomes. Using this competency assessment tool, institutions or clinical trialists can define quality standards of sentinel lymph node dissection and measure individual surgical performance.

Significant efforts are made by chief investigators and trial management committees to minimize the variability in surgical techniques and outcomes, including limiting the trial to sites with a high surgical volume. Recently, principal investigators completed a site visit and all surgeons were observed in-person^{18,31} or unedited videos were reviewed to confirm standardization of the technique.³² In other trials, participating surgeons were required to have completed a minimum number of procedures, before the initiation of enrollment.³³ While these measures were valuable within institutions, volume, minimal number, or observation of one surgery, they may be inaccurate without application of a standardized assessment tool.

Conclusion

The output from this work includes a list of mandatory and prohibited steps of a SLN dissection that independent assessors can use to check for both surgical proficiency as well as whether SLN dissection has been performed in accordance to an agreed standard. The work provides specific steps of SLN dissection, and the quality assurance criteria developed as part of this process will help with selection of prospective surgeons into surgical trials evaluating SLN dissection. The goal is to shorten the learning curve³⁴ but also to control for heterogeneity in surgical performance that could override the true efficacy.⁴

Author affiliations

- ¹Gynaecologic Oncology, Royal Brisbane and Women's Hospital, Herston, Queensland, Australia
- ²Centre for Health Services Research, The University of Queensland Faculty of Medicine, Brisbane, Queensland, Australia
- ³Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA
- ⁴Gynecology Service Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, New York, USA
- ⁵Obstetrics and Gynecology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA
- ⁶Gynaecological Oncology, Royal Brisbane and Women's Hospital, Herston, Queensland, Australia
- ⁷Faculty of Medicine, The University of Queensland, St Lucia, Queensland, Australia
- ⁸Gynecology Oncology Service, Centre Hospitalier Universitaire de Québec-Université Laval, Québec, Québec, Canada
- ⁹Surgical Oncology, Institute Curie, Paris, France
- ¹⁰Surgical Oncology Department for Breast and Gynecology, Université de Paris, Paris, Île-de-France, France
- ¹¹Department of Obstetrics and Gynecology, Università degli Studi Milano-Bicocca, San Gerardo Hospital, Monza, Italy
- ¹²Division of Gynecologic Oncology Italy, Ospedale Michele e Pietro Ferrero, Verduno (CN), Italy
- ¹³Gynecologic Surgery, Department of Obstetrics and Gynecology, Mayo Clinic Rochester, Rochester, Minnesota, USA
- ¹⁴Obstetrics and Gynaecology, The University of Western Australia Faculty of Health and Medical Sciences, Perth, Western Australia, Australia
- ¹⁵Gynecologic Oncology, University Health Network, Toronto, Ontario, Canada
- ¹⁶Obstetrics and Gynaecology, University of Toronto, Toronto, Ontario, Canada
- ¹⁷Gynecologic Oncology, Instituto Nacional de Cancerología, Bogotá, Colombia
- ¹⁸Gynecologic Oncology, Clínica De Oncología Astorga, Medellín, Colombia
- ¹⁹Gynecology and Obstetrics, University of Essen, Essen, Germany
- ²⁰Gynaecologic Oncology, National University Health System, Singapore
- ²¹Obstetrics and Gynaecology, Royal Women's Hospital, Parkville, Victoria, Australia
- ²²Victorian Comprehensive Cancer Centre, University of Melbourne, Parkville, Victoria, Australia
- ²³Gynaecologic Oncology, Mater Health Services Brisbane, South Brisbane, Queensland, Australia
- ²⁴Gynecology and Obstetrics, Shanghai Jiao Tong University School of Medicine Affiliated Renji Hospital, Shanghai, China
- ²⁵Gynaecologic Oncology, John Hunter Hospital, New Lambton Heights, New South Wales, Australia
- ²⁶Gynecology Oncology, Hospital General San Juan de Dios, Guatemala, Guatemala
- ²⁷Gynecology Oncology, University of San Carlos de Guatemala Faculty of Medical Sciences, Guatemala, Guatemala
- ²⁸Obstetrics and Gynaecology, National University of Singapore, Singapore
- ²⁹Women's and Children's Health, Karolinska Institute, Stockholm, Sweden
- ³⁰Women's and Children's Health, Karolinska University Hospital, Stockholm, Sweden
- ³¹Obstetrics and Gynaecology, Skanes Universitetssjukhus Lund, Lund, Skåne, Sweden
- ³²Clinical Sciences, Obstetrics and Gynaecology, Lund University Faculty of Medicine, Lund, Sweden
- ³³Gynaecologic Oncology, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy
- ³⁴Gynecologic Oncology, University of Manitoba, Winnipeg, Manitoba, Canada
- ³⁵Gynecologic Oncology, CancerCare Manitoba, Winnipeg, Manitoba, Canada
- ³⁶Gynaecological Oncology, Mercy Hospital for Women, Heidelberg, Victoria, Australia
- ³⁷Gynecology and Obstetrics, Charles University First Faculty of Medicine, Praha, Praha, Czech Republic
- ³⁸Gynecology and Obstetrics, General University Hospital in Prague, Praha, Czech Republic
- ³⁹Gynecology Oncology, Universidad de Chile, Santiago de Chile, Chile
- ⁴⁰Gynecology Oncology, Hospital Clínico San Borja Arriaran, Santiago, Chile
- ⁴¹Gynaecology Oncology, Mater Misericordiae University Hospital, Dublin, Ireland
- ⁴²School of Medicine, University College Dublin, Dublin, Ireland
- ⁴³Gynaecological Oncology, Royal Marsden NHS Foundation Trust, London, UK
- ⁴⁴Gynaecology, St George's University of London, London, UK

⁴⁵Gynecology, Leiden University Medical Center, Leiden, Zuid-Holland, The Netherlands

⁴⁶Obstetrics and Gynaecology, University of Hong Kong Li Ka Shing Faculty of Medicine, Hong Kong, Hong Kong

⁴⁷Surgery and Cancer, Imperial College London, London, UK

⁴⁸Center for Clinical Research, Faculty of Medicine, University of Queensland, Herston, Queensland, Australia

⁴⁹Queensland Centre for Gynaecologic Cancer Research, Royal Brisbane and Women's Hospital, Herston, Queensland, Australia

Twitter Michael Frumovitz @frumovitz, Mario Leitao @leitaomd, Julio Lau @drjuliolau, Fabio Martinelli @DrFMartinelli and Thomas Edward Ind @ThomasInd

Acknowledgements We thank all participants for their expertise in developing this tool. We also thank Vanessa Behan, Queensland Centre for Gynaecological Cancer Research, University of Queensland, Centre for Clinical Research, and QLD Australia for administrative support with the project.

Contributors KM, AO, MJ, and GH contributed to the design of the trial. KM, AO, and MJ contributed to manuscript writing. All authors contributed to data acquisition, interpretation of data, revising the draft for intellectual content, final approval of the version to be published, and agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy and integrity of any part of the work were appropriately investigated and resolved.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests AO reports grants and personal fees from Surgical Performance PTY LTD, grants from Medtronic, outside the submitted work; NRA-R reports grants from Stryker/Novadaq, outside the submitted work; MF reports grants from Astra Zeneca, grants from Tesaro/GSK, grants and personal fees from Stryker, grants from Biom'Up, outside the submitted work; ML reports Ad hoc consulting from Intuitive Surgical, serves on advisory board for Ethicon, partial grant support from NIH/NCI Memorial Sloan Kettering Cancer Center Support, outside the submitted work; TEI reports personal fees from Medtronic, personal fees from Intuitive Surgical, outside the submitted work; RK reports personal fees from Intuitive Surgical Inc., personal fees from Medtronic, personal fees from Medcaroid, outside the submitted work, and President of SERGS and Council Member of IGCS; HF reports personal fees from Intuitive Surgical Inc, outside the submitted work; JP reports personal fees from Intuitive Surgical Inc., outside the submitted work; AA reports grants and a site PI, speaker fees and serves on advisory board for Astrazeneca, serves on the advisory board for GSK, grants and grants and site Co-PI and speaker fees from Merck, speaker fees from Sanofi, grants from Pfizer, grants from Clovis, grants from CancerCare Manitoba Foundation, grants from Canadian Clinical Trials group, outside the submitted work.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. All data relevant to the study are included in the article or uploaded as supplementary information. The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Michael Frumovitz <http://orcid.org/0000-0002-0810-2648>

Nadeem R Abu-Rustum <http://orcid.org/0000-0001-9689-1298>

Alessandro Buda <http://orcid.org/0000-0002-7093-6862>

Rene Pareja <http://orcid.org/0000-0003-0093-0438>

Henrik Falconer <http://orcid.org/0000-0003-4427-9421>

Fabio Martinelli <http://orcid.org/0000-0002-4863-1747>

Thomas Edward Ind <http://orcid.org/0000-0001-5260-199X>

REFERENCES

- Cook JA. The challenges faced in the design, conduct and analysis of surgical randomised controlled trials. *Trials* 2009;10:1–9.
- Ergina PL, Cook JA, Blazeby JM, et al. Challenges in evaluating surgical innovation. *Lancet* 2009;374:1097–104.
- Simon C, Caballero C, Gregoire V, et al. Surgical quality assurance in head and neck cancer trials: an EORTC Head and Neck Cancer Group position paper based on the EORTC 1420 'Best of' and 24954 'larynx preservation' study. *Eur J Cancer* 2018;103:69–77.
- Tsai AY-C, Mavroveli S, Miskovic D, et al. Surgical quality assurance in COLOR III: standardization and competency assessment in a randomized controlled trial. *Ann Surg* 2019;270:768–74.
- Claassen YHM, de Steur WO, Hartgrink HH, et al. Surgicopathological quality control and protocol adherence to lymphadenectomy in the CRITICS Gastric Cancer Trial. *Ann Surg* 2018;268:1008–13.
- Harris A, Butterworth J, Boshier PR, et al. Development of a reliable surgical quality assurance system for 2-stage esophagectomy in randomized controlled trials. *Ann Surg* 2020. doi:10.1097/SLA.0000000000003850. [Epub ahead of print: 27 Mar 2020].
- Markar SR, Wiggins T, Ni M, et al. Assessment of the quality of surgery within randomised controlled trials for the treatment of gastro-oesophageal cancer: a systematic review. *Lancet Oncol* 2015;16:e23–31.
- Amant F, Mirza MR, Koskas M, et al. Cancer of the corpus uteri. *Int J Gynaecol Obstet* 2015;131 Suppl 2:S96–104.
- de Boer SM, Powell ME, Mileskin L, et al. Adjuvant chemoradiotherapy versus radiotherapy alone in women with high-risk endometrial cancer (PORTEC-3): patterns of recurrence and post-hoc survival analysis of a randomised phase 3 trial. *Lancet Oncol* 2019;20:1273–85.
- Boronow RC, Morrow CP, Creasman WT, et al. Surgical staging in endometrial cancer: clinical-pathologic findings of a prospective study. *Obstet Gynecol* 1984;63:825–32.
- Creasman WT, Morrow CP, Bundy BN, et al. Surgical pathologic spread patterns of endometrial cancer. A Gynecologic Oncology Group study. *Cancer* 1987;60:2035–41.
- Mikuta JJ. International Federation of Gynecology and Obstetrics staging of endometrial cancer 1988: FIGO staging of endometrial cancer. *Cancer* 1993;71:1460–3.
- Benedetti Panici P, Basile S, Maneschi F, et al. Systematic pelvic lymphadenectomy vs. no lymphadenectomy in early-stage endometrial carcinoma: randomized clinical trial. *J Natl Cancer Inst* 2008;100:1707–16.
- Kitchener H, Swart AMC, group Astudy, et al. ASTEC Swart group. Efficacy of systematic pelvic lymphadenectomy in endometrial cancer (MRC ASTEC trial): a randomised study. *Lancet* 2009;373:125–36.
- National comprehensive cancer network practice guidelines in oncology: uterine neoplasms 2020.
- Rossi EC. Current state of sentinel lymph nodes for women with endometrial cancer. *Int J Gynecol Cancer* 2019;29:613–21.
- Abu-Rustum NR, Khoury-Collado F, Pandit-Taskar N, et al. Sentinel lymph node mapping for grade 1 endometrial cancer: is it the answer to the surgical staging dilemma? *Gynecol Oncol* 2009;113:163–9.
- Rossi EC, Kowalski LD, Scalici J, et al. A comparison of sentinel lymph node biopsy to lymphadenectomy for endometrial cancer staging (FIRES trial): a multicentre, prospective, cohort study. *Lancet Oncol* 2017;18:384–92.
- Casarin J, Multinu F, Abu-Rustum N, et al. Factors influencing the adoption of the sentinel lymph node technique for endometrial cancer staging: an international survey of gynecologic oncologists. *Int J Gynecol Cancer* 2019;29:60–7.
- Holloway RW, Abu-Rustum NR, Backes FJ, et al. Sentinel lymph node mapping and staging in endometrial cancer: a Society of Gynecologic Oncology literature review with consensus recommendations. *Gynecol Oncol* 2017;146:405–15.
- Obermair A, Abu-Rustum NR. Sentinel lymph node mapping in endometrial cancer – areas where further research is needed. *Int J Gynecol Cancer* 2020;30:283–4.
- Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401–9.
- Harris PA, Taylor R, Minor BL, et al. The REDCap Consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208:1–9.
- Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.

- 25 Foster JD, Mackenzie H, Nelson H, *et al.* Methods of quality assurance in multicenter trials in laparoscopic colorectal surgery: a systematic review. *Ann Surg* 2014;260:220–9.
- 26 Sprangers MA, Cull A, Bjordal K, *et al.* The European Organization for Research and Treatment of Cancer. approach to quality of life assessment: guidelines for developing questionnaire modules. EORTC Study Group on quality of life. *Qual Life Res* 1993;2:287–95.
- 27 Miskovic D, Ni M, Wyles SM, *et al.* Is competency assessment at the specialist level achievable? A study for the National Training Programme in laparoscopic colorectal surgery in England. *Ann Surg* 2013;257:476–82.
- 28 Curtis NJ, Foster JD, Miskovic D, *et al.* Association of surgical skill assessment with clinical outcomes in cancer surgery. *JAMA Surg* 2020;155:590–7.
- 29 Cook JA, McCulloch P, Blazeby JM, *et al.* IDEAL framework for surgical innovation 3: randomised controlled trials in the assessment stage and evaluations in the long term study stage. *BMJ* 2013;346:f2820.
- 30 Harold JA, Uyar D, Rader JS, *et al.* Adipose-only sentinel lymph nodes: a finding during the adaptation of a sentinel lymph node mapping algorithm with indocyanine green in women with endometrial cancer. *Int J Gynecol Cancer* 2019;29:53–9.
- 31 Janda M, Gebiski V, Davies LC, *et al.* Effect of total laparoscopic hysterectomy vs total abdominal hysterectomy on disease-free survival among women with stage I endometrial cancer: a randomized clinical trial. *JAMA* 2017;317:1224–33.
- 32 Ramirez PT, Frumovitz M, Pareja R, *et al.* Minimally invasive versus abdominal radical hysterectomy for cervical cancer. *N Engl J Med* 2018;379:1895–904.
- 33 Frumovitz M, Plante M, Lee PS, *et al.* Near-Infrared fluorescence for detection of sentinel lymph nodes in women with cervical and uterine cancers (FILM): a randomised, phase 3, multicentre, non-inferiority trial. *Lancet Oncol* 2018;19:1394–403.
- 34 Cusimano MC, Walker R, Bernardini MQ, *et al.* Implementing a cervical sentinel lymph node biopsy program: quality improvement in gynaecologic oncology. *J Obstet Gynaecol Can* 2017;39:659–67.