Abstract 218 Table 1 Multivariate analysis for overall survival. Recurrence-free survival, and disease-free survival of cervical cancer patients

	Hazard Ratio (CI 95%)	P-value
5-year mortality		
HPV negative	2.19 (0.99-5.01)	0.062
Advanced age	1.09 (1.01-1.05)	0.008
Higher stage at diagnosis (183-48)	7.27 (2.52-20.99)	<0.001
5-year recurrence		
HPV negative	1.75 (0.66-4.65)	0.261
Adenocarcinoma	2.77 (1.28-6.03)	0.010
Chemo/Radiation as primary treatment	3.06 (1.29-7.29)	0.011
Advanced age	1.023 (0.99-1.05)	0.084
Disease-free survival		
HPV negative	1.57 (0.73-3.39)	0.246
Advanced age	1.03 (1.01-1.05)	0.003
Chemo/Radiation as primary treatment	3.79 (1.89-7.62)	<0.001
Adenocarcinoma	1.65 (0.88-3.12)	0.120

Abstract 218 Figure 1 Kaplan Meier curve of overall survival, comparing women with HPV-negative and HPV-positive cervical cancer

negativity ($P=0.062$). Median overall survival for HPV-positive CC was not reached, compared to 24 months for HPV-negative CC. Kaplan-Meier curves showed lower rates of overall survival ($P=0.011$), recurrence-free survival ($P=0.005$) and disease-free survival ($P=0.008$) for women with HPV-negative compared to HPV-positive CC.

Conclusion* The relatively poor prognosis of HPV-negative CC is important in light of its relatively high prevalence, which could increase proportionally to HPV-positive CC due to increased HPV screening and vaccination. Further studies are needed to confirm if HPV status is truly an independent prognostic factor in CC.

226

CLEARANCE OF HPV AFTER CONIZATION OF CERVICAL CANCER AND ADENOCARCINOMA IN SITU CORRELATES WITH ABSENCE OF CANCER

¹N Kugelman*, ¹E Siegler, ¹Y Goldberg, ²Y Siegler, ³P Shaked Mishan, ⁴S Mazareb, ¹L Mackuli, ⁴E Sabo, ²O Lavie, ²Y Segev. ¹Carmel Medical Center, Obstetrics and Gynecology, Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ²Rambam Health Care Campus, Obstetrics and Gynecology, Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ³Carmel Medical Center, Clinical Serology and Virology Laboratory, Haifa, Israel; ⁴Carmel Medical Center, Pathology, Haifa, Israel

10.1136/ijgc-2021-ESGO.19

Introduction/Background* About 40% of CC are in women under age 40 years, for whom fertility-sparing treatment might be very important and should always be considered. More than half the patients who undergo radical surgery (trachelectomy or hysterectomy) have no remnant tumor in the final pathology specimen. HPV clearance is a test of cure after conisation, but it is not a test of cure in CC. Our objective was to assess correlation of clearance of high-risk human papillomavirus (HR-HPV) after large loop excision of the transformation zone (LLETZ) with absence of residual disease, in women diagnosed with cervical cancer (CC) and Adenocarcinoma in Situ (AIS).

Methodology Data was collected from 92 women diagnosed with CC and AIS who were positive to High-Risk HPV (HR-HPV), and had a repeat cervical HPV test 3-12 weeks post-LLETZ, and before final surgical treatment. We compared characteristics of women with negative and positive HR-HPV post-LLETZ.

Result(s)* The pathological results of women who were HR-HPV negative ($n=40$) compared to HR-HPV positive ($n=52$) at the post-LLETZ follow-up visit included a significantly higher incidence of AIS: 14 (35%) vs 5 (9.6%) ($p<0.006$). In the negative HR-HPV post-LLETZ group, 36 (90%) had normal histology and only 2 (5%) had cancer in the final histological specimen. Among women who underwent radical hysterectomy/trachelectomy after LLETZ, a normal final histology was observed in 75% and 9% of those who were HR-HPV negative and HR-HPV positive, respectively ($p<0.0005$). The negative predictive value for residual cancer, with clearance of HR-HPV after LLETZ was 95%.

Conclusion* Clearance of HR-HPV from the cervix a short time after LLETZ has a high correlation with the absence of residual cancer in the final outcome, either in the pathology or the follow up. Testing for HR-HPV a short time after LLETZ might serve as a parameter for risk assessment.

245

THE SIGNIFICANCE OF SURGICAL ASSESSMENT IN ONCOLOGICAL OUTCOMES AFTER RADICAL HYSTERECTOMY FOR EARLY-STAGE CERVICAL CANCER. A MULTICENTER STUDY

¹S Fernandez-Gonzalez*, ²M Barahona, ³A Gil-Moreno, ³NR Gómez-Hidalgo, ⁴B Diaz-Feijoo, ⁵P Coronado, ⁵V González, ⁵A Casajuana, ⁶JM Silvan, ⁶L Melero, ⁶MA Martinez, ⁷J Hilario de la Rosa, ⁷I Lobo, ⁷E Beiro, ⁸J Frias-Gomez, ⁸Y Benavente, ²C Ortega, ⁹E Martinez, ²L Marti, ²J Ponce. ¹University Hospital of Bellvitge (IDIBELL), Gynecology, L'Hospitalet de Llobregat, Spain; ²University Hospital of Bellvitge (IDIBELL), Gynecology, L'Hospitalet de Llobregat (Barcelona), Spain; ³Hospital Universitari Vall d'Hebron, Department of Gynecologic Oncology, Barcelona; ⁴Hospital Clínic de Barcelona, Gynecology, Barcelona, Spain; ⁵Hospital Clínic San Carlos, Instituto de Salud de la Mujer, Madrid, Spain; ⁶Hospital Universitario Virgen del Rocío, Gynecology, Sevilla, Spain; ⁷Hospital Universitario de Basurto, Gynecology, Bilbao, Spain; ⁸Catalan Institute of Oncology (IDIBELL), Cancer Epidemiology Research Programme, L'Hospitalet de Llobregat, Spain; ⁹University Hospital of Bellvitge (IDIBELL), Nursery, L'Hospitalet de Llobregat (Barcelona), Spain, Spain

10.1136/ijgc-2021-ESGO.20

Introduction/Background* Patients with early-stage cervical cancer [CC] had worse prognosis when operated by minimal invasive surgery according to LACC trial. Different hypothesis were suggested such as the intrauterine manipulator, the CO₂ or the lack of protective manoeuvre. However, the effect of surgical expertise among patients who underwent radical hysterectomy by the same approach has not been evaluated.