Conclusion* In neoplasms where the initial symptoms are mild, such as endometrial cancer, the number of cases has been reduced, which may be due to the lack of consultation by patients. Regarding ovarian cancer, we have not seen a decrease in cases, although we have seen an increase in the stage. It is expected that the consequences of the COVID pandemic will continue to be felt in the future with an increase in the advanced stages of these neoplasms.

695 ACS NSQIP – PERSONALISED RISK PREDICTION TOOL FOR POSTOPERATIVE COMPLICATIONS IN GYNAEONCOLOGY SURGERY?

¹L Sevinyan*, ²S Jones, ³J Horne, ³R Bharathan, ¹A Tailor, ¹S Butler-Manuel, ⁴P Williams, ^{1,5}T Madhuri. ¹Royal Surrey County Hospital, Gynaecology, Guildford, UK; ²Cardiff University, UK; ³University Hospitals of Leicester, UK; ⁴University Of Surrey, UK; ⁵University Of Brighton, UK

10.1136/ijgc-2021-ESGO.315

Introduction/Background* Despite the informed consent process, patients' understanding of potential post-operative complications is often limited, making it difficult to call the decision an informed one, so estimating the risk of postoperative complications is important for shared decision making and to help multidisciplinary teams plan postoperative care. Increased incidence of gynaecological cancers and operations, especially technically challenging minimally invasive surgery (MIS) in older, obese and patients with multiple comorbidities, requires accurate prediction of the likelihood of mortality and morbidity and patient involvement in joint decision making about the management. ACS-NSQIP (American College of Surgeons National Surgical Quality Improvement Program) risk calculator is a validated web-based tool based on 21 preoperative risk factors to predict 8 post-operative outcomes. The objective of our study was to explore the validity of ACS NSQIP in gynaecology for perioperative prediction of postoperative complications.

Abstract 695 Figure 1

Methodology A retrospective multicentre cohort study evaluated 1552 patients who underwent surgery at a tertiary oncology centre. Data collection undertaken through dedicated database. Data collated on 764 patients undergoing robotic, 248 laparoscopic and 540 open surgery for suspected or confirmed gynaecological malignancy. All missing data collated from patient notes. Following data lock with the actual post-op event/complication that occurred in this retrospective cohort, ACS NSQIP used to count predictive scores for each patient. Data analysis evaluating ACS-NSQIP validity and relevance in gynaecological oncology patients and its ability to predict postoperative complications performed.

Result(s)* ACS-NSQIP was found to best predict mortality (AUC - 0.908), it was most accurate for prediction of complications as follows: discharge to rehabilitation (AUC-0.875), cardiac complications (AUC-0.854), sepsis (AUC-0.795), pneumonia (AUC-0.764), return to theatre (AUC-0.696), surgical site infection (AUC-0.686), VTE (AUC-0.676), readmission (AUC-0.669), renal failure (AUC-0.601). Poor result in the prediction of UTI (AUC-0.543) was noted.

Conclusion* ACS-NSQIP risk calculator appears to predict major complications and post-operative mortality making it useful as an informed consent tool. Preliminary data suggests that further validation is required to fully evaluate if the risk scores may be used to inform patients pre-operatively of their risk of complications and is currently being undertaken.

726 APPLICABILITY OF PRE-OPERATIVE PATIENT REPORTED DUKE ACTIVITY SCALE INDEX IN PREDICTION OF POSTOPERATIVE COMPLICATIONS IN GYNAEONCOLOGICAL ONCOLOGY

¹L Sevinyan*, ¹A Tailor, ²P Prabhu, ³P Williams, ^{1,4}T Madhuri. ¹Royal Surrey County Hospital, Gynaecology, UK; ²Royal Surrey County Hospital, Anaesthetics, UK; ³University Of Surrey, UK; ⁴University Of Brighton, UK

10.1136/ijgc-2021-ESGO.316

Introduction/Background* Increase in the incidence of gynaecological cancers has resulted in increased operative procedures, specifically in patients with multiple comorbidities including obesity and frailty. This is often associated with prolonged admission and higher rates of postoperative mortality and morbidity and presents a challenge with an unmet need for an accurate, personalised risk prediction. Duke Activity Scale Index (DASI) is a 12 item scale in the form of self-reported questionnaire based around commonly performed activities of daily living. Currently, DASI is used to evaluate patients with cardiovascular diseases, however there is growing interest in utilising it in preoperative setting in different specialities. This study investigates the accuracy of DASI in pre-operative prediction of postoperative outcomes in gynaecology.

Methodology A retrospective cohort study of 486 patients who had undergone an operative treatment at a tertiary oncology centre. Data collection undertaken through dedicated gynaecology database and missing data collected through patients' records. All patients had completed the DASI questionnaire prior to their consultation. Actual postoperative 30 day complications and the length of stay also recorded. DASI was then compared with the occurrence of postoperative complications.

Result(s)* 242 patients had a *Da Vinci* robot-assisted procedure, 44 - laparoscopic and 200 - open surgery. 133/486 were