

Diagonal segments are produced by ties.

Abstract 726 Figure 1

classified as having any type of complications within 30 days of the operation. Our results have shown that the higher DASI score the less likely patients were to have postoperative complications. This result was statistically significant with odds ratio of 0.973 and confidence interval between 0.961 and 0.986. We were also able to demonstrate that for every 10 points further up the DASI score a patient was 0.761 times less likely to have a postoperative complication. Hence general morbidity prediction of DASI score has been found to statistically significantly predict postoperative complications (AUC-0.740).

Conclusion* Our study has shown that DASI self-reported score is a useful predictive tool of perioperative estimation of postoperative complications in the gynaecology setting. Further analysis with a larger sample size and a multicentre prospective study is currently underway to validate the findings.


757

EUROPEAN ENHANCED RECOVERY AFTER SURGERY (ERAS) GYNECOLOGICAL ONCOLOGY SURVEY: CURRENT STATE OF PERIOPERATIVE PRACTICE

¹NR Gómez-Hidalgo*, ²A Pletnev, ³Z Razumova, ⁴N Bizzari, ⁵I Selcuk, ⁶C Theofanakis, ⁷K Zalewski, ⁸T Nikolova, ⁹M Lanner, ¹⁰J Kacperczyk-Bartnik, ¹¹U Acosta, ¹²PB Asuncion, ¹³A Gil-Moreno, ¹⁴G Nelson, ¹⁵C Fotopoulou, ¹⁶JL Sánchez Iglesias, ¹⁷Hospital Universitari Vall d'Hebron, Gynecology Oncology, Barcelona, Spain; ¹⁸Alexandrov National Cancer Centre of Belarus, Gynecology Oncology, Belarus; ¹⁹Karolinska Institute, Gynecology Oncology, Sweden; ²⁰Fondazione Policlinico Universitario A. Gemelli, Roma, Italy; ²¹Maternity Hospital, Ankara City Hospital, Ankara, Turkey; ²²Alexandra General Hospital, Athens, Greece; ²³Świętokrzyskie Cancer Centre, Kielce, Poland; ²⁴Heidelberg University, Heidelberg, Germany; ²⁵Cardinal Schwarzenberg Hospital, Schwarzach im Pongau, Austria; ²⁶Medical University of Warsaw, Warszawa, Poland; ²⁷Cumming School of Medicine U C, Calgary, Canada; ²⁸Imperial College Healthcare NHS Trust, UK

10.1136/ijgc-2021-ESGO.317

Introduction/Background* Enhanced Recovery After Surgery (ERAS) reduces complication rates after surgery, decreases length of hospital stay (LOS) and reduces costs. A correct



Abstract 757 Figure 1 Participating European Centres




Figure 2. 2. Intra-operative ERAS Items





Figure 2. 3. Postoperative ERAS Items



Abstract 757 Figure 2 Centres Practicing ERAS Items

implementation of the ERAS protocol and adherence to all ERAS items contribute to better outcomes in gynecological oncology surgery.

The aim of this survey is to acquire a comprehensive picture of the current status of implementation of the ERAS protocol among European centers.

Methodology A 45-item questionnaire survey, investigating ERAS practice preferences in pre-, intra-, and post-operative management was launched between December 2020 and May 2021. An on-line questionnaire link was e-mailed to the national representatives (NATREPs) of the European Network of Young gynecologic oncologists (ENYGO), who then shared the survey with their centers. In order to avoid duplicate data, NATREPs selected one referral person from the highest volume centers performing complex surgery per country. Two weeks later a reminder was sent to non-responders.

Result(s)* A total of 116 responses were collected. 73% of centers were academic/teaching hospitals. Overall, 70% of respondents reported that ERAS was implemented at their institution. The median LOS for advanced ovarian cancer surgery was between 5-7 days according to 63% of respondents. 81% of respondents reported a median LOS between 2-4 days among patients who underwent surgery for early-stage gynecological cancer. An overall compliance rate between 60% and 80% was reported by 44,29% of centers.

ERAS items with well adherence to the guidelines were: deep vein thrombosis prophylaxis, antibiotic prophylaxis, prevention of hypothermia and early mobilization. Regarding pre-operative ERAS items, 28% of respondents reported bowel preparation as 'sometimes-normally' performed. 60% of respondents described ERAS implementation as a challenge and 76% reported being reluctant to change clinical practice.

Conclusion* This European survey of ERAS in gynecologic oncology surgery shows that there is still a lack of implementation of the ERAS protocol across Europe and European centers need to increase ERAS gynecologic oncology guideline compliance to improve patient outcomes. Therefore, European centers need a further protocol and guidelines that encompass the way of ERAS implementation

798

THE LOCKDOWN EFFECT ON GYNAECOLOGICAL CANCER SURGERIES DURING THE COVID-19 PANDEMIC

V Theodoulidis*, DE Vlachos, C Theofanakis, V Pergialiotis, N Thomakos, A Rodolakis, D Haidopoulos. *National University of Athens, First Department of Obstetrics and Gynecology, Athens, Greece*

10.1136/ijgc-2021-ESGO.318


Introduction/Background* The impact of COVID-19 pandemic caused a disruption of the healthcare systems and led to significant delays in diagnosis and treatment of gynecological cancer patients. New algorithms that aim to sustain balance between management of oncological patients and the need to maintain a sufficient amount of resources were adapted.

Methodology This retrospective study reviewed the patients with gynecological cancer operated in our hospital during the first lockdown period in Greece (between 13 March 2020 and 30 May 2020) and compared the results with the corresponding time period in 2019 before COVID-19 pandemic. We also examine the number of patients that were referred for neoadjuvant chemotherapy or radiotherapy between those periods.

Result(s)* The gynecological oncological operations performed during the lockdown period of the first pandemic wave were

not altered by the outbreak (153 in 2019 vs 130 in 2020) (figure 1). There was no difference in ovarian cancer surgeries (34 vs 31) and the number of primary debulking was not affected (20 vs 19). The patients referred to neoadjuvant chemotherapy was the same between the two periods. No significant difference was obtained in the endometrial, vaginal and cervical cancer surgeries and the number of surgically treated recurrences. Contrastingly there was a significant decrease in endoscopic procedures and diagnostic biopsies (72 vs 53) and the fertility sparing surgeries were postponed from patients at a later date (table 1).

Conclusion* The findings of our study suggest that at the first wave of the pandemic lockdown, the operations conducted in our department did not alter. In accordance with international consensus guidelines the stage, the grade and the type of cancer, and the potential comorbidities were the main factors that accounted for the decision of the optimal mode of treatment.



Abstract 798 Figure 1 Gynecological cancer surgeries on both periods

Abstract 798 Table 1 Type of procedures on both periods

	13/3/2019 – 30/05/2019	13/3/2020 – 30/5/2020
OVARIAN CANCER	34	31
Primary Cytoreduction	20	19
Interval – Late Debulking	5	6
Neoadjuvant Chemotherapy	7	7
Final Benign Histology	9	6
ENDOMETRIAL CANCER	28	23
CERVICAL CANCER	2	2
Radical Hysterectomy	1	2
Trachylectomy	1	-
RECURRENCE	7	7
Ovarian Cancer	3	3
Endometrial Cancer	2	-
Cervical Cancer	-	1
Vulvar Cancer	2	3
REOPERATION	2	5
DIAGNOSTIC LAPAROSCOPY	11	8
MINOR PROCEDURE	61	45
Pigtail - Cystoscopy	42	31
EUA - Biopsy - D&G	19	14
FERTILITY SPARRING SURGERY	5	0