4 cm; 2 (25%) had a 1a1 stage. No patient had a tumor larger than 4 cm on examination, MRI, and histological examination. No patient received neoadjuvant chemotherapy. 2 patients (25%) underwent a biopsy for the purpose of establishing a diagnosis, 6 patients (75%) conization of the cervix. The average age of the patients was 28 years (from 26 to 37 years). 5 (62.5%) patients were nulliparous, 2 patients had 2 children, 1 patient had 1 child.

LVSI was negative in preoperative histological examination, and resection margins were also negative. In 1 patient (12.5%) after histological examination LVSI was positive, in 7 it was negative. Of the 8 patients who retained fertility, there were 5 pregnancies, 2 miscarriages at 9–10 weeks, and 3 deliveries at 36–37 weeks of gestation.

Conclusion* Thus, in 2013–2021, 8 radical trachelectomy operations were successfully performed. The data presented in this publication demonstrate that patients with stage IB1 tumors ranging in size from 2 to 4 cm and with favorable histology are acceptable candidates for attempted radical trachelectomy. These patients must be clearly informed of all the options if they want to consider maintaining fertility.

77

SINGLE INSTITUTION OUTCOMES FOR CERVICAL CANCER PATIENTS TREATED WITH PALLIATIVE SYSTEMIC ANTI CANCER THERAPY

¹C Cotronei*, ²A Stillie, ²M Zahra. ¹University of Edinburgh, UK; ²Western General Hospital Edinburgh Cancer Centre, Edinburgh, UK

10.1136/ijgc-2021-ESGO.4

Aim Bevacizumab, in combination with chemotherapy, has been implemented as first line treatment of patients with metastatic cervical cancer since 2015. This is a single institution review to evaluate the overall survival (OS) and progression free survival (PFS) for patients with advanced cervical cancer treated with palliative systemic anti cancer therapy (SACT) over a 5-year period from 2015-2020.

Methodology Patient records were reviewed retrospectively for all cervical cancer patients who received palliative SACT in a single institution between 2015-2020. Patient demographics, previous treatment for cervical cancer (if applicable), palliative SACT regimen received, toxicities, and outcome data were recorded. OS and PFS were calculated for the whole cohort and for the following subgroups: patients treated with Bevacizumab in combination with palliative chemotherapy; patients treated with chemotherapy alone. Toxicities were recorded by Common Toxicity Criteria (CTC).

Result(s)* 52 patients received palliative SACT during the 5-year period following a diagnosis of metastatic cervical cancer. The median age was 51 (range 25-76). The majority of patients had a squamous cell histology (75%). 40 patients received Bevacizumab in combination with chemotherapy, 12 were treated with chemotherapy alone due to comorbidities. The median OS for the whole patient cohort was 64 weeks and PFS was 45 weeks. Patients who received bevacizumab in combination with chemotherapy had a median OS of 76 weeks compared to 50 weeks in the chemotherapy only subgroup. Grade 1 and 2 toxicities were comparable in patients receiving chemotherapy alone and those treated with Bevacizumab. 10% of patients in the cohort receiving bevacizumab developed a thromboembolic event compared with 5.5% in the chemotherapy only cohort.

Conclusion* Results from this study of real world data are comparable with the published data. The results support the continued use of Bevacizumab, in combination with chemotherapy, in our patient population as the combination regimen improves OS with acceptable toxicities.

106

COMPARISON OF LOCALLY ADVANCED CERVICAL CANCER TREATMENT GUIDELINES IN EUROPE

¹A Leary*, ²B Monk, ³J Takyar, ⁴A Nunes, ⁴JD Hernández Chagüi, ⁴K Rabon-Stith, ⁵E Pujade-Lauraine. ¹Gustave Roussy Cancer Center, University Paris Saclay, Department of Medical Oncology, Villejuif, France; ²University of Arizona College of Medicine, Creighton University School of Medicine, Division of Gynecologic Oncology, Phoenix, USA; ³Parexel International, HEOR Evidence Evaluation Regulatory and Access, Mohali, India; ⁴AstraZeneca, Gaithersburg, USA; ⁵Arcady-Gineco, Paris, USA

10.1136/ijgc-2021-ESGO.5

Introduction/Background* From our previously conducted global systematic literature review (SLR) assessing recommended treatment for locally advanced cervical cancer (LACC) it was observed that there was international consensus on the use of concurrent chemoradiotherapy (cCRT) as standard of care (SoC) for Stage IIB-IVA LACC, but recommendations for Stage IB2-IIA LACC varied. Here, we present a subanalysis of European guidelines to determine if recommended LACC treatment is consistent across Europe.

Methodology English-language cervical cancer treatment guidelines and consensus statements were identified through literature databases (1999-2020), international and national health organizations, and general internet searches. Included guidelines represented the latest updates from Europe. Non-English guidelines from EU-5 countries were translated for comparison (Italian [AIOM], German [S3], and French [HAS]) to the 4 European guidelines obtained through this SLR protocol (ESTRO/ESGO/ESP, ESMO, Spain [SEOM], UK [NHS]). An English-language guideline from Germany was excluded because a newer German-language only version was available.

Result(s)* The 7 guidelines were updated between 2010 and 2021 (table 1). ESTRO/ESGO/ESP guidelines used AJCC 8th edition TNM criteria for staging. All others, except SEOM, used FIGO 2009 criteria; SEOM used FIGO 2018 criteria. Table 1 shows recommended treatment by LACC stage. For Stage IB2-IIA2 disease, cCRT was the SoC or an alternative to radical hysterectomy. AIOM was the only guideline to recommend radiotherapy alone for Stage IIA1 disease. cCRT followed by brachytherapy was SoC for suitable patients with Stage IIB-IVA LACC among all European guidelines. For cCRT, a platinum-based regimen was recommended by all, and most guidelines recommended an external beam radiation therapy (EBRT) dose of 45-50 Gy (table 2).

Conclusion* In line with global SLR findings, consensus on the use of cCRT as primary treatment for Stage IIB-IVA LACC was recommended by European guidelines. The ideal cCRT regimen was weekly cisplatin at 40 mg/m² concurrent with EBRT at 45-50 Gy and followed by brachytherapy. For Stage IB-IIA LACC, recommendations mainly varied between radical hysterectomy and cCRT as options or cCRT alone. Alignment of guidelines with FIGO 2018 staging criteria may help reduce variation in recommended treatment for early-stage LACC.

Funding AstraZeneca

Klikněte nebo klepněte sem a zadejte text.