Abstract 186 Figure 1 Kaplan-Meier plots of overall survival within tumour types according to complete (TR0) and largely (TR1-10mm) debulking; number of patients included into the clear cell cohort was too small to allow meaningful OS analyses

Abstract 186 Table 1 OR: odds ratio; OS: overall survival; HR: hazard ratio; *number of patients included into the clear cell cohort was too small to allow meaningful OS analyses

	High grade	Low grade	Mucinous	Clear cell
n	5,156	299	219	71
TR0 (%)	32.7%	52.2%	31.1%	49.3%
TR1-10 (%)	31.7%	26.1%	26.0%	21.1%
OR (95%CI) for TR0	1 (reference)	1.84 (1.44–2.36)	0.95 (0.70–1.29)	1.69 (1.03–2.79)
Deaths n (%)	3,122 (60.6%)	115 (38.5%)	155 (70.8%)	46 (64.8%)
OS HR(95%CI) TR0 vs TR>10	0.48 (0.43–0.52)	0.39 (0.24–0.64)	0.64 (0.40–1.03)	*
OS HR(95%CI) TR1-10 vs TR>10	0.91 (0.84–0.99)	0.87 (0.52–1.44)	1.09 (0.71–1.67)	*
OS HR(95%CI) TR0 vs TR1-10	0.52 (0.48–0.58)	0.45 (0.28–0.73)	0.59 (0.36–0.96)	*
OS HR(95%CI) TR0 vs TR>0	0.50 (0.46–0.54)	0.42 (0.28–0.64)	0.62 (0.40–0.95)	*

serous, endometrioid, undifferentiated histology). Multiple logistic regression of achieving TR0 in the full population and multiple Cox-regression of OS separately in each histological subpopulation adjusting potential confounders (treatment arm within each study, ECOG performance status, age, FIGO stage) were undertaken.

Result(s)* 5,745 pts were eligible (5,156 high grade, 299 low grade, 219 mucinous, 71 clear cell). Differences in resection rates between histological subtypes are displayed in the table

1. Adjusted odds ratios showed significantly higher odds for achieving TR0 for low grade and clear cell compared to high-grade, (table 1). Median follow-up was 60.7 months. The figure shows OS within tumour types according to resection status. Hazard ratios between TR0, TR1-10 and TR>10 obtained from multiple Cox regression of OS are shown in the table 1.

Conclusion* Our analysis confirmed the value of complete resection in all subtypes of ovarian cancer. The role of largely resection (TR1-10mm) seemed to be relevant in low grade OC, but marginal although significant in HGSOc and inconclusive in mucinous and clear cell OC. Larger databases are necessary to gain more reliable data in these subgroups.

187

PLATINUM RESISTANCE IN OVARIAN CARCINOMA LONG-TERM SURVIVORS: A RETROSPECTIVE STUDY AT THE TUEBINGEN UNIVERSITY WOMEN'S HOSPITAL

¹S Mittelstadt*, ¹M Grube, ¹T Engler, ^{1,2}E Oberlechner, ¹S Wörz, ¹S Matovina, ¹A Stöhr, ¹L Volmer, ¹EM Grischke, ¹CB Walter, ¹A Hartkopf, ³S Kommos. ¹Tuebingen University Hospital Department of Women's Health; ²University Hospital Tuebingen, Department of Women's Health, Tuebingen, Germany; ³Tuebingen University Hospital, Department of Women's Health, Tuebingen, Germany

10.1136/ijgc-2021-ESGO.353

Introduction/Background* Despite the majority of ovarian carcinoma patients (OvCa) initially responding to primary therapy, relapse occurs frequently in early and advanced stages (25 and 80% respectively) and OvCa is still associated with an overall poor prognosis. Nevertheless, therapy response and disease progression can be quite variable and long-term survival is reported in up to 30% of patients. It is known that,

even after multiple chemotherapy lines, platinum sensitivity is associate with a longer therapy-free interval. The aim of this study was to investigate platinum resistance in long-term OvCa survivors.

Methodology Patients diagnosed with OvCa at the Tuebingen University Women's Hospital between 2000 and 2012 were retrospectively identified and follow-up data was collected. For patients surviving at least 8 years a detailed chart review was performed. Platinum resistance was defined as disease progression during platinum-based chemotherapy or 6 months after the last platinum dose administration.

Result(s)* From a total n=745 of patients with adequate follow-up data, n=223(30%) survived at least 8 years after primary diagnosis. Median follow-up was 12.1 years and median age at diagnosis was 54.7 years. Relapse was recorded in 87/223(39%) patients, of which 28/87(32%) developed platinum resistance at some point. Platinum resistance was observed after the second line chemotherapy in most patients (median: 2, range: 1–4). 23/28 patients received further chemotherapy lines (median: 2, range: 0–8). Rechallenge with a platinum-based chemotherapy has been used in 9/28(32%) patients. Median overall survival of these 28 patients was 9.3 years (8.2–18.4, IQR3.7) and 4.1 (0.4–16.8, IQR3.8) years after platinum resistance.

Conclusion* We were able to demonstrate long-term survival after platinum resistance in a substantial number of ovarian carcinoma patients. Suggesting therefore, that other parameters influence the disease behaviour. Further research of patient characteristics, environmental and genetic features and treatment modalities will help to further understand factors contributing to long-term survival.

188

PLASMAJET GIVES LESS TISSUE DAMAGE THAN ELECTROCOAGULATION DEVICE DURING CYTOREDUCTIVE SURGERY FOR ADVANCED-STAGE OVARIAN CANCER

^{1,2}G Nieuwenhuyzen-de Boer*, ¹N Van de Berg, ¹XS Gao, ³PC Ewing-Graham, ¹HJ Van Beekhuizen. ¹Erasmus University Medical Center, Gynaecologic Oncology, Rotterdam, Netherlands; ²Albert Schweitzer Hospital, Gynaecology, Dordrecht, Netherlands; ³Erasmus University Medical Center, Pathology, Rotterdam, Netherlands

10.1136/ijgc-2021-ESGO.354

Introduction/Background* The aim of surgery for advanced-stage ovarian cancer is complete cytoreductive surgery, because

this is the most important independent prognostic factor for prolonged survival. Yet this can be difficult to achieve when there are micrometastases on the mesentrium or intestines. The PlasmaJet® Surgical device is an instrument to remove these micrometastases, but little is known about the depth of tissue damage compared to electrocoagulation devices.

Methodology A prospective study was performed to determine the histological depth of thermal damage in tissues in a series of 106 histological slides of 17 advanced-stage ovarian cancer patients to compare thermal damage depth after use of the PlasmaJet® Surgical device with electrocoagulation devices. Depending on which tissue types were resected during complete cytoreductive surgery, samples were collected from reproductive organs (uterus, ovaries), intestines (ileum, colon, rectum) and omentum, mesentrium and peritoneum. A systematic review of the literature on tissue damage with the PlasmaJet® Surgical device was carried out.

Result(s)* Average thermal damage depth was 0.15 mm (range 0.03-0.60 mm) after use of the PlasmaJet and 0.33 mm (range 0.08-1.80 mm) after use of electrocoagulation (P<0.001). Greater disruption of the tissue surface was often observed after electrocoagulation use. These results correspond with the available literature.

Conclusion* Based on our case series and review of literature, we conclude that PlasmaJet treatment produces significantly less thermal damage than electrocoagulation treatment. It is therefore considered a safe alternative, aiding in the achievement of complete cytoreductive surgery in patients with advanced-stage ovarian cancer without complications due to perforation of hollow organs.

190

PERIOPERATIVE DYNAMIC EVALUATION OF FOLATE RECEPTOR ALPHA EXPRESSION IN OVARIAN CARCINOMA

^{1,2}MP Kubelac*, ^{1,2}C Vlad, ^{1,2}B Pop, ¹B Fetica, ¹E Fischer-Fodor, ¹O Soritau, ¹O Balacescu, ^{1,2}F Ignat, ^{1,2}G Lazar, ^{1,2}C Lisencu, ^{1,2}A Rancea, ^{1,2}A Irimie, ^{1,2}P Achimas-Cadariu. ¹The Oncology Institute "Prof. Dr. Ion Chiricuta" Cluj-Napoca, Cluj-Napoca, Romania; ²Iuliu Hatieganu University of Medicine and Pharmacy, Cluj-Napoca, Romania

10.1136/ijgc-2021-ESGO.355

Introduction/Background* The human folate receptor α (FOLR1) is overexpressed in ovarian cancer and has been correlated with the transformed phenotype in ovarian cancer, platinum sensitivity, and tumor growth. The highly restricted

Abstract 188 Figure 1