grade serous carcinoma (HGSC) in patients with a BRCA1/2 pathogenic variant (PV), but results in significant short and long-term health risks. New interventions, such as risk-reducing salpingectomy with delayed oophorectomy are promising. In this alternative approach, the detection of serous tubal intraepithelial carcinoma (STIC) as precursor to HGSC, has become more important. The detection of STIC indicates an increased risk for HGSC and would prompt for an immediate oophorectomy. Unfortunately, reproducibility of STIC diagnosis is only moderate, even among experienced gynecological pathologists. The aim of this pilot study is to develop and validate an AI algorithm for automated detection of potential STIC lesions in scanned H&E slides, to aid the pathologist in diagnosing STIC.

Methodology We collected and digitalised 60 cases of STIC and 65 control cases. STIC diagnosis was confirmed using p53 and Ki-67 immunohistochemical stainings (IHC). The dataset was split into 50 cases for training, five for validation and five for testing. We developed a Convolutional Neural Network and compared two approaches: directly detecting STIC (one-step) or first detecting all epithelium and subsequently detecting STIC within epithelial regions (two-step). Additionally, we evaluated whether we could improve the network by enriching the training data with hard negative examples.

Abstract 1009 Figure 1

Abstract 1009 Figure 2

Result(s)* We found that the optimal configuration for detection of STIC was the two-step approach, with training set enrichment by hard negatives. This network reached an area under the receiver operating curve of 0.90 (figure 1). Visual inspection of cases in the test set showed concordance between the model output, p53 and Ki67 IHC, and pathologists' annotations (see example model output on H&E stained tissue in figure 2).

Conclusion* We present a convolutional neural network that can successfully detect STIC lesions in whole slide images. AI has the potential to aid the pathologist in the detection of STIC and assist in producing more accurate and consistent diagnosis. Additional performance and robustness is expected to be achieved by expansion of the dataset.

1029 UNUSUALLY BREAST METASTASES

¹S Sakhr, ²K Tlili, ³S Kammoun, ⁴M Bouhani*, ⁵Y Houcine, ³K Hamza, ²H Bouaziz, ⁶A Ghoucha, ¹I Bettaib, ⁶M Idriss. ¹Institut Salah Azaiz, Surgical Department, Tunis, Tunisia; ²Azaiez Institute of Oncology, Department of Pathology, Tunisia; ³Salah Azaiez Institute, anatomopathology, Tunis, Tunisia; ⁴Salah Azaiez Institute, surgical department, Tunisia, Tunisia; ⁵Institut Salah Azaiz, anatomopathology, Tunis, Tunisia; ⁶Institute Salah Azaiz, anatomopathology, Tunisia, Tunisia

10.1136/ijgc-2021-ESGO.534

Introduction/Background* The breast is the most common site of primary malignancies in adult women, but is an uncommon site for metastasis from extramammary malignancies. Breast metastases from extramammary malignancies have both haematogenous and lymphatic routes. We report 10 cases of breast metastases in our pathology lab between 2014 and 2020.

Methodology We have identified all the cases of breast metastases diagnosed between 2014 and 2020.

Result(s)* We found 10 cases of breast metastases from extramammary malignancies. All patients were female with a middle age of 42 years. The primary malignancies were a melanoma in 3 cases, an ovarian carcinoma also in 3 cases, a fibrosarcoma, a rhabdomyosarcoma and a nasopharyngeal carcinoma in 1 case each. The diagnosis was based on morphological features but also on immunohistochemistry especially with the ovarian carcinoma which exhibits some similar features to a primary breast carcinoma.

Conclusion* Breast metastases from extramammary malignancies show variable histological features. Therefore, the possibility of metastatic lesion should be considered in evaluating breast lesions in patients with primary malignancy in other organs. Awareness of typical and atypical imaging features of such lesions may be helpful to diagnose metastatic lesions in the breast.

1111 ATYPICAL (C3) AND SUSPICIOUS (C4) CATEGORIES IN BREAST FINE NEEDLE ASPIRATION

¹S Sakhr, ²K Hamza, ²Y Houcine, ²S Kammoun, ²K Tlili, ³M Bouhani*, ²A Goucha, ²I Bettaieb, ²M Driss. ¹Institut Salah Azaiz, surgical department, Tunis, Tunisia; ²Institut Salah Azaiz, anatomopathology, Tunis, Tunisia; ³Institut Salah Azaiz, surgical department, Tunis, Tunisia

10.1136/ijgc-2021-ESGO.535

Introduction/Background* The present study was carried out over a 1-year period (January 2017-December 2017) and conducted at the cytology unit of Pathology Department (Salah Azaiez