227

INVASIVE MOLE: A RARE CAUSE OF HEMOPERITONEUM

¹M Zamani, ²M Mollabashi, ³S Alizadeh*. ¹Hamadan University of Medical Science, Associate Professor of Onco-Gynecology, Department of Obstetrics and Gynecology, Hamadan, Iran; ²Hamadan University of Medical Science, Assistant professor of Radiology, Department of Radiology, Hamadan, Iran; ³Board Certified of Obstetrics and Gynecology at Tehran university of medical science, Tehran, Iran

10.1136/ijgc-2021-ESGO.607

Introduction/Background* Gestational trophoblastic neoplasia comprises a unique group of human neoplastic diseases that derive from fetal trophoblastic tissues. The hydatidiform mole is the most common form of GTD, representing 80 percent of cases. An invasive mole is a hydatidiform mole characterized by the enlarged hydropic villi invading into the myometrium, into vascular spaces, or into extrauterine sites.

Methodology Case presentation: Here is a case with invasive mole after the evacuation of complete molar pregnancy, presented with an acute abdomen. We desired to preserve the uterine because our 21 years old patient doesn't have a child.

Result(s)*

Clinical Discussion An emergency abdominal ultrasound scan showed a 47*34*55 mm ill-defined hyperechoic heterogeneous mass with anechoic cystic vascular spaces within it, in the posterior wall of the uterus away from the endometrium that extended to the serous layer of the uterus. Laparotomy was done. After the evacuation of 2 L of hemoperitoneum, an approximately 5×4 metastatic, vesicular mass was seen in the posterior wall of the uterus, which was resected and uterine preservation was successful.

Conclusion* This case report describes the clinical, imaging, surgical and histopathological findings of Invasive mole after a hydatidiform molar pregnancy. Our case highlights the feasibility of fertility-preserving surgery in the case who experienced life-threatening hemorrhage due to a ruptured uterus.

273

AVELUMAB IN PATIENTS WITH GESTATIONAL TROPHOBLASTIC TUMORS RESISTANT TO POLYCHEMOTHERAPY: EFFICACY OUTCOMES OF COHORT B OF TROPHIMMUN PHASE II TRIAL

¹B You*, ¹PA Bolze, ²JP Lotz, ¹J Massardier, ³L Gladieff, ⁴A Floquet, ¹T Hajri, ¹P Descargues, ⁵C Langlois-Jacques, ⁵C Mercier, ⁶S Bin, ⁶L Villeneuve, ⁷A Roux, ⁶M Alves-Ferreira, ⁸D Grazziotin-soares, ⁹P Rousset, ¹⁰G Freyer, ¹F Golfier. ¹Centre de Référence des Maladies Trophoblastiques; Hospices Civils de Lyon, CITOHL, CICLY, Univ Lyon, Medical Oncology, LYON, France; ²Hôpital Tenon, Pôle Onco-Hématologie Hôpitaux Universitaires de l'Est Parisien, APHP, Université Pierre et Marie Curie, Medical Oncology, Paris, France; ³Institut Claudius Regaud; IUCT-ONCOPOLE, Département d'oncologie médicale; Toulouse, France; ⁴Institut Bergonié, Medical Oncology, Bordeaux, France; ⁵Hospices Civils de Lyon, CNRS UMR5558, Laboratoire de Biométrie et Biologie Evolutive, Equipe Biostatistique-Santé, Service de Biostatistique, Lyon, France; ⁶Hospices Civils de Lyon, Unité Recherche et Epidémiologie Cliniques – Pôle de Santé Publique, Centre Hospitalier Lyon Sud, Lyon, France; ⁷Hospices Civils de Lyon, Pôle de Santé Publique, Lyon, France; ⁸Hôpital Tenon, Hôpitaux Universitaires de l'Est Parisien, APHP, Université Pierre et Marie Curie, Pôle Onco-Hématologie, Paris, France; ⁹Hospices Civils de Lyon, Radiologie, Lyon, France; ¹⁰Hospices Civils de Lyon, Medical Oncology, CITOHL, Lyon, France

10.1136/ijgc-2021-ESGO.608

Introduction/Background* In patients with gestational trophoblastic tumors (GTT) with a FIGO score ≥ 7 , or GTT resistant to both standard monotherapies, the recommended polychemotherapy regimen is EMA-CO. In case of resistance to polychemotherapy, the prognosis is poor. The anti-PD-L1

monoclonal antibody avelumab may be effective for GTT resistant to monochemotherapy (You et al JCO 2020). The efficacy data of avelumab in patients with GTT resistant to polychemotherapy enrolled in cohort B of TROPHIMMUN trial (NCT03135769) are presented.

Methodology In cohort B, patients with GTT resistant to polychemotherapy received avelumab 10 mg/kg Q2W until hCG normalization, and for 3 additional cycles thereafter. The primary endpoint was the rate of patients with hCG normalization, following a 2-step Simon design. The cohort was closed prematurely for futility.

Result(s)* 2017-2020 : seven patients were treated with the French Gestational Trophoblastic Center (median age was 37 ; choriocarcinoma: 4; placental-site: 1; epithelioid: 1; other: 1) ; stage I/III: 43%/57%; FIGO score 8-10: 43%; score 11-15: 57%. Patients had experienced previous failures to monochemotherapy (n=5), pelvis surgery (n=2), and polychemotherapy (EMA-CO, n=5; EMA-EP, n=1; TP/TE, n=1; APE, n=1). They received a median of 6 avelumab cycles (range: 3-13). Six (85.7%) patients achieved initial hCG stabilization/decline, and one patient (14.2%) had successful hCG normalization after 13 cycles. Another patient experienced favorable hCG decline, but avelumab was discontinued for hemostatic hysterectomy, followed by sustained hCG normalization. The 5 other patients (71.4%) experienced hCG re-increase suggesting avelumab resistance, including two patients who developed brain hemorrhage after 4 cycles (brain metastases in one patient; arteriovenous malformation in one patient who died). The 4 remaining patients were subsequently treated with hysterectomy, other polychemotherapy, including high-dose/bone-marrow-transplant for two; pembrolizumab for one (who died).

Conclusion* TROPHIMMUN is the first trial of immunotherapy in GTT. Contrarily to avelumab suggested effectiveness in patients with monotherapy resistance (Cohort A), avelumab activity was limited in patients with polychemotherapy resistance. Despite initial changes in hCG kinetics in most patients, eventual hCG normalization was rare (14%). The prognosis of patients experiencing polychemotherapy resistance remains poor. Combination treatments with immunotherapy should be considered.

314

PARTIAL MOLAR PREGNANCY WITH A COEXISTING FETUS; CASE SERIES AND REVIEW OF THE LITERATURE

¹L Coopmans*, ¹C Lok, ¹N Van Trommel, ²M Seckl. ¹Antoni van Leeuwenhoek, Gynecologic oncology, Amsterdam, Netherlands; ²Charing Cross Gestational Trophoblastic Disease Centre, Medical oncology, London, UK

10.1136/ijgc-2021-ESGO.609

Introduction/Background* Partial molar pregnancy with a coexisting fetus (PHMCF) is a very rare entity leading to limited understanding of its natural course and optimal diagnostics and treatment. The aim of this study was to describe a case series of patients with PHMCF and to review the current available literature.

Methodology We searched the databases of the Charing Cross Trophoblastic Disease Center and the Dutch Working Party on Trophoblastic Tumors. Secondly, we performed a systematic literature review to evaluate the incidence of PHMCF, the clinical presentation, obstetrical and maternal outcomes of a pregnancy complicated by PHMCF.