Vulvar International Tumor Analysis Evaluation Form

Terms and definitions for sonographic assessment of inguinal lymph nodes in vulvar cancer

Abstract 744 Figure 1

problems of standardization and to formulate terms and procedures to derive morphologic end-points by B-mode imaging and end-point of vascularity and blood flow by color Doppler imaging. The recommendations of the steering committee were distributed to each participating center and subsequently refined. The following consensus is going to be tested in international multicentric interobserver and diagnostic accuracy studies.

Result(s)* We present a consensus of international multidisciplinary VITA group on terms, definitions and measurements which may be used to describe inguinal lymph nodes on gray-scale and color/power Doppler ultrasound. The proposed nomenclature is not limited to inguinal lymph nodes as part of vulvar cancer staging; it can be used for the assessment of peripheral lymph nodes in general, as well as non-peripheral (i.e. parietal or visceral) lymph nodes.

Conclusion* VITA terms and definitions were developed as a basis for international standardization of lymph node assessment by ultrasound and lay the foundations for prospective studies aiming to identify ultrasound features typical of metastases and other pathology in lymph nodes.

905

PRIMARY MALIGNANT MELANOMA INVOLVING THE VULVA – THE IMPORTANCE OF EARLY DIAGNOSIS

¹LP Castillo Rabazo, ¹MDLR Oliver*, ¹JM Seoane-Ruiz, ¹C Alvarez, ¹G Lopez Gonzalez, ¹B Gil Ibanez, ²J Sanz Pablos, ¹P Carmona Payan, ¹A Tejerizo. ¹Hospital 12 de octubre, ginecología y obstetricia, madrid, Spain; ²Hm hospitales montepríncipe, ginecología y obstetricia, boadilla del monte, Spain

10.1136/ijgc-2021-ESGO.627

Introduction/Background* Primary melanomas originating from the gynecological tract are rare and aggressive cancers. The majority of patients experience poor long-term survival. Our objective is to describe the diagnosis, treatment and survival of patients diagnosed with vulvar melanoma in a tertiary Spanish hospital during a 9-year period.

Methodology Retrospective descriptive cohort study of patients diagnosed with vulvar melanoma between 01/2011 and 12/2020 at University Hospital 12 de Octubre, Madrid, Spain.

Result(s)* A total of 7 women were included. The mean age at diagnosis was 74.7 ± 9.7 years. All women consulted for a mass located in genital tract mainly at labia minora. In all

patients gynecologic examination revealed pigmented lesion from which a biopsy was taken. Histologic examination indicated the diagnose of a melanoma. The mean tumor's size was 3.5 ± 2.1 cm with clinically negative nodes and a mean Breslow depth 11.2 ± 8.7 mm. PET-CT was solicited as pre-surgical imaging workup.

According to the AJCC Cancer Staging Manual, 3/7 (42.8%) of patients were diagnosed at IIC stage, 1/7 (14.3%) at IIIB stage, 2/7 (28.6%) at IIIC stage and 1/7 (14.3%) at IV stage. The treatment was surgery: 3/7 (42.8%) of patients underwent wide local excision and 4/7 (57.2%) of patients underwent hemivulvectomy. Negative margins were obtained in 4/7 (57.2%) of surgeries. Sentinel lymph node (SLN) biopsy was performed in 5/7 (71.4%) of patients and SLN confirmed nodal metastases in 4/5 (80%) of patients. Immunotherapy was the adjuvant treatment for patients with $\geq$ IIIB stage (4/7), one of them combined with radiotherapy.

During the follow up, 5/7 (71.4%) of patients relapsed. The mean time to relapsed was 5.5 ± 2.3 months. The treatment for the recurrences was 3/5 (60%) cytoreductive surgery combined with immunotherapy in two patients. One woman (20%) received a combination of radiotherapy and immunotherapy and other woman (20%) received palliative treatment. At the time of last follow up, 4/7 (57.2%) women died because of melanoma, 2/7 (28.6%) was tumor free and 1/7 (14.3%) was alive with the disease.

Conclusion* As reported in literature, primary vulvar melanomas have biologically aggressive characteristics. The treatment consists of a surgery with appropriate free surgical margins. However, early recognition is what brings the maximal benefit to survival.

1021

V-Y FLAP RECONSTRUCTION FOR VULVAR CANCER RECURRENCE AFTER PRIMARY RADIOTHERAPY

¹S Fernandez-Gonzalez*, ¹C Ortega, ¹L Marti, ²JO Bermejo, ²JA Palacin, ³Y Perez, ³S Martorell, ³M Alvarez, ¹M Barahona, ¹J Ponce. ¹University Hospital of Bellvitge (IDIBELL), Gynecology, L'Hospitalet de Llobregat, Spain; ²University Hospital of Bellvitge (IDIBELL), Plastic and reconstructive surgery, L'Hospitalet de Llobregat, Spain; ³University Hospital of Bellvitge (IDIBELL), Nursery, L'Hospitalet de Llobregat, Spain

10.1136/ijgc-2021-ESGO.628

Introduction/Background* Vulvar cancer is a rare disease specially in young women. When a large tumour is diagnosed, reconstruction flap should be considered in order to achieve the best cosmetic and functional result.

Methodology This surgical video shows the excision of a perianal recurrence secondary to a vulvar cancer after primary radiotherapy. Rectal amputation and terminal colostomy were performed through a laparoscopic approach. A bilateral V-Y flap was designed for perineal reconstruction. This procedure was recorded step by step.

Result(s)* A 47 years old woman was diagnosed with perianal VIN III. It measured 25mm and had no extraperineal disease. Primary radiotherapy was indicated due to the invasion of anal mucosa. After 6 months of treatment, perianal recurrence of squamous vulvar carcinoma was diagnosed. Consequently, wide surgical excision, anal amputation, bilateral V-Y flap reconstruction and terminal colostomy were performed under lithotomy position and general anaesthesia. Despite surgical wound infection by pseudomonas and partial dehiscence, very good cosmetic and functional results were achieved.

Abstract 1021 Figure 1

Conclusion* V-Y flap reconstruction may be a good option for the treatment of perineal defects that are too large for primary intention wound closure.

1048

TRANSINGUINAL PELVIC LYMPHADENECTOMY IN VULVAR CANCER PATIENTS (TRIPLE STUDY): A SINGLE-INSTITUTION, PROSPECTIVE, PILOT STUDY

¹SM Fragomeni*, ²L Della Corte, ¹C Conte, ¹A Federico, ³B Marinucci, ⁴L Tagliaferri, ⁵S Gentileschi, ¹G Corrado, ⁶I Romito, ³A Zampolini Faustini, ¹LC Turco, ¹V Gallotta, ^{1,3}A Fagotti, ^{1,3}G Scambia, ^{3,6}G Garganese. ¹Fondazione Policlinico Universitario A. Gemelli IRCCS, Unità di Ginecologia Oncologica, Dipartimento Scienze della Salute della Donna, del Bambino e di Sanità Pubblica, Rome, Italy; ²University of Naples Federico II, Department of Neuroscience, Reproductive Sciences and Dentistry, School of Medicine, Naples, Italy; ³Università Cattolica del Sacro Cuore, Istituto di Clinica Ostetrica e Ginecologica, Rome, Italy; ⁴Fondazione Policlinico Universitario A. Gemelli IRCCS, Unità Operativa Complessa di Radioterapia, Dipartimento di Scienze Radiologiche, Radioterapiche ed Ematologiche, Rome, Italy; ⁵Fondazione Policlinico Universitario A. Gemelli IRCCS, Unità di Chirurgia Plastica, Dipartimento Scienze della Salute della Donna, del Bambino e di Sanità Pubblica, Rome, Italy; ⁶Mater Olbia Hospital, Gynecology and Breast Care Center, Olbia, Italy

10.1136/ijgc-2021-ESGO.629

Introduction/Background* Vulvar cancer (VC) is a lymphotropic disease. The preferred lymphatic pathway involves the inguinofemoral lymph nodes (IFLNs), followed by the pelvic lymph nodes (PLNs). In selected cases, surgical approach to PLNs may be the first choice when histologic confirmation is required for suspicious involvement on imaging or when radiation therapy is out of indication (e.g., in cases of previous radiation therapy or in specific histotypes).

Given the anatomic continuity between the inguinal and pelvic sites, a novel retrograde trans-inguinal pelvic access was explored to provide concomitant lymphadenectomy, with a single skin incision and no change in surgical position or instruments.

Our objective was to evaluate the feasibility and safety of the extraperitoneal trans-inguinal novel approach to pelvic lymphadenectomy (TRIPLE).

Methodology All consecutive patients referred to our Institution in the last 18 months, affected by primary/recurrent VC, candidate to concomitant groin and pelvic lymph node surgery were included. After conventional IFLN dissection, ipsilateral extraperitoneal trans-inguinal pelvic lymphadenectomy (TRIPLE) was performed. Clinical data, type of treatment,