Methodology A 32-item questionnaire addressing multiple issues related to the SUCCOR study and surgical management of ECC was designed and sent from the ESGO account through a newsletter on February 23rd to all ESGO members from 54 countries of the ESGO directory. The survey was completely confidentially and could be completed in less than 5 minutes. Responses from 353 members were obtained. Descriptive statistical analysis and frequency tables, as well as two-sided Fisher's exact test, were used.

Result(s)* 92.9% of responders were aware of the findings of the SUCCOR study. 73.4% expected the results obtained. 5 out of 10 of members confirmed that have changed their practice after the publication of the manuscript.

Currently, 34.8% of responders admit that they usually perform MIS (laparoscopy/robotic) for cervical tumours smaller than 2 cm, while for tumours larger than 2 cm (FIGO IB2) 15.1% state that they continue to carry out MIS. Between 63 and 64.1% apply protective maneuvers during this procedure. In ECC who have undergone conization, 56.1% would performed MIS for tumours smaller than 2 cm while only 30% would offer it for tumours larger than 2 cm.

Finally, 84.1% of ESGO members consider ethically acceptable the development of a new European prospective observational study (Image 1) to evaluate the surgical approach in ECC (IB1-IB2).

Conclusion* According to the ESGO members' perspectives regarding the approach to ECC, it seems that MIS still has an important role to play, which should be evaluated in a new prospective study that reflects the European reality (Image 2).

785

INCIDENCE OF CERVICAL CANCER AND MANAGEMENT OF HIGH GRADE SMEARS IN WOMEN AGED 50 YEARS AND ABOVE

R Shah*, E Leung, D Mukhopadhyay. *Queen's Hospital, UK*

10.1136/ijgc-2021-ESGO.69

Introduction/Background* Interpretation of smears is a challenge in older women owing to atrophic changes. Colposcopy can be difficult and views are repeatedly unsatisfactory due to a stenotic os. LLETZ procedures are therefore often performed based on initial high-grade smear cytology. However, approximately 60% of cervical cancers occur in women aged 45 plus; evidence shows a second peak of high-risk HPV in postmenopausal women. This study aims to review the management of women aged 50 years and above with high-grade smears and establish the incidence of cervical cancer in this cohort.

Methodology A retrospective study was conducted; an electronic search was undertaken to identify all women aged 50 years and above referred from the national screening programme with a high grade (moderate or severely dyskaryotic) smear to the Colposcopy unit at Queen's Hospital & King George Hospital Barking, Havering & Redbridge NHS Trust between 1st January 2016 – 31st December 2019. Inflex and cyberlab were used to collate data to establish colposcopy findings, histology of biopsy, LLETZ, or further surgical intervention plus results following tests of cure. Data was analysed using Microsoft Excel.

Result(s)* Of the 99 women referred, smear cytology demonstrated 1 suspicious of glandular neoplasia, 50 of severe dyskaryosis and 48 of moderate dyskaryosis. 11 patients were

excluded due to incomplete data. Colposcopic views were undiagnostic for 27 (31%) of patients. 82 (93%) patients underwent LLETZ, while 11% of patients underwent radical surgery based on their smear +/- histology results. This study identified a 5% incidence of cancer (3 SCC; 1 adenocarcinoma). 61% exhibited high-grade histology (23% CINII and 38% CINIII) and 13% exhibited low-grade histology (CINI). 11% had no abnormality and a further 10% displayed other benign changes.

Conclusion* Our local analyses demonstrate a 5% incidence of cancer in women aged 50 years as per histological diagnosis. 61% of referred patients exhibited high-grade histology; 11% of the sample population underwent radical surgery. This may justify a lower threshold to performing LLETZ procedures where colposcopy findings are indeterminate.

787

EARLY OUTCOMES OF NEOADJUVANT CHEMOTHERAPY AND SUBSEQUENT RADIATION THERAPY IN NON METASTATIC ADVANCED CERVICAL CANCER

¹R Mittal, ¹L Gurram, ¹C Johny, ²S Gupta, ²J Ghosh, ²S Gulia, ¹S Chopra, ¹R Engineer, ³ST S, ³A Maheshwari, ²S Rath, ³P Poddar, ⁴P Popat, ¹D A, ⁴U Mahantshetty. ¹TATA MEMORIAL HOSPITAL, Homi Bhabha Block, Radiation Oncology, Mumbai, India; ²TATA MEMORIAL HOSPITAL, Homi Bhabha Block, Medical Oncology, Mumbai, India; ³TATA MEMORIAL HOSPITAL, Homi Bhabha Block, Surgical Oncology, Mumbai, India; ⁴TATA MEMORIAL HOSPITAL, Homi Bhabha Block, Radiodiagnosis, Mumbai, India

10.1136/ijgc-2021-ESGO.70

Introduction/Background* To report outcomes of non metastatic advanced cervical cancer patients treated with neoadjuvant chemotherapy (NACT) followed by chemoradiation/radiation.

Methodology 104 patients treated between January 2015 and December 2019 were analyzed and restaged as per FIGO 2018. RECIST criteria v1.1 was used to assess response to NACT and toxicity was assessed using CTCAE v4.0. Survival analysis was done using Kaplan-Meier method. Univariate and multivariate analysis was done using Log-rank test and Cox-regression model respectively.

Result(s)* 104 patients (median age: 52 years) of advanced cervical cancer (92% SCC) received NACT for following reasons: bulky pelvic nodes (n=20), bulky para-aortic nodes (n=10), bulky pelvic + para-aortic nodes (n=31), bladder invasion (n=43). Baseline median pelvic and para-aortic nodal size was 2.5 cm (range: 1-8) and 1.8 cm (range: 1-3.5) respectively. Baseline median tumor infiltration size into bladder on cystoscopy was 3 cm (range 0.5-7). NACT regimen was Paclitaxel (175 mg/m²) and Carboplatin (AUC 5) 3 weekly IV infusion for 3 (median) cycles (range: 1-6). Post NACT completion, 9 (8.6%) defaulted from subsequent treatment, 8 (7.7%) were labeled as best supportive care, 1 received palliative chemotherapy for progressive disease and 13 (12.5%) patients received palliative RT. 73 (70.2%) patients received radical pelvic EBRT (median dose of 46Gy/23Fr) of which 68 (93.1%) received brachytherapy boost (median dose 21Gy/3Fr). 32 (43.8%) and 28 (38.3%) patients also received extended field RT (EFRT) and nodal boost (median dose 5.4Gy/3Fr) respectively. 64 (87.6%) patients received concurrent cisplatin. On univariate analysis, patients with comorbidities, bladder invasion, pelvic nodal disease, NACT dose reduction, incomplete radical treatment post-NACT and pelvic nodal size ≥ 2.5cm at baseline had poor outcomes (table 1). On multivariate

Abstract 787 Figure 1 Kaplan meier curves for patients who completed NACT followed by radical local treatment (CTR/RT + brachytherapy boost)

Abstract 787 Table 1 Univariate analysis of correlation of prognostic factors with LC, RC, DMC, RFS and OS

Variables		Local Control	Regional Control	Distant Metastasis Control	Disease Free Survival	Overall Survival
Age (years)	<52	0.260	0.482	0.937	0.623	0.792
	≥ 52					
Comorbidities	No	0.590	0.019	0.942	0.349	0.139
	Yes					
Stage	IIIC1	0.109	0.826	0.196	0.944	0.671
	IIIC2					
	IVA					
Reason for giving NACT	Bladder invasion	0.012	0.750	0.076	0.986	0.701
	Others					
Nodal Site	Pelvic nodes	0.503	0.037	0.088	0.041	0.110
	Others					
Maximum Pelvic node size (cm)	< 2.5	0.911	0.234	0.137	0.232	0.007
	≥ 2.5					
Maximum Para-aortic node size (cm)	< 1.8	0.996	0.669	0.121	0.459	0.507
	≥ 1.8					
NACT dose reduction	No	0.001	0.061	0.386	0.003	0.022
	Yes					
Post NACT	Completed radical treatment	0.000	0.000	0.436	0.000	0.000
	Not completed radical treatment					
Nodal Boost	No	0.143	0.986	0.259	0.939	0.642
	Yes					

analysis, none of the variables reached levels of significance. At median follow-up of 26 months (IQR: 11-45), 34 patients had died of disease, 51 were alive, of which 34 were disease-free. The 2 year LC, RC, DMC, DFS and OS for the entire cohort was 75%, 68%, 72%, 48% and 60% respectively whereas for the patents who completed NACT and radical local treatment was 92%, 82%, 76%, 62% and 75% respectively (figure 1).

Conclusion* NACT followed by radiation therapy in non metastatic advanced cervical cancer is a feasible option with good survival outcomes and acceptable toxicity.

820

LATERALLY EXTENDED PELVIC RESECTION WITH COMPLETE RESECTION OF URETER AND NEPHRECTOMY

M Zikan*, O Dubova, P Koliba, V Student, T Brtnicky, P Kabele. *Charles University – First Faculty of Medicine and Bulovka University Hospital, Dept. of Gynecology and Obstetrics, Praha 8, Czech Republic*

10.1136/ijgc-2021-ESGO.71

Introduction/Background* Laterally extended endopelvic resection became a standard surgical approach for pelvic tumors, recurrences specifically, localized laterally in the pelvis. This