EPV208/#485

THE CLINICAL DEMAND FOR PRESSURIZED INTRAPERITONEAL AEROSOL CHEMOTHERAPY IN SOUTH KOREA: AN ELECTRONIC SURVEY-BASED STUDY

¹EJ Lee*, ¹SJ Park, ²J Mun, ²H Paik, ²J Lee, ²A Seol, ²J Kim, ³N Lee, ⁴GW Yim, ⁵S-H Shim, ¹HS Kim, ⁶S-J Chang. ¹Seoul National University College of Medicine, Obstetrics and Gynecology, Seoul, Korea, Republic of; ²Seoul National University College of Medicine, Department of Obstetrics and Gynecology, Seoul, Korea, Republic of; ³CHA Gangnam Medical Center, Obstetrics and Gynecology, Seoul, Korea, Republic of; ⁴Dongguk University Ilsan Hospital, Department of Obstetrics and Gynecology, Goyang, Korea, Republic of; ⁵Konkuk University School of Medicine, Obstetrics and Gynecology, Seoul, Korea, Republic of; ⁶Ajou University School of Medicine, Obstetrics and Gynecology, Suwon, Korea, Republic of

10.1136/ijgc-2021-IGCS.279

Objectives Pressurized intraperitoneal aerosol chemotherapy (PIPAC) is effective for treating solid tumors with peritoneal metastasis. However, PIPAC is not a standard treatment globally and is currently only used in the limited areas. Thus, we performed a survey of surgical oncologists related to PIPAC to evaluate the clinical desire for PIPAC in South Korea, one of the many countries where PIPAC has not yet been introduced. **Methods** We performed an online survey of Korean surgical oncologists between November and December 2019. The questionnaire consisted of 20 questions related to PIPAC, which were divided into comprehensive inquiry (5 questions), procedure inquiry (13 questions), and cost inquiry (2 questions).

Results A total of 164 respondents answered the questionnaire. Among all respondents, 41.7–50% majoring in ovarian cancer, pseudomyxoma peritonei, and malignant mesothelioma preferred PIPAC for the curative treatment of primary diseases, whereas 32.7–33.3% majoring in colorectal and hepatobiliary cancers chose PIPAC for the palliative treatment of recurrent diseases. Moreover, 66.7–95.2% of the respondents considered PIPAC appropriate for the cancers the specialized in. About 70% expected a treatment response of more than 50% through the repeated implementation of PIPAC under general anesthesia. Respondents also considered grade 1 or 2 minor surgical complications an acceptable risk. Finally, respondents considered the reasonable costs to purchase and implement PIPAC once at between 1,000,000–5,000,000 KRW.

Conclusions Although the treatment scope for applying PIPAC was different among Korean surgical oncologists, most of them expected relatively high tumor response rates with minor toxicities through the repeated implementation of PIPAC.

EPV209/#49

SURVIVAL IN CASE OF CARDIOPHRENIC LYMPHADENOPATHY IN ADVANCED STAGE EPITHELIAL OVARIAN CANCER PATIENTS WHO UNDERWENT CYTOREDUCTIVE SURGERY; A SYSTEMATIC REVIEW AND META- ANALYSIS

¹M Kengsakul*, ²G Nieuwenhuijzen-De Boer, ³A Bijleveld, ⁴S Udomkarnjananun, ⁵S Kerr, ⁶C Niehot, ²H Van Beekhuizen. ¹Panyanantaphikku Chonprathan Medical Center, Srinakharinwirot University, Gynecologic Oncology, Nonthaburi, Thailand; ²Erasmus MC Cancer institute, University Medical Center Rotterdam, Gynecologic Oncology, Rotterdam, Netherlands; ³Albert Schweitzer Hospital, Obstetrics and Gynecology, Dordrecht, Netherlands; ⁴King Chulalongkorn Memorial Hospital, Chulalongkorn University, Department of Medicine, Bangkok, Thailand; ⁵King Chulalongkorn Memorial Hospital, Chulalongkorn University, Research Affairs, Bangkok, Thailand; ⁶Erasmus MC Cancer institute, University Medical Center Rotterdam, Medical Library, Rotterdam, Netherlands

10.1136/ijgc-2021-IGCS.280

Objectives Favorable survival outcomes for patients with advanced stage epithelial ovarian cancer (ASEOC) is associated with complete cytoreduction. In this meta-analysis we evaluate the therapeutic role of cardiophrenic lymph nodes (CPLNs) resection ASEOC who have undergone cytoreductive surgery.

Methods Embase, Medline, Web of science, Cochrane Library and Google scholar were searched for articles published in English from their inception to November 2020. Meta-analysis was conducted to determine the prognostic impact of surgical outcome, postoperative complications and survival.

Results Fifteen relevant articles, 727 patients with CPLNs adenopathy and 981 patients without CPLNs adenopathy, were analyzed. Higher percentage of ascites, intra and extra abdominal metastases was observed in CPLNs adenopathy group. The mean size of pre-operative CPLNs was 9.1 ± 3.75 mm. Eighty-two percent of enlarged CPLN were histological confirmed. No difference in surgical outcome and perioperative complication was observed between both groups. Meta-analysis showed that patients with CPLNs adenopathy had a significantly increased risk of disease recurrence (OR 4.56, 95% CI 1.98–10.51, $P < 0.001$) and dying from disease (OR 2.96, 95% CI 2.08– 4.22, $p < 0.001$) in comparison to those without CPLNs adenopathy.

Conclusions Patients with CPLNs adenopathy had higher tumor burden intra and extra-abdominally and decreased survival compared to patients without CPLNs adenopathy. There is not enough available data to confirm the therapeutic role of CPLNs resection. Therefore, a randomized controlled trial should be conducted to demonstrate the benefit of CPLNs resection in cytoreductive surgery.

EPV210/#490

PROGNOSTIC IMPLICATIONS OF HEMODYNAMIC INSTABILITY DURING OVARIAN CANCER SURGERY

¹J Kim*, ¹SI Kim, ¹A Seol, ¹HS Kim, ¹HH Chung, ¹J-W Kim, ¹NH Park, ¹Y-S Song, ^{1,2}M Lee. ¹Seoul National University College of Medicine, Department of Obstetrics and Gynecology, Seoul, Korea, Republic of; ²Seoul National University Hospital, Department of Obstetrics and Gynecology, Seoul, Korea, Republic of

10.1136/ijgc-2021-IGCS.281

Objectives To evaluate the impact of intraoperative hypotension and hemodynamic instability on survival outcomes in patients with high-grade serous ovarian carcinoma (HGSOC).

Methods We retrospectively identified patients with HGSOC, who underwent primary or interval debulking surgery between August 2013 and December 2019. We collected anesthesia-related variables, including the arterial blood pressure measurements (at 1-min interval) during surgery of patients. The cumulative duration of mean arterial blood pressure (MAP) readings under 65 mmHg and two performance measurements (median performance error [MDPE] and wobble) were calculated. We investigated associations between the factors indicating hemodynamic instability and prognosis.

Results In total, 338 patients were included. Based on the cumulative duration of MAP under 65 mmHg, we divided patients into two groups: ≥ 30 min and < 30 min. The progression-free survival (PFS) was worse in the ≥ 30 min group ($n=107$) than the < 30 min group ($n=231$) (median, 18.2 vs.