p53 immunohistochemistry on 39.4% with a sensitivity of 98.5% to detect p53abn (99.6% negative predictive value). Cytologic features including tumor giant cells, smudged chromatin, cherry-red/macronucleoli, and atypical mitoses accurately predicted p53abn. In 7/292, p53abn upgraded ESGO risk groups (2 to intermediate-risk, 5 to high-risk). EEC12/stage IA patients had an excellent cause-specific 5-year survival of 98.5%.

Conclusions Pathologists can select cases for p53 testing with high sensitivity and low risk of false negativity. Molecular characterization of endometrial carcinomas has great potential to refine ESGO risk classification for a small subset but offers little value for approximately half of endometrial carcinomas, namely, EEC12/stage IA.

EPV104/#228

EMERGING IMMUNOTHERAPY PARADIGMS IN ADVANCED ENDOMETRIAL CANCER: THE EFFECT OF ONLINE EDUCATION ON CLINICIAN KNOWLEDGE AND CONFIDENCE

¹G Fisher*, ¹A Furedy, ¹J Vandenbroucq, ²N Colombo. ¹Medscape, Medical Education, London, UK; ²Instituto Europeo di Oncologia, Department of Gynecology, Milan, Italy

10.1136/ijgc-2021-IGCS.174

Objectives This study determined whether online continuing medical education (CME) could improve the knowledge of oncologists (oncs) and obstetricians/gynaecologists (obs/gyns) regarding the rationale and evidence for immunotherapy paradigms in advanced endometrial cancer.

Methods A 30-minute online video lecture was launched for physicians outside the USA August 2020 with data collected to November 2020. Educational effect assessed with repeated-pairs pre-/post-activity- individual participants serving as own control. 3 multiple-choice, knowledge questions and 1 self-efficacy, 5-point Likert scale confidence question were analyzed. Chi-squared test assessed pre- to post-activity change (5% significance level, $P < .05$). Magnitude of change in total number of correct responses overall, and for each question, determined with Cramer's V ($< .06$ =Modest, 0.06 – 0.15 =Noticeable, 0.16 – 0.26 =Considerable, $> .26$ =Extensive).

Results 142 obs/gyns and 60 oncs completed pre- and post-activity questions. Positive educational effect was observed for obs/gyns (noticeable effect, $V = .092$, $P < .01$; average% of correct responses increasing from 33 to 42%) and oncs (noticeable effect, $V = .150$, $P = .0043$; average% of correct responses increasing from 47 to 62%). Increases in correct responses post-activity seen for questions on response to 2nd line chemotherapy (% relative improvement, obs/gyn: 23%, oncs: 22%), rationale for immunotherapy (obs/gyns: 24%, oncs: 72%), data for the dostarlimab GARNET trial (obs/gyns: 36%, oncs: 21%). Confidence in knowledge of the evidence for immunotherapy strategies increased post-activity (total average confidence shift: 27% obs/gyns and 40% oncs). Overall, 22% of learners' responses were improved and 39% of learners' responses were reinforced.

Conclusions This online CME activity resulted in a positive educational impact for both clinical specialties. However, education gaps remained evident post-activity.

EPV105/#237

COMBINATION TARGETED TREATMENT WITH MEK AND PAN-ERBB INHIBITORS ENHANCES ANTITUMOR ACTIVITY IN ERBB AMPLIFIED EX-VIVO SEROUS ENDOMETRIAL CANCER CELLS

¹A Larish*, ²W-H Lin, ³J Smadbeck, ³S Sotiriou, ³J Cheville, ¹F Harris, ¹R Feathers, ¹J Lynch, ⁴A Kumar, ⁵M Block, ¹L Jones, ¹G Karagouga, ¹A Mccune, ⁵A Mansfield, ³S Murphy, ⁴A Mariani, ³G Vasmatazis, ²P Anastasiadis, ⁵J Weroha. ¹Mayo Clinic, Obstetrics and Gynecology, Rochester, USA; ²Mayo Clinic, Department of Laboratory Medicine and Pathology, Jacksonville, USA; ³Mayo Clinic, Department of Laboratory Medicine and Pathology, Rochester, USA; ⁴Mayo Clinic, Division of Gynaecologic Surgery, Department of Obstetrics and Gynaecology, Rochester, USA; ⁵Mayo Clinic, Medical Oncology, Rochester, USA

10.1136/ijgc-2021-IGCS.175

Objectives ERBB pathway alterations present therapeutic targets in high grade endometrial cancer (EC), but efficacy can be limited by persistent co-activation of other ERBB binding partners. The efficacy of dual-inhibition MEK+pan-ERBB or BET+pan-ERBB in an ERBB2/ERBB3 amplified EC was investigated via 3D microcancer ex-vivo cell assay.

Methods Tumor was prospectively collected from a patient with stage IIIc1 serous EC. Whole exome, mRNA, and Mate-Pair genomic characterization was performed. Tumor cells were grown in 3D culture and subjected to titrating drug treatments. Cell viability was determined by the CellTiter-Glo Luminescent Assay. Data transformation and dose-response curves were generated using GraphPad PRISM using the variable slope model. CalcuSyn software with the Chou-Talalay method analyzed drug interactions and synergy. Afatinib, binimetanib, and JQ1 were used to inhibit pan-ERBB, MEK1/2, BET, respectively. For translational relevance, inhibitory effect was defined as percent reduction in ATP from baseline at the

Abstract EPV105/#237 Figure 1 Microcancer ex vivo exposure to MEK+pan-ERBB inhibitors. Dose response curves of single and combination treatments (left) were 10-fold titrated across 8 log doses for each agent. The highest concentration (i.e. fraction of full (FoF)=1) of afatinib, binimetanib and afatinib+JQ1 was 3 μ M, 10 μ M, and 50 μ M, respectively. The physiologically achievable concentration of afatinib is indicated (dotted lines). A combination index (CI, right) was used to assess synergy with afatinib+binimetanib and afatinib+JQ1 as shown by Fa-CI plots.

Abstracts

physiologically achievable concentration (maximum plasma concentration (C_{max}) value).

Results Sequencing revealed amplifications of ERBB2 (17q12), RAF1, c-myc, and ERBB3 (12q13.2) low-level gain. Inhibition of viability was moderate by single agents: Afatinib, binimetanib, JQ1, as shown by inhibitory effect values of 14.4%, 47.8%, 8.8%, respectively at physiologically achievable concentrations (C_{max}) of afatinib. Combinations demonstrated increasing inhibitory effect values: 99.7% for Afatinib+ binimetanib, and 99.5% for Afatinib+JQ1. Synergy was evidenced for both combinations by a combination index <1 (figure 1).

Conclusions Combined inhibition of pan-ERBB with inhibition of MEK or BET proteins synergistically suppress viability in patient-derived serous EC harboring ERBB amplifications.

EPV106/#249

ENDOMETRIAL CANCER IMMUNOHISTOCHEMICAL RISK STRATIFICATION IN A LARGE UTERINE-CONFINED CANCER SERIES

¹E Perrone*, ²F De Felice, ³I Capasso, ⁴D Arciuolo, ⁵E Distefano, ¹D Lorusso, ⁴GF Zannoni, ¹G Scambia, ¹F Fanfani. ¹Fondazione Policlinico Universitario A. Gemelli, IRCCS, Uoc Ginecologia Oncologica, Dipartimento Per La Salute Della Donna E Del Bambino E Della Salute Pubblica, Rome, Italy; ²Policlinico Umberto I, 'Sapienza' University of Rome, Department of Radiotherapy, Rome, Italy; ³Policlinico universitario Agostino Gemelli, Uoc Ginecologia Oncologica, Roma, Italy; ⁴Fondazione Policlinico Universitario A. Gemelli, IRCCS, Gyneco-pathology and Breast Pathology Unit, Dipartimento Per La Salute Della Donna E Del Bambino E Della Salute Pubblica, Rome, Italy

10.1136/ijgc-2021-IGCS.176

Objectives The aim of this study is to assess the clinical reproducibility and the oncological validity of the Endometrial cancer (EC) risk stratification based on the molecular information given by the immunohistochemistry (IHC).

Methods Retrospective IHC analyses were conducted in a large series of 778 pre-operative uterine-confined ECs, studying the presence/absence of MLH1, MSH2, MSH6, to define the mismatch repair (MMR) stable or unstable phenotype; the presence of p53 mutations and other molecular features. The molecular profile was correlated with histological, clinical and prognostic EC patients' data.

Results Based on the IHC, we defined 3 EC populations: MMR stable (MMRs), unstable (MMRi) and p53 mutated (p53+) patients. Our result demonstrated that the IHC stratification statistically correlated with the most relevant anatomical features: FIGO stage (p<0.001), grading (12.5% G3 in MMRs vs 22.9% in MMRi vs 95.3% in p53+, p<0.001), histotype (Type II 6.2% in MMRs vs 5.3% in MMRi vs 87.5% in p53+, p<0.001), presence of LVSI (positive in 16.3% in MMRs vs 23.8% in MMRi vs 38.7% in p53+,

p<0.001), myometrial invasion and tumor dimension (p=0.003 and p<0.001 respectively). Again, the 3 IHC populations statistically reflected the EC risk class ESGO-ESMO-ESP classification 2020 (p<0.001). These results were confirmed also in Kaplan-Meier curves in terms of over-all survival (OS) and disease-free survival (DFS) (p<0.001) (figure 1).

Conclusions In this large series, we demonstrated that the pragmatic and systematic use of IHC may have an important role to properly stratify, in terms of histological features and clinical outcome, the uterine-confined EC patients.

EPV107/#258

ENGOT-EN11/GOG-3053/KEYNOTE-B21: PHASE 3 STUDY OF PEMBROLIZUMAB OR PLACEBO IN COMBINATION WITH ADJUVANT CHEMOTHERAPY WITH/WITHOUT RADIOTHERAPY IN PATIENTS WITH NEWLY DIAGNOSED HIGH-RISK ENDOMETRIAL CANCER

¹T Van Gorp*, ²M Mirza, ³A Lortholary, ⁴I Vergote, ⁵D Cibula, ⁶A Walther, ⁷A Savarese, ⁸M-P Barretina-Ginesta, ⁹F Ortaç, ¹⁰C Papadimitriou, ¹¹L Bodnar, ¹²C-H Lai, ¹³K Hasegawa, ¹⁴X Xie, ¹⁵EL Barber, ¹⁶RL Coleman, ¹⁷S Keefe, ¹⁷R Orlowski, ¹⁸B Slomovitz. ¹UZ Leuven, Gynaecological Oncology, Leuven, Belgium; ²NSGO-CTU and Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark; ³Centre Catherine de Sienne, Hôpital Privé Du Confluent, Nantes, France; ⁴BGOG and University Hospitals Leuven, Leuven Cancer Institute, Leuven, Belgium; ⁵Department of Obstetrics and Gynecology, General Faculty Hospital In Prague, First Faculty of Medicine, Charles University, Prague, Czech Republic; ⁶Bristol Cancer Institute, University Hospitals Bristol, Bristol, UK; ⁷Department of Medical Oncology, Istituto Nazionale Tumori Regina Elena, Rome, Italy; ⁸Catalan Institute of Oncology and Girona Biomedical Research Institute, Medical School University of Girona, Girona, Spain; ⁹Department of Obstetrics and Gynecology, Ankara University School of Medicine, Ankara, Turkey; ¹⁰Aretaieio University Hospital, National and Kapodistrian University of Athens, Athens, Greece; ¹¹Department of Oncology and Immunooncology, Warmian-masurian Cancer Center of The Ministry of The Interior and Administration's Hospital, Olsztyn, Poland; ¹²TGOG and Department of Gynecology and Obstetrics, Chang Gung Memorial Hospital, Linkou Branch, Taoyuan, Taiwan; ¹³Department of Gynecologic Oncology, Saitama Medical University, Hidaka, Saitama Prefecture, Japan; ¹⁴Women's Hospital, School of Medicine, Zhejiang University, Hangzhou, China; ¹⁵Department of Gynecologic Oncology, Northwestern University Feinberg School of Medicine, Chicago, USA; ¹⁶Department of Gynecologic Oncology, Us Oncology Research, The Woodlands, USA; ¹⁷Oncology, Merck and Co., Inc., Kenilworth, USA; ¹⁸Department of Gynecologic Oncology, Broward Health, Fort Lauderdale, USA

10.1136/ijgc-2021-IGCS.177

Objectives Pembrolizumab, an anti-PD-1 antibody, has demonstrated activity as monotherapy and in combination with lenvatinib in patients with previously treated mismatch repair (MMR) deficient and MMR proficient endometrial cancer (EC). ENGOT-en11/GOG-3053/KEYNOTE-B21 (NCT04634877) is a phase 3, randomized, double-blind study of pembrolizumab or

Abstract EPV106/#249 Figure 1