EPV154/#117

HIV TESTING IN CERVICAL DYSPLASIA, PRACTITIONERS' OPINION

¹R van de Laar*, ²C Jordans, ³C Rokx, ⁴H van Beekhuizen, ⁵HC van Doorn. ¹Erasmus MC Cancer Institute, Gynaecological Oncology, Rotterdam, Netherlands; ²Erasmus MC, Microbiology and Infectious Diseases, Rotterdam, Netherlands; ³Erasmus MC, Internal Medicine, Section Infectious Diseases, Rotterdam, Netherlands; ⁴Erasmus MC Cancer Institute, University Medical Center Rotterdam, Gynecologic Oncology, Rotterdam, Netherlands; ⁵Erasmus MC Cancer Institute, Gynecologic Oncology, Rotterdam, Netherlands

10.1136/ijgc-2021-IGCS.224

Objectives Cervical dysplasia is an HIV indicator condition and according international recommendations HIV testing is strongly advised in women with cervical dysplasia, because the risk of an undiagnosed HIV is thought to be >0.1%. Therefore an HIV test should be offered to all women with cervical dysplasia. There is no literature about the opinion of Gynaecologist on HIV screening in patients with cervical dysplasia.

Methods We sent an online questionnaire to gynecologist in South West Netherlands to investigate 1) what they know about this issue, 2) their opinion and willingness on active HIV testing for this cervical dysplasia.

Results The questionnaire was sent to 103 gynaecologists of whom fifty-six participants replied (54%). Forty-eight (86%) think patients are not offended when HIV testing is offered and 50 (89%) have no difficulty to address HIV testing. Thirty-nine (70%) gynaecologist think that the prevalence of undiagnosed HIV infection is lower than 0.1%, and only

Abstract EPV154/#117 Figure 1

Abstract EPV155/#210 Figure 1 Age-specific incidence of cervical cancer in US and Taiwan, 2001-2017

seven (12,5%) accept HIV testing in case of a prevalence of 0.1% or less. Thirty-two (57%) are willing to test with a prevalence of 1% or higher.

Conclusions To address and offer HIV testing seems not an issue for the gynaecologists questioned in our study. However, the willingness to routinely perform an HIV test for cervical dysplasia at the assumed 0.1% prevalence looks insufficient and differs from the recommendations of international policy makers. Discussion is needed to change the threshold or the willingness for testing.

EPV155/#210

DISPARITIES IN CERVICAL CANCER INCIDENCE IN NATIVE ASIANS VS. US ASIANS – A POPULATION ANALYSIS

¹A Moon*, ²C-I Liao, ³E Thayer, ⁴J Chan. ¹Stanford University School of Medicine, Obstetrics and Gynecology, Division of Gynecologic Oncology, Stanford, USA; ²Kaohsiung Veterans General Hospital, Obstetrics and Gynecology, Kaohsiung City, Taiwan; ³University of Massachusetts Medical School, Obstetrics and Gynecology, Worcester, USA; ⁴California Pacific Medical Center, Obstetrics and Gynecology, San Francisco, USA

10.1136/ijgc-2021-IGCS.225

Objectives To evaluate the incidence and trends of cervical cancer in native Asians compared to Asians in the United States.

Methods Data were obtained from Taiwan Cancer Registry in Health and Welfare Data Center and United States Cancer Statistics between 2001 and 2017. SEER*Stat 8.3.8, Joinpoint regression program 4.8.0.1, Microsoft Excel calculated the age-adjusted incidence (AAI, per 100,000 women), age-specific incidence (ASI, per 100,000 women), and trends (average annual percent change, AAPC).

Results Compared to US Asians, native Asians had a significantly higher cervical cancer incidence at 7.8 vs. 5.1 per 100,000. Over time, the incidence in Taiwan is improving at a rate of 6% per year but remains high. Based on age groups, the incidence increased at a younger age in Whites (25–29 years) compared to an older age in US Asians (30–34 years) and native Asians (35–39 years). Although new cases peaked in Whites and US Asians after age 40 and then plateaued in the older age groups, native Asians continued to have an increase into age 80. In fact, the incidence of cancer in native Asians age 85 and older was 53.6 vs. 12.7 in US Asians, a four-fold difference.

Conclusions After age 40, cervical cancer incidence was increasing every 5-age years in native Asians while plateauing in the US. Native Asians aged 85+ years had a four-fold higher incidence of cervical cancer compared to age-matched US Asians. The lack of screening may explain these disparities.

EPV156/#363

ADEQUACY OF INFORMATION RECEIVED FOR PATIENTS TREATED OVERSEAS AND ITS IMPACT ON CONTINUING TREATMENT AND FOLLOW-UP

¹MT Alsayed*, ¹S Brich, ²H Khaldi, ¹A Magzoub, ¹A Al -Ansari, ²S Chandramouli, ²JJO Herod. ¹Hamad Medical Corporation, Women Wellness and Research Center, DOHA, Qatar; ²Hamad Medical Corporation, Women Wellness and Research Center, Doha, Qatar

10.1136/ijgc-2021-IGCS.226

Objectives The gynaecological oncology service in Doha treats all women living in or visiting Qatar. Despite the quality and affordability of the service many women travel overseas for