



Incidence of lymph node metastasis in early-stage low-grade serous ovarian cancer: a systematic review

David Viveros-Carreño ^{1,2}, Juliana Rodriguez ^{1,3}, Rene Pareja ^{1,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ijgc-2022-003618>).

For numbered affiliations see end of article.

Correspondence to

Dr David Viveros-Carreño, Gynecologic Oncology, Instituto Nacional de Cancerología, Bogota, Colombia; dviveros@gmail.com

Received 13 April 2022

Accepted 27 June 2022

Published Online First

13 July 2022

HIGHLIGHTS

- ⇒ The rate of positive lymph nodes in patients with clinically presumed early-stage low-grade serous ovarian cancer was 12.6%.
- ⇒ The upstaging rate due to lymph node metastasis alone was 10.5%.
- ⇒ None of the studies included report specific complications of the lymph node dissection.

ABSTRACT

Objective The objective of this systematic review was to assess the incidence of lymph node metastasis in patients with clinically presumed early-stage low-grade serous ovarian cancer that underwent primary surgical treatment.

Methods This study was registered in PROSPERO (CRD42022308923). A systematic literature review was conducted following the Meta-analyses Of Observational Studies in Epidemiology (MOOSE) checklist. PubMed/MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, Ovid, and Scopus databases were searched since inception and up to March 2022. The inclusion criteria were: pathological confirmation of low-grade serous ovarian cancer (clinically presumed FIGO 2014 stages I-IIA at time of surgery) that underwent primary surgical treatment, including pelvic and/or para-aortic lymph node dissection.

Results The search identified 3763 articles; 59 were considered potentially eligible after removing duplicates, and eight studies finally met the selection criteria. In total, 35 of 277 (12.6%) patients had lymph node metastasis, and only four studies reported upstaging due to lymph node metastasis in 16 of 153 (10.5%) patients. None of the eight studies included reported the rate of complications or complications specifically for the subgroup of patients with early-stage low-grade tumors.

Conclusion In patients with early-stage low-grade serous ovarian cancer, lymph node assessment should be discussed when counseling for primary surgical staging.

INTRODUCTION

Ovarian cancer is the eighth most common cancer among women worldwide, with 6.6 cases per 1 000 women and 313 959 new cases reported in GLOBOCAN 2020. It has a disproportionately high mortality rate of 4.2 cases per 100.000 women,¹ and only about a third of the patients present with early-stage disease.² Primary surgical treatment is the preferred therapy for this population.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Low-grade epithelial ovarian tumors have a smaller risk of lymph node metastasis compared with high-grade epithelial tumors. However, the risk differs among histological subtypes (endometrioid, mucinous, or serous) and could be higher for low-grade serous tumors compared with the other low-grade epithelial tumors.

WHAT THIS STUDY ADDS

⇒ The risk of lymph node metastasis of low-grade serous tumors is higher than that previously reported for other low-grade epithelial tumors and the upstaging rate due to lymph node metastasis alone is higher than 10%.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The reported rate of lymph node involvement and the impact of its identification in adjuvant management decisions may justify the performance of lymph node assessment; however, the effect of this finding on oncological outcomes has not been demonstrated and given the rarity of this tumor is highly unlikely to be evaluated in prospective trials.

Low-grade serous ovarian cancer represents about 10% of epithelial ovarian cancer cases.³ Although there are differences with high-grade serous ovarian cancer in molecular biology, clinic characteristics, and prognosis, these patients are treated similarly. Low-grade serous ovarian cancer is considered relatively insensitive to systemic chemotherapy; however, it remains a standard treatment in advanced disease after maximum surgical effort.^{4,5}

The indication for lymph node dissection in ovarian cancer patients has been debated during the last several years for high-grade epithelial ovarian cancer. Recently a prospective trial⁶ demonstrated no benefit



© IGCS and ESGO 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Viveros-Carreño D, Rodriguez J, Pareja R. *Int J Gynecol Cancer* 2022;**32**:1283–1288.

Original research

of systematic lymphadenectomy in advanced disease with macroscopically complete resection and clinically apparent normal lymph nodes. On the other hand, for clinical early stages (International Federation of Gynecology and Obstetrics (FIGO) 2014 stage I-IIA) comprehensive surgical staging, including systematic bilateral pelvic and para-aortic lymphadenectomy up to the renal veins, still could have a crucial role due to the occult lymph node metastasis in apparent early-stage disease of nearly 15%⁷ and the subsequent change in the adjuvant treatment.^{5 8 9}

The incidence of lymph node metastasis may change according to histologic subtype and grading. A previous report showed that the incidence is only 2.9% in low-grade early-stage epithelial ovarian cancer.¹⁰ As the incidence for low-grade serous ovarian cancer could be higher than for low-grade endometrioid or mucinous histological types, the objective of this systematic review was to assess the incidence of lymph node metastasis in patients with clinically apparent early-stage low-grade serous ovarian cancer who underwent primary surgical treatment.

METHODS

The Meta-analyses Of Observational Studies in Epidemiology (MOOSE) checklist was used to conduct this systematic review and the protocol was registered in PROSPERO (CRD42022308923). MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, Ovid, and Scopus databases were searched since inception and up to March 2022. The search strategy is reported in the Online supplemental file 1. We also searched the reference lists of eligible studies and used forward citation tracking with Google Scholar to identify further eligible manuscripts. Similar methodology has been reported by the authors previously.¹¹

All types of studies published in the English language literature were included, except for case reports, abstracts, commentaries, and unpublished studies. The inclusion criteria were pathologically confirmed low-grade serous ovarian cancer (clinically presumed FIGO 2014 stages I-IIA at time of surgery) that underwent primary surgical treatment, including pelvic and/or para-aortic lymph node dissection. Any surgical route (laparotomy or minimally invasive surgery) was allowed. Only grade 1 tumors were considered if the tumor grade was specified, given significant variations in their classification into low- and high-grade for grade 2 tumors. For studies that did not specify tumor grade, all low-grade tumors were included.¹² Criteria for exclusion were patients under 18 years of age, neoadjuvant chemotherapy administration, and evidence of lymph node metastasis, either on pre-operative images, or at the time of surgery.

When two or more articles were published by the same authors or using the same data source, only the most recent article was included. The measured outcomes were the incidence of pelvic and/or para-aortic lymph node metastasis as defined by the authors, the rate of upstaging exclusively due to lymph node involvement, and the rate of complications associated with lymph node dissection as defined by the authors. Studies not reporting the number of resected nodes could be included in the review, and for studies reporting the number of lymph nodes resected at least 10 nodes per patient had to be resected to be included in the review.

Study Selection

Two authors (JR, DV-C) independently assessed all titles and abstracts of records retrieved from the search strategy for inclusion. The final selection of studies for inclusion was undertaken independently by three authors (RP, JR, DV-C) and any disagreement was resolved through discussion. We designed a form to extract data, which was pilot tested. Two authors (JR, DV-C) extracted the data independently using the form for eligible studies. Any disagreement about extracted data was resolved through discussion until a consensus was reached.

The information was presented as medians or means (according to its normal distribution) and percentages with absolute counts, if it was a quantitative or qualitative variable, respectively. The descriptive statistics were performed in SPSS 2.0. According to local regulations, no institutional review board agreement was required for this type of study. Ethical approval was not required as only data from previously published studies were retrieved and analyzed. No new data are presented.

RESULTS

The search identified 3763 articles; after duplicates were removed, 1874 articles were evaluated, and the title and abstract screening of these references identified 59 studies as potentially eligible for this review. The full-text screening excluded 51 studies, and eight studies finally met the selection criteria. Four studies were excluded because they did not provide information according to histological types, one was a literature review and did not have new patients reported, four did not include low-grade tumors, and 42 did not provide separate information for early-stage low-grade serous tumors (Figure 1 and Online supplemental file 2).

The total number of patients included in the eight studies was 2964, but only 277 patients had early-stage low-grade serous tumors and were considered for the analysis. Seven studies were retrospective series^{13–19} and one study was a population-based study.²⁰ The accrual time was from 1975 up to 2018. Three studies included only FIGO stage I tumors^{14 19 20} and five included FIGO stage I and II tumors.^{13 15 16} We selected only patients with tumors up to FIGO stage IIA tumors for the review. Tumor grade was reported and classified as low grade in five studies^{13–15 17 18} and reported as grade 1 serous tumors in three studies.^{16 19 20}

The details about the performed surgeries were not described in three studies,^{15 17 20} it was described as staging surgery or primary surgery in three studies,^{16 18 19} as a surgery that included hysterectomy, bilateral salpingo-oophorectomy, peritoneal random peritoneal biopsies, and omentectomy in one study,¹³ and as bilateral salpingo-oophorectomy (97.5%), hysterectomy (82.8%), omentectomy (98.8%), and appendectomy (68.1%) in one study.¹⁴ Two studies^{13 14} included patients with primary surgery and reclassification surgery; four included only primary surgery treatment,^{15 16 18 19} and it was not reported in two of the studies.^{17 20} Details about included studies are provided in Table 1. Given the nature of the studies and the intervention, with only single-arm studies, we did not evaluate the risk of bias for included studies as this would not provide reliable information.

Six studies reported pelvic and para-aortic lymph node dissection,^{13–16 18 19} one²⁰ did not report specifically if pelvic and para-aortic dissection was performed, and one¹⁷ reported some patients with




Figure 1 PRISMA flow diagram for study selection.

pelvic and others with pelvic and para-aortic dissection. Two studies included the number of resected lymph nodes; one reported for inclusion a minimum of 10 pelvic and 10 para-aortic nodes,¹⁴ and one at least 20 nodes in total.²⁰ Five studies reported the median number of resected nodes, although three studies^{14 16 19} reported data only for the complete cohort of included patients and not independent information on early stages. The median number of resected nodes were 45,^{14 19} 69,^{16 27, 20} and 41.¹⁸

In total, 35 (12.6%) of 277 patients had lymph node metastasis. All studies reported at least one case of lymph node involvement, and the location was reported only in three studies,^{13 17 19} one patient with pelvic and para-aortic involvement in one study, 14 of 14 patients with pelvic and 4 of 14 patients with para-aortic metastasis in the second study; and in the third study four only pelvic and eight both pelvic and para-aortic region. The rate of lymph node metastasis for each study ranged from 2.7%¹³ to 25%.¹⁸ Three studies^{14 17 19} reported the number of patients upstaged exclusively because of lymph node metastasis. Of 153 patients, 25 (16.3%) had lymph node metastasis and 16 (10.5%) were upstaged only due to lymph node involvement. Unfortunately, none of the eight included studies reported the rate of complications or complications specifically for the subgroup of interest of patients with early-stage low-grade serous tumors. Details about lymph node procedures are provided in [Table 2](#).

According to our protocol, we did two subgroup analyses. For the first analysis, excluding the population-based study, in seven studies^{13–19} 210 patients were reported and 29 patients had confirmed lymph node metastasis (13.8%). When only patients with stage I were analyzed, 183 patients from three studies^{14 19 20} were included and 24 (13.1%) had

lymph node metastasis. The results of the subgroup analyses showed no relevant effect on the general results.

DISCUSSION

Summary of Main Results

In our review, the overall rate of lymph node metastasis in patients with early-stage low-grade serous ovarian cancer was 12.6%. Patients were upstaged exclusively because of lymph node metastasis in 10.5% of cases. The rate of involvement in this group of patients may justify lymphadenectomy in primary surgical management.

Results in the Context of Published Literature

For all low-grade early-stage epithelial tumors, the incidence of lymph node metastasis has been reported previously in less than 3% of cases,^{10 20} unlike high-grade tumors where a lymph node metastasis rate of 10% or higher^{7 13 20} has been reported. In high-grade tumors, a plausible impact on oncologic prognosis²¹ is assumed by the lymph node dissection and the recommendation of systematic lymphadenectomy in clinical early stages as part of the primary surgical treatment in international guidelines is recommended.^{5 9}

The ovarian epithelial tumors are heterogeneous, in mucinous and endometrioid tumors molecular and clinical behavior can differ from that of serous tumors.^{22 23} These differences are reflected among others on the risk of lymph node metastasis; for expansile (low-grade) mucinous tumors the rate of involvement could be nearly zero^{24 25} and it is less than 2% for low-grade endometrioid tumors.^{14 20} However, this risk is up to 10% for low-grade serous tumors.

Table 1 Major characteristics of studies included in the systematic review

Author/year	Accrual time	Study type	N total	N early (I-IIA) low-grade serous	Age (years)	FIGO stage clinically presumed	Histologic subtype grade	Procedure/fertility-preserving	Procedure/primary vs re-staging
Bogani 2017 ¹³	1975–2016	Retrospective	290	37	49.8 (18.1–76) median	I-II (85.9%, 14.1%)	Low-grade	Hysterectomy, bilateral salpingo-oophorectomy, peritoneal random peritoneal biopsies; omentectomy	Both
Minig 2017 ¹⁴	2000–2016	Retrospective	163	56	50 median (18–81) whole cohort	I	Low-grade	Bilateral salpingo-oophorectomy (97.5%), hysterectomy (82.8%), omentectomy (98.8%), and appendectomy (68.1%)	Both
Heitz 2018 ¹⁶	2000–2016	Retrospective	762	5	57 median (18–82) whole cohort	pT1a-pT2aM0 (I-II)	Grade 1	Staging surgery	Primary
Nasioudis 2018 ²⁰	1988–2014	Population-based SEER	1242	67	51 median	IA, IB, IC	Grade 1	NR	NR
Wafa 2019 ¹⁸	2000–2017	Retrospective	37	4	48 (26–76) whole cohort	pT1a-pT2aM0 (I-II)	Low-grade	Primary surgery	Primary
Simon 2020 ¹⁵	2000–2017	Retrospective	126	11	53 median (19–82)	I-II	Low-grade	NR	Primary
Chen 2021 ¹⁷	2005–2015	Retrospective	196	37	NR as median or average	I-IIA	Low-grade	NR	NR
Durmuş 2022 ¹⁹	1999–2018	Retrospective	148	60	49.1 mean (21–84)	I	Grade 1/low-grade	Both (complete surgery consisted of total hysterectomy, bilateral salpingo-oophorectomy, omentectomy, and peritoneal biopsies)	Primary
Total			2964	277					

FIGO, International Federation of Gynecology and Obstetrics; NR, not reported; SEER, Surveillance, Epidemiology, and End Results Program.

Table 2 Lymph node procedures and pathology report.

Author/year	Lymph node procedure	Lymph node count: total	N N(+)	Location N(+)	N N(+)/total as percentage (%)	N upstaged due to N(+)
Bogani 2017 ¹³	Pelvic and para-aortic dissection	NR	1	Pelvic and para-aortic	2.72	NR
Minig 2017 ¹⁴	Pelvic and para-aortic dissection (minimum number of 10 pelvic nodes and 10 para-aortic nodes)	45 median (whole cohort)	6	NR	10.71	3
Heitz 2018 ¹⁶	Pelvic and para-aortic dissection	69 median (whole cohort)	1	NR	20	NR
Nasioudis 2018 ²⁰	NR (at least 20 nodes)	27 median	6	NR	8.95	NR
Wafa 2019 ¹⁸	Pelvic and para-aortic dissection	41 median	1	NS	25	NR
Simon 2020 ¹⁵	Pelvic and para-aortic dissection	NR	1	NR	9.09	NR
Chen 2021 ¹⁷	Pelvic (more than 10 nodes) in 25 patients and systematic lymph node dissection (more than 10 pelvic and 5 para-aortic nodes) in 12 patients	20 (pelvic) in 25 patients and 25 plus 8 (pelvic and para-aortic in 12 patients)	7	14/14 pelvic, 4/14 para-aortic in the whole cohort.	18.91	4
Durmuş 2022 ¹⁹	Bilateral pelvic lymphadenectomy and para-aortic lymphadenectomy	45 median (whole cohort)	12	4 only pelvic, 8 both pelvic and para-aortic	20	9
Total			35		12.64	

NR, not reported; NS, not specified.

The relevance of lymph node metastasis identification is associated with the requirement for adjuvant treatment and the oncological prognosis of patients. Guidelines^{5,9} recommend abstaining from adjuvant chemotherapy in patients with early stages (FIGO 2014 IA-IB) of low-grade serous ovarian cancer but recommend using combined systemic chemotherapy in cases of lymph node metastasis or if other advanced-stage features are present (FIGO 2014 IC-IV stage). Lymph node metastasis in advanced low-grade epithelial ovarian tumors is frequent with involvement in more than 70% of patients.^{15,18} However, it has been suggested that the lymph node dissection of microscopic disease has no impact on overall survival or progression-free survival in this population.¹⁵ The reporting of only early stages is limited in the literature; most studies report all low-grade early-stage tumors together or with all stages, and the risk of lymph node metastasis could have been underestimated previously.

Among the most relevant outcomes remains the rate of upstaging exclusively due to lymph node metastasis as a justification for the procedure; in our review it was 10.5%, whereas a previous systematic review that assessed the upstaging rate of the procedures that are performed in the primary surgical treatment of patients with epithelial ovarian cancer found that the upstaging rate for hysterectomy, omentectomy, peritoneal biopsies, appendectomy, and cytology did not exceed 10%.²⁶

Unfortunately, in our review the studies included did not specifically report adverse events associated with lymphadenectomy; however, it is well known even from prospective randomized trials that lymph node dissection raises the risk of complications related to the surgery as

evidenced by higher median blood loss, requirement for transfusions, intensive care support, higher infections, and reintervention rates.^{6,27}

Strengths and Weaknesses

The main strength of our review is the rigorous process for data collection, and we have a pre-registered protocol with specified selection criteria and a rigorous process for data analysis. Our review is limited for several reasons. The clinical heterogeneity of the studies must be highlighted as we included case series studies and a population database study. The retrospective nature of all studies included in the review, with potential selection and publication bias, and the relatively small number of patients included in the results given the limited report of this specific population, are limiting factors, particularly since we did not have information from most studies for adverse events, for upstaging exclusively for lymph node metastasis, and lymph node metastasis location. There was no central pathology review, surgical quality report beyond lymph node count for all studies, or complete information about pre-operative imaging studies. As we limited the search to published data in the English language, we could have missed some relevant references published in other languages, and as the studies were conducted over a long period, there could be performance bias.

Implications for Practice and Future Research

This systematic review confirms that the risk of lymph node metastasis for early-stage low-grade serous ovarian cancer is different than for other low-grade epithelial tumors. As the identification of lymph node

Original research

involvement will change the recommendation of adjuvant chemotherapy, lymphadenectomy should be recommended. We encourage collection of prospective information in this population with subsequent data analysis according to the tumor type for future publications.

CONCLUSION

Given the rate of lymph node metastasis in clinically presumed early-stage low-grade serous ovarian cancer and the impact of its identification on adjuvant treatment decisions, lymph node assessment should be considered as part of primary surgical staging.

Author affiliations

¹Department of Gynecologic Oncology, Instituto Nacional de Cancerología, Bogotá, Colombia

²Gynecologic Oncology, Clínica Universitaria Colombia and Clínica Los Nogales, Bogotá, Colombia

³Department of Gynecology and Obstetrics, section of Gynecologic Oncology, Fundación Santa Fe de Bogotá, Bogotá, Colombia

⁴Gynecologic Oncology, Clínica ASTORGA, Medellín, Colombia

Contributors DV-C: conceptualization, investigation, methodology, writing - original draft, writing - review, editing and responsible for the overall content as guarantor. JR: data curation, formal analysis, investigation, methodology, writing - review, and editing. RP: conceptualization, methodology, formal analysis, writing - review and editing, supervision.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

David Viveros-Carreño <http://orcid.org/0000-0001-9395-0627>

Juliana Rodríguez <http://orcid.org/0000-0002-5472-4093>

Rene Pareja <http://orcid.org/0000-0003-0093-0438>

REFERENCES

- Sung H, Ferlay J, Siegel RL, *et al*. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71:209–49.
- Young RC, Walton LA, Ellenberg SS, *et al*. Adjuvant therapy in stage I and stage II epithelial ovarian cancer. Results of two prospective randomized trials. *N Engl J Med* 1990;322:1021–7.
- Seidman JD, Horkayne-Szakaly I, Haiba M, *et al*. The histologic type and stage distribution of ovarian carcinomas of surface epithelial origin. *Int J Gynecol Pathol* 2004;23:41–4.
- Kaldawy A, Segev Y, Lavie O, *et al*. Low-grade serous ovarian cancer: a review. *Gynecol Oncol* 2016;143:433–8.
- Armstrong DK, Alvarez RD, Bakkum-Gamez JN, *et al*. Ovarian cancer, version 2.2020, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2021;19:191–226.
- Harter P, Sehouli J, Lorusso D, *et al*. A randomized trial of lymphadenectomy in patients with advanced ovarian neoplasms. *N Engl J Med* 2019;380:822–32.
- Kleppe M, Wang T, Van Gorp T, *et al*. Lymph node metastasis in stages I and II ovarian cancer: a review. *Gynecol Oncol* 2011;123:610–4.
- Moore KN, Pothuri B. Poly(ADP-ribose) polymerase inhibitor inhibition in ovarian cancer: a comprehensive review. *Cancer J* 2021;27:432–40.
- Colombo N, Sessa C, du Bois A, *et al*. ESMO-ESGO consensus conference recommendations on ovarian cancer: pathology and molecular biology, early and advanced stages, borderline tumours and recurrent disease†. *Ann Oncol* 2019;30:672–705.
- Lago V, Minig L, Fotopoulou C. Incidence of lymph node metastases in apparent early-stage low-grade epithelial ovarian cancer: a comprehensive review. *Int J Gynecol Cancer* 2016;26:1407–14.
- Viveros-Carreño D, Rodríguez J, Pareja R. Incidence of metastasis in circumflex iliac nodes in patients with cervical cancer: a systematic review. *Int J Gynecol Cancer* 2021;31:1530–4.
- Malpica A, Deavers MT, Lu K, *et al*. Grading ovarian serous carcinoma using a two-tier system. *Am J Surg Pathol* 2004;28:496–504.
- Bogani G, Tagliabue E, Ditto A, *et al*. Assessing the risk of pelvic and para-aortic nodal involvement in apparent early-stage ovarian cancer: a predictors- and nomogram-based analyses. *Gynecol Oncol* 2017;147:61–5.
- Minig L, Heitz F, Cibula D, *et al*. Patterns of lymph node metastases in apparent stage I low-grade epithelial ovarian cancer: a multicenter study. *Ann Surg Oncol* 2017;24:2720–6.
- Simon V, Ngo C, Pujade-Lauraine E, *et al*. Should we abandon systematic pelvic and paraaortic lymphadenectomy in low-grade serous ovarian cancer? *Ann Surg Oncol* 2020;27:3882–90.
- Heitz F, Harter P, Ataseven B, *et al*. Stage- and histologic subtype-dependent frequency of lymph node metastases in patients with epithelial ovarian cancer undergoing systematic pelvic and paraaortic lymphadenectomy. *Ann Surg Oncol* 2018;25:2053–9.
- Chen J, Yin J, Li Y, *et al*. Systematic lymph node dissection may be abolished in patients with apparent early-stage low-grade mucinous and endometrioid epithelial ovarian cancer. *Front Oncol* 2021;11:705720.
- Wafa M, Braicu EI, Muallem MZ, *et al*. Incidence and pattern of spread of lymph node metastasis in patients with low-grade serous ovarian cancer. *Anticancer Res* 2019;39:5617–21.
- Durmuş Y, Meydanlı MM, Akıllı H, *et al*. Factors associated with the involvement of lymph nodes in low-grade serous ovarian cancer. *J Surg Oncol* 2022;125:264–72.
- Nasioudis D, Mastroiannis SA, Ko EM, *et al*. Does tumor grade influence the rate of lymph node metastasis in apparent early stage ovarian cancer? *Arch Gynecol Obstet* 2018;298:179–82.
- Trimbos JB, Vergote I, Bolis G, *et al*. Impact of adjuvant chemotherapy and surgical staging in early-stage ovarian carcinoma: European Organisation for Research and Treatment of Cancer-Adjuvant ChemoTherapy in Ovarian Neoplasm Trial. *J Natl Cancer Inst* 2003;95:113–25.
- Dion L, Carton I, Jaillard S, *et al*. The landscape and therapeutic implications of molecular profiles in epithelial ovarian cancer. *J Clin Med* 2020;9:2239.
- Romero I, Leskelä S, Mies BP, *et al*. Morphological and molecular heterogeneity of epithelial ovarian cancer: therapeutic implications. *EJC Suppl* 2020;15:1–15.
- Gouy S, Saidani M, Maulard A, *et al*. Staging surgery in early-stage ovarian mucinous tumors according to expansile and infiltrative types. *Gynecol Oncol Rep* 2017;22:21–5.
- Schmeler KM, Tao X, Frumovitz M, *et al*. Prevalence of lymph node metastasis in primary mucinous carcinoma of the ovary. *Obstet Gynecol* 2010;116:269–73.
- van de Vorst REWM, Hoogendam JP, van der Aa MA, *et al*. The attributive value of comprehensive surgical staging in clinically early-stage epithelial ovarian carcinoma: a systematic review and meta-analysis. *Gynecol Oncol* 2021;161:876–83.10.1016/j.ygyno.2021.04.007
- Panici PB, Maggioni A, Hacker N, *et al*. Systematic aortic and pelvic lymphadenectomy versus resection of bulky nodes only in optimally debulked advanced ovarian cancer: a randomized clinical trial. *J Natl Cancer Inst* 2005;97:560–6.

Supplement 1. Search strategy

Database	Strategy search	Field search	Filters
PubMed	((("ovarian cancer*" OR "ovary cancer*" OR "ovarian neoplasm*" OR "ovary neoplasm*" OR "ovarian carcinoma*" OR "ovarian epithelial cancer*" OR "epithelial ovarian cancer*" OR "carcinoma, ovarian epithelial" OR "low-grade ovarian cancer" OR "low grade ovarian cancer" OR "low-grade serous*" OR "low grade serous*" OR "early-stage low-grade*") AND ("primary surgery" OR laparoscopy OR laparotomy OR "open approach" OR "robotic approach" OR "laparoscopic approach" OR "surgical procedures" OR "staging procedures" OR "surgical staging" OR surgical* OR "neoplasm staging" OR "cytoreduction surgical procedures*")) AND ("lymph nodes*" OR "lymph node dissection*" OR "lymph node excision*" OR "para-aortic lymph node*" OR "paraaortic lymphadenectomy*" OR "pelvic lymph node*" OR "lymphadenectomy*" OR "systematic lymphadenectomy*" OR "systematic pelvic*" OR "systematic paraaortic*")) AND (metastas* OR "neoplasm metastas*" OR "lymphatic spread" OR "nodal involvement*" OR "lymph node metastas*" OR "lymphatic metastasis" OR "intraoperative complication") Filters: English	All Fields	Languages (English)
Embase	('ovarian cancer*' OR 'ovary cancer*' OR 'ovarian neoplasm*' OR 'ovary neoplasm*' OR 'ovarian carcinoma*' OR 'ovarian epithelial cancer*' OR 'epithelial ovarian cancer*' OR 'carcinoma, ovarian epithelial' OR 'low-grade ovarian cancer' OR 'low grade ovarian cancer' OR 'low-grade serous*' OR 'low grade serous*' OR 'early-stage low-grade*') AND ('primary surgery' OR laparoscopy OR laparotomy OR 'open approach' OR 'robotic approach' OR 'laparoscopic approach' OR 'surgical procedures' OR 'staging procedures' OR 'surgical staging' OR surgical* OR 'neoplasm staging' OR 'cytoreduction surgical procedures*') AND ('lymph nodes*' OR 'lymph node dissection*' OR 'lymph node excision*' OR 'para-aortic lymph node*' OR 'paraaortic lymphadenectomy*' OR 'pelvic lymph node*' OR 'lymphadenectomy*' OR 'systematic lymphadenectomy*' OR 'systematic pelvic*' OR 'systematic paraaortic*') AND (metastas* OR 'neoplasm metastas*' OR 'lymphatic spread' OR 'nodal involvement*' OR 'lymph node metastas*' OR 'lymphatic metastasis' OR 'intraoperative complication') AND [english]/lim	All Fields	Languages (English)

Database	Strategy search	Field search	Filters
Cochrane	("ovarian cancer*" OR "ovary cancer*" OR "ovarian neoplasm*" OR "ovary neoplasm*" OR "ovarian carcinoma*" OR "ovarian epithelial cancer*" OR "epithelial ovarian cancer*" OR "carcinoma, ovarian epithelial" OR "low-grade ovarian cancer" OR "low grade ovarian cancer" OR "low-grade serous*" OR "low grade serous*" OR "early-stage low-grade*") AND ("primary surgery" OR laparoscopy OR laparotomy OR "open approach" OR "robotic approach" OR "laparoscopic approach" OR "surgical procedures" OR "staging procedures" OR "surgical staging" OR surgical* OR "neoplasm staging" OR "cytoreduction surgical procedures*") in All Text AND "lymph nodes*" OR "lymph node dissection*" OR "lymph node excision*" OR "para-aortic lymph node*" OR "paraaortic lymphadenectom*" OR "pelvic lymph node*" OR "lymphadenectom*" OR "systematic lymphadenectom*" OR "systematic pelvic*" OR "systematic paraaortic*" in All Text AND metastas* OR "neoplasm metastas*" OR "lymphatic spread" OR "nodal involvement*" OR "lymph node metastas*" OR "lymphatic metastasis" OR "intraoperative complication" in All Text - (Word variations have been searched)	All text	None
Scopus	(TITLE-ABS-KEY (("ovarian cancer*" OR "ovary cancer*" OR "ovarian neoplasm*" OR "ovary neoplasm*" OR "ovarian carcinoma*" OR "ovarian epithelial cancer*" OR "epithelial ovarian cancer*" OR "carcinoma, ovarian epithelial" OR "low-grade ovarian cancer" OR "low grade ovarian cancer" OR "low-grade serous*" OR "low grade serous*" OR "early-stage low-grade*") AND ("primary surgery" OR laparoscopy OR laparotomy OR "open approach" OR "robotic approach" OR "laparoscopic approach" OR "surgical procedures" OR "staging procedures" OR "surgical staging" OR surgical* OR "neoplasm staging" OR "cytoreduction surgical procedures*")) AND TITLE-ABS-KEY ("lymph nodes*" OR "lymph node dissection*" OR "lymph node excision*" OR "para-aortic lymph node*" OR "paraaortic lymphadenectom*" OR "pelvic lymph node*" OR "lymphadenectom*" OR "systematic lymphadenectom*" OR "systematic pelvic*" OR "systematic paraaortic*") AND TITLE-ABS-KEY (metastas* OR "neoplasm metastas*" OR "lymphatic spread" OR "nodal involvement*" OR "lymph node metastas*" OR "lymphatic metastasis" OR "intraoperative complication")) AND (LIMIT-TO (LANGUAGE , "English"))	Article title, Abstract, Keywords (All fields 12.788)	Language (English)

Database	Strategy search	Field search	Filters
Ovid	<p>1 (("ovarian cancer*" or "ovary cancer*" or "ovarian neoplasm*" or "ovary neoplasm*" or "ovarian carcinoma*" or "ovarian epithelial cancer*" or "epithelial ovarian cancer*" or "carcinoma, ovarian epithelial" or "low-grade ovarian cancer" or "low grade ovarian cancer" or "low-grade serous*" or "low grade serous*" or "early-stage low-grade*") and ("primary surgery" or laparoscopy or laparotomy or "open approach" or "robotic approach" or "laparoscopic approach" or "surgical procedures" or "staging procedures" or "surgical staging" or surgical* or "neoplasm staging" or "cytoreduction surgical procedures*")).ab. and ("lymph nodes*" or "lymph node dissection*" or "lymph node excision*" or "para-aortic lymph node*" or "paraaortic lymphadenectom*" or "pelvic lymph node*" or "lymphadenectom*" or "systematic lymphadenectom*" or "systematic pelvic*" or "systematic paraaortic*").af. and (metastas* or "neoplasm metastas*" or "lymphatic spread" or "nodal involvement*" or "lymph node metastas*" or "lymphatic metastasis" or "intraoperative complication").af.</p> <p>2 limit 3 to english [Limit not valid in Books@Ovid,CLCMR,DARE,CLEED; records were retained]</p>	All Fields	<p>Language Publication types Ovid Full Text Available Remove duplicates</p>

Supplement 2. Excluded articles

Did not provide information according to histological types

1. Burghardt E, Girardi F, Lahousen M, Tamussino K, Stettner H. Patterns of pelvic and paraaortic lymph node involvement in ovarian cancer. *Gynecol Oncol*. Feb 1991;40(2):103-6. doi:10.1016/0090-8258(91)90099-q
2. Kanazawa K, Suzuki T, Tokashiki M. The validity and significance of substage IIIC by node involvement in epithelial ovarian cancer: impact of nodal metastasis on patient survival. *Gynecol Oncol*. May 1999;73(2):237-41. doi:10.1006/gyno.1999.5349
3. Pickel H, Lahousen M, Stettner H, Girardi F. The spread of ovarian cancer. *Baillieres Clin Obstet Gynaecol*. Mar 1989;3(1):3-12. doi:10.1016/s0950-3552(89)80036-7
4. Suzuki M, Ohwada M, Yamada T, Kohno T, Sekiguchi I, Sato I. Lymph node metastasis in stage I epithelial ovarian cancer. *Gynecol Oncol*. Nov 2000;79(2):305-8. doi:10.1006/gyno.2000.5951

Literature review

1. Lago V, Minig L, Fotopoulou C. Incidence of Lymph Node Metastases in Apparent Early-Stage Low-Grade Epithelial Ovarian Cancer: A Comprehensive Review. *Int J Gynecol Cancer*. Oct 2016;26(8):1407-14. doi:10.1097/igc.0000000000000787

Did not include low-grade tumors

1. Timmers PJ, Zwinderman K, Coens C, Vergote I, Trimbos JB. Lymph node sampling and taking of blind biopsies are important elements of the surgical staging of early ovarian cancer. *Int J Gynecol Cancer*. Oct 2010;20(7):1142-7. doi:10.1111/igc.0b013e3181ef8e03
2. Bizzarri N, du Bois A, Fruscio R, et al. Is there any therapeutic role of pelvic and para-aortic lymphadenectomy in apparent early stage epithelial ovarian cancer? *Gynecologic Oncology*. 2021;160(1):56-63.
3. Brockbank EC, Harry V, Kolomainen D, et al. Laparoscopic staging for apparent early stage ovarian or fallopian tube cancer. First case series from a UK cancer centre and systematic literature review. *Eur J Surg Oncol*. Aug 2013;39(8):912-7. doi:10.1016/j.ejso.2013.05.007
4. Knipscheer RJ. Para-aortal lymph nodes dissection in 20 cases of primary epithelial ovary carcinoma stage I (Figo): influence on staging. *Eur J Obstet Gynecol Reprod Biol*. Aug 1982;13(5):303-7. doi:10.1016/0028-2243(82)90053-3

Did not provide separate information for early-stage low-grade serous tumors

1. Bacalbasa N, Balescu I, Vilcu M, et al. The Risk of Para-Aortic Lymph Node Metastases in Apparent Early Stage Ovarian Cancer. *Medicina (Kaunas)*. Mar 3 2020;56(3)doi:10.3390/medicina56030108
2. Baiocchi, Raspagliesi, Grosso, Fontanelli, Cobellis, Re D. Early ovarian cancer: Is there a role for systematic pelvic and para-aortic lymphadenectomy? *International Journal of Gynecological Cancer*. 1998;8(2):103-108. doi:<https://doi.org/10.1046/j.1525-1438.1998.09758.x>
3. Benedetti-Panici P, Greggi S, Maneschi F, et al. Anatomical and pathological study of retroperitoneal nodes in epithelial ovarian cancer. *Gynecol Oncol*. Nov 1993;51(2):150-4. doi:10.1006/gyno.1993.1263
4. Cass I, Li AJ, Runowicz CD, et al. Pattern of lymph node metastases in clinically unilateral stage I invasive epithelial ovarian carcinomas. *Gynecol Oncol*. Jan 2001;80(1):56-61. doi:10.1006/gyno.2000.6027
5. Chan JK, Munro EG, Cheung MK, et al. Association of lymphadenectomy and survival in stage I ovarian cancer patients. *Obstet Gynecol*. Jan 2007;109(1):12-9. doi:10.1097/01.AOG.0000249610.95885.ef
6. Chang SJ, Bristow RE, Ryu HS. Analysis of para-aortic lymphadenectomy up to the level of the renal vessels in apparent early-stage ovarian cancer. *J Gynecol Oncol*. Jan 2013;24(1):29-36. doi:10.3802/jgo.2013.24.1.29
7. Deng T, Huang Q, Wan T, et al. The impact of lymph node dissection on survival in patients with clinical early-stage ovarian cancer. *J Gynecol Oncol*. May 2021;32(3):e40. doi:10.3802/jgo.2021.32.e40
8. Desteli GA, Gultekin M, Usubutun A, Yuce K, Ayhan A. Lymph node metastasis in grossly apparent clinical stage Ia epithelial ovarian cancer: Hacettepe experience and review of literature. *World J Surg Oncol*. Nov 30 2010;8:106. doi:10.1186/1477-7819-8-106
9. Garcia-Soto AE, Boren T, Wingo SN, Heffernen T, Miller DS. Is comprehensive surgical staging needed for thorough evaluation of early-stage ovarian carcinoma? *Am J Obstet Gynecol*. Mar 2012;206(3):242.e1-5. doi:10.1016/j.ajog.2011.08.022
10. Grabowski JP, Glajzer J, Richter R, et al. Lymphovascular space invasion and Ki67 as predictors of lymph node metastasis in primary low grade serous ovarian cancer. *Int J Gynecol Cancer*. Jan 2021;31(1):98-103. doi:10.1136/ijgc-2020-001950
11. Harter P, Gnauert K, Hils R, et al. Pattern and clinical predictors of lymph node metastases in epithelial ovarian cancer. *Int J Gynecol Cancer*. Nov-Dec 2007;17(6):1238-44. doi:10.1111/j.1525-1438.2007.00931.x
12. Lee CL, Kusunoki S, Huang CY, Wu KY, Lee PS, Huang KG. Surgical and survival outcomes of laparoscopic staging surgery for patients with stage I ovarian cancer. *Taiwan J Obstet Gynecol*. Feb 2018;57(1):7-12. doi:10.1016/j.tjog.2017.12.002

13. Lepinay K, Szubert S, Lewandowska A, et al. The association between lymph node metastases and long-term survival in patients with epithelial ovarian cancer. *Contemp Oncol (Pozn)*. 2020;24(3):163-171. doi:10.5114/wo.2020.99029
14. Li X, Xing H, Li L, et al. Clinical significance of para-aortic lymph node dissection and prognosis in ovarian cancer. *Front Med*. Mar 2014;8(1):96-100. doi:10.1007/s11684-014-0316-4
15. Maggioni A, Benedetti Panici P, Dell'Anna T, et al. Randomised study of systematic lymphadenectomy in patients with epithelial ovarian cancer macroscopically confined to the pelvis. *Br J Cancer*. Sep 18 2006;95(6):699-704. doi:10.1038/sj.bjc.6603323
16. Mattes MD, Lee JC, Metzger DJ, Ashamalla H, Katsoulakis E. The incidence of pelvic and para-aortic lymph node metastasis in uterine papillary serous and clear cell carcinoma according to the SEER registry. *J Gynecol Oncol*. Jan 2015;26(1):19-24. doi:10.3802/jgo.2015.26.1.19
17. Morice P, Joulie F, Camatte S, et al. Lymph node involvement in epithelial ovarian cancer: analysis of 276 pelvic and paraaortic lymphadenectomies and surgical implications. *J Am Coll Surg*. Aug 2003;197(2):198-205. doi:10.1016/s1072-7515(03)00234-5
18. Nomura H, Tsuda H, Susumu N, et al. Lymph node metastasis in grossly apparent stages I and II epithelial ovarian cancer. *Int J Gynecol Cancer*. Apr 2010;20(3):341-5. doi:10.1111/IGC.0b013e3181cf6271
19. Onda T, Yoshikawa H, Yokota H, Yasugi T, Taketani Y. Assessment of metastases to aortic and pelvic lymph nodes in epithelial ovarian carcinoma. A proposal for essential sites for lymph node biopsy. *Cancer*. Aug 15 1996;78(4):803-8. doi:10.1002/(sici)1097-0142(19960815)78:4<803::aid-cnrcr17>3.0.co;2-z
20. Oshita T, Itamochi H, Nishimura R, et al. Clinical impact of systematic pelvic and para-aortic lymphadenectomy for pT1 and pT2 ovarian cancer: a retrospective survey by the Sankai Gynecology Study Group. *Int J Clin Oncol*. Dec 2013;18(6):1107-13. doi:10.1007/s10147-012-0483-8
21. Petru E, Lahousen M, Tamussino K, et al. Lymphadenectomy in stage I ovarian cancer. *Am J Obstet Gynecol*. Feb 1994;170(2):656-62. doi:10.1016/s0002-9378(94)70244-6
22. Powless CA, Aletti GD, Bakkum-Gamez JN, Cliby WA. Risk factors for lymph node metastasis in apparent early-stage epithelial ovarian cancer: implications for surgical staging. *Gynecol Oncol*. Sep 2011;122(3):536-40. doi:10.1016/j.ygyno.2011.05.001
23. Svolgaard O, Lidegaard O, Nielsen ML, et al. Lymphadenectomy in surgical stage I epithelial ovarian cancer. *Acta Obstet Gynecol Scand*. Mar 2014;93(3):256-60. doi:10.1111/aogs.12322

24. Ulker V, Kuru O, Numanoglu C, Akbayır O, Polat I, Uhri M. Lymph node metastasis in patients with epithelial ovarian cancer macroscopically confined to the ovary: review of a single-institution experience. *Arch Gynecol Obstet*. May 2014;289(5):1087-92. doi:10.1007/s00404-013-3078-3
25. Zhou J, Sun JY, Wu SG, et al. Risk factors for lymph node metastasis in ovarian cancer: Implications for systematic lymphadenectomy. *Int J Surg*. May 2016;29:123-7. doi:10.1016/j.ijsu.2016.03.039
26. Erdem B, Yüksel IT, Peker N, et al. Evaluation of Factors Affecting Lymph Node Metastasis in Clinical Stage I-II Epithelial Ovarian Cancer. *Oncol Res Treat*. 2018;41(7-8):444-448. doi:10.1159/000488082
27. Haller H, Mamula O, Krasevic M, et al. Frequency and distribution of lymph node metastases in epithelial ovarian cancer. *Int J Gynecol Cancer*. Feb 2011;21(2):245-50.
28. Fournier M, Stoeckle E, Guyon F, et al. Lymph node involvement in epithelial ovarian cancer: sites and risk factors in a series of 355 patients. *Int J Gynecol Cancer*. Nov 2009;19(8):1307-13. doi:10.1111/IGC.0b013e3181b8a07c
29. Signorelli M, Guerra L, Pirovano C, et al. Detection of nodal metastases by 18F-FDG PET/CT in apparent early stage ovarian cancer: a prospective study. *Gynecol Oncol*. Nov 2013;131(2):395-9. doi:10.1016/j.ygyno.2013.08.022
30. Bachmann C, Krämer B, Brucker SY, et al. Relevance of pelvic and para-aortic node metastases in early-stage ovarian cancer. *Anticancer Res*. Nov 2014;34(11):6735-8.
31. Ditto A, Martinelli F, Reato C, et al. Systematic para-aortic and pelvic lymphadenectomy in early stage epithelial ovarian cancer: a prospective study. *Ann Surg Oncol*. Nov 2012;19(12):3849-55. doi:10.1245/s10434-012-2439-7
32. Faught W, Le T, Fung Kee Fung M, Krepart G, Lotocki R, Heywood M. Early ovarian cancer: what is the staging impact of retroperitoneal node sampling? *J Obstet Gynaecol Can*. Jan 2003;25(1):18-21. doi:10.1016/s1701-2163(16)31078-7
33. Baiomy TA, OHK, WMA, et al. Ovarian Cancer Surgical Staging, Laparoscopy Versus Laparotomy: A Comparative Study. *Journal of Gynecologic Surgery*. 2020;36(4):179-183. doi:10.1089/gyn.2019.0109
34. Hengeveld EM, Zusterzeel PLM, Lajer H, Høgdall CK, Rosendahl M. The value of surgical staging in patients with apparent early stage epithelial ovarian carcinoma. *Gynecol Oncol*. Aug 2019;154(2):308-313. doi:10.1016/j.ygyno.2019.06.006
35. Matsuo K, Wong KK, Fotopoulou C, et al. Impact of lympho-vascular space invasion on tumor characteristics and survival outcome of women with low-grade serous ovarian carcinoma. *Journal of surgical oncology*. 2018;117(2):236-244.

36. Khouly NI, Elkelani OA, Elhalaby AF. Lymph node involvement in clinically apparent early ovarian cancer: a prospective study. *Eur J Gynaecol Oncol.* 2017;38(2):232-235.
37. Brown JV, Mendivil AA, Abaid LN, et al. The safety and feasibility of robotic-assisted lymph node staging in early-stage ovarian cancer. *Int J Gynecol Cancer.* Oct 2014;24(8):1493-8. doi:10.1097/IGC.0000000000000224
38. Ercelep O, Ozcelik M, Gumus M. Association of lymphadenectomy and survival in epithelial ovarian cancer. *Curr Probl Cancer.* 04 2019;43(2):151-159. doi:10.1016/j.currprobcancer.2018.08.002
39. Wang J, Li J, Chen R, Lu X. Survival effect of different lymph node staging methods on ovarian cancer: An analysis of 10 878 patients. *Cancer Med.* Sep 2018;7(9):4315-4329. doi:10.1002/cam4.1680
40. Di Re F, Fontanelli R, Raspagliesi F, Di Re E. Pelvic and para-aortic lymphadenectomy in cancer of the ovary. *Baillieres Clin Obstet Gynaecol.* Mar 1989;3(1):131-42. doi:10.1016/s0950-3552(89)80048-3
41. Chen SS, Lee L. Incidence of para-aortic and pelvic lymph node metastases in epithelial carcinoma of the ovary. *Gynecol Oncol.* Aug 1983;16(1):95-100. doi:10.1016/0090-8258(83)90013-6
42. Díaz-Feijoo B, Bebia V, Hernández A, et al. Surgical complications comparing extraperitoneal vs transperitoneal laparoscopic aortic staging in early stage ovarian and endometrial cancer. *Gynecol Oncol.* Jan 2021;160(1):83-90. doi:10.1016/j.ygyno.2020.10.038