



Perioperative ovarian cancer guidelines: prevention and management of upper abdominal complications

Cagatay Taskiran ,¹ Pawel Knapp,² Christina Fotopoulou ³

Christina Fotopoulou

³Gynaecologic Oncology, Imperial College London Faculty of Medicine, London, UK



Biography: Christina Fotopoulou is the Professor of Gynaecological Cancer Surgery in the Department of Surgery and Cancer, Faculty of Medicine of Imperial College London, UK. She is the Deputy director of the Ovarian Cancer Action Research Centre at Imperial College. She holds an honorary chair in the Gynaecology Department at the Charité' University of Berlin, where she was trained and then later took the role of the Vice Director of the Gynecological Department. Her surgical and scientific expertise focuses on the management of patients with advanced and relapsed ovarian cancer, profiling of tumor heterogeneity and integration of tumor biology factors with surgical effort under the umbrella of individualization of surgical care. She has served as the Chair of the guidelines committees of the British Gynaecological Cancer Society (BGCS) and of ESGO (European Society of Gynaecologic Oncology). She has been an elected member of the ESGO Council and is also a member of the German AGO- Ovarian Cancer Group. She is on the editorial board and reviewer of numerous international gynaecological and oncological journals and is a member of various international oncological committees, including BGCS, ASCO, ESGO, IGCS, ESMO, ENGOT, AGO, SGO and NOGGO.

For numbered affiliations see end of article.

Correspondence to

Professor Christina Fotopoulou, Gynaecologic Oncology, Imperial College London Faculty of Medicine, London W12 0HS, UK; chfotopoulou@gmail.com

For 'Presented at statement' see end of article.

Accepted 8 August 2022
Published Online First
3 October 2022



© IGCS and ESGO 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Taskiran C, Knapp P, Fotopoulou C. *Int J Gynecol Cancer* 2022;32:1486–1487.

More than half of patients with advance ovarian cancer will require upper abdominal and multivisceral debulking techniques so that total macroscopic tumor clearance can be achieved, including diaphragmatic surgery, liver resection, splenectomy, distal pancreatectomy, gastric resection, and porto-celiac-cardiophrenic lymphadenectomy.^{1,2} Extensive surgery may be associated with significant morbidity, such

as biliary duct injury, biliary leaks, bleeding, pancreatic fistula formation, ureteric and bladder injuries, and fistula.^{3,4} Precise surgery based on anatomical dissection planes is crucial to prevent complications. In addition, early recognition and management of complications have vital importance. Close cooperation between surgeons, radiologists, and interventional radiologists is required to determine the best




Figure 1 Thorough knowledge of the liver and biliary tree anatomy, associated vascular supply, and the necessary mobilization techniques are crucial for a successful and safe cytoreduction that includes upper abdominal resection techniques.

Stomach – Gastric Perforation

- Gastroparesis is a syndrome of objectively delayed gastric emptying of solids in the absence of a mechanical obstruction.
- Main symptoms are nausea, vomiting, early satiety, bloating, and/or upper abdominal pain.
- Gastric stasis is usually due to surgical injury to the vagus nerve.
- Vagal injury can be demonstrated by measurement of the plasma pancreatic polypeptide response to modified sham feeding.

ESGO eAcademy
Gynecologic Oncology
Cagatay Taskiran

13

@IJGOnline

Copyright © 2020 BMI Publishing Group Ltd, International Gynecologic Cancer Society, & European Society of Gynaecological Oncology. All rights reserved.

Video 1 Thorough knowledge of the liver and biliary tree anatomy, associated vascular supply, and the necessary mobilization techniques are crucial for a successful and safe cytoreduction that includes upper abdominal resection techniques.

treatment modality and to avoid unnecessary surgical interventions in patients that can be solved endoscopically or with imaging guidance (Figure 1).

Recent advances in interventional radiology have facilitated the management of surgical complications with reduced morbidity. Biliary leaks and pancreatic fistulas can be resolved with radiological drains, and also stenting of ureters is indicated in fistulas to induce healing. A multidisciplinary approach is mandatory to achieve the best outcome and to avoid detrimental long term impacts on the quality of life and functionality of patients .

Author affiliations

¹Department of Obstetrics and Gynecology, Chair, Koc University School of Medicine and VKV American Hospital, Istanbul, Turkey

²University Oncology Center University Hospital of Bialystok, University of Bialystok Institute of History and Political Sciences, Bialystok, Podlaskie, Poland

³Gynaecologic Oncology, Imperial College London Faculty of Medicine, London, UK

Presented at

Published in partnership with the European Society for Gynecologic Oncology and BMJ

Contributors All of the authors collected the data, wrote the article, and produced the videos.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

Data availability statement Data are available in a public, open access repository.

ORCID iDs

Cagatay Taskiran <http://orcid.org/0000-0002-0936-552X>

Christina Fotopoulou <http://orcid.org/0000-0001-6375-9645>

REFERENCES

- 1 Aletti GD, Dowdy SC, Gostout BS, *et al*. Aggressive surgical effort and improved survival in advanced-stage ovarian cancer. *Obstet Gynecol* 2006;107:77–85.
- 2 Bund V, Lecoindre L, Velten M, *et al*. Impact of lymphadenectomy on survival of patients with serous advanced ovarian cancer after neoadjuvant chemotherapy: a French national multicenter study (FRANCOGYN). *J Clin Med* 2020;9.
- 3 Tozzi R, Traill Z, Garruto Campanile R, *et al*. Porta hepatis peritonectomy and hepato-celiac lymphadenectomy in patients with stage IIIC-IV ovarian cancer: diagnostic pathway, surgical technique and outcomes. *Gynecol Oncol* 2016;143:35–9.
- 4 Martinez A, Pomel C, Mery E, *et al*. Celiac lymph node resection and porta hepatis disease resection in advanced or recurrent epithelial ovarian, fallopian tube, and primary peritoneal cancer. *Gynecol Oncol* 2011;121:258–63.