




Peri-operative ovarian cancer guidelines: psycho-oncology

Annette Hasenburg,¹ Jalid Sehouli,² Christina Fotopoulou ³

Christina Fotopoulou

³Gynaecologic Oncology, Imperial College London Faculty of Medicine, London, UK



Biography: Christina Fotopoulou is the Professor of Gynaecological Cancer Surgery in the Department of Surgery and Cancer, Faculty of Medicine of Imperial College London, UK. She is the Deputy director of the Ovarian Cancer Action Research Centre at Imperial College. She holds an honorary chair in the Gynaecology Department at the Charite' University of Berlin, where she was trained and then later took the role of the Vice Director of the Gynecological Department. Her surgical and scientific expertise focuses on the management of patients with advanced and relapsed ovarian cancer, profiling of tumor heterogeneity and integration of tumor biology factors with surgical effort under the umbrella of individualization of surgical care. She has served as the Chair of the guidelines committees of the British Gynaecological Cancer Society (BGCS) and of ESGO (European Society of Gynaecologic Oncology). She has been an elected member of the ESGO Council and is also a member of the German AGO-Ovarian Cancer Group. She is on the editorial board and reviewer of numerous international gynaecological and oncological journals and is a member of various international oncological committees, including BGCS, ASCO, ESGO, IGCS, ESMO, ENGOT, AGO, SGO and NOGGO.

For numbered affiliations see end of article.

Correspondence to

Professor Christina Fotopoulou, Gynaecologic Oncology, Imperial College London Faculty of Medicine, London W12 0HS, UK; chfotopoulou@gmail.com

For 'Presented at statement' see end of article.

Accepted 23 August 2022
Published Online First
3 October 2022

The diagnosis of a life-threatening disease such as cancer is associated with enormous distress, which can manifest at the somatic (eg, fatigue, pain) and psychological levels with significant social/financial implications as well as spiritual and existential concerns.¹

Holistic support with evaluation of psychological distress symptoms, sexual function, psychiatric comorbidities, and psychosocial needs should therefore be offered at the time of diagnosis, during

treatment, and at follow-up and survivorship to all affected patients (Figure 1).

Treatment approaches should be tailored to individual needs and availability of interventions.

The National Comprehensive Cancer Network (NCCN) distress thermometer serves as an easily manageable first-stage screening tool to evaluate a patient's distress in areas such as practical, family, emotional, spiritual, sexual, and physical problems.² A cut-off of >4 is recommended

Evaluation for distress, sexual dysfunction and psychiatric comorbidity

- Screening time points:
 - at the time of diagnosis,
 - during treatment, follow-up and survivorship
- Screening tools
 - NCCN Distress Thermometer (cut off ≥ 4 : elevated level of distress)
 - Hospital Anxiety and Depression Scale (HADS)
 - Patient Reported Outcomes (PROMS)
- Survivorship care plan




Figure 1 Standardized evaluation tools for the peri-operative psychosexual assessment of patients with advanced ovarian cancer are now an integral part of peri-operative algorithms in an effort to routinely establish more holistic care for these patients.



© IGCS and ESGO 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Hasenburg A, Sehouli J, Fotopoulou C. *Int J Gynecol Cancer* 2022;32:1626–1628.

Lenovo  <

Breaking Bad News

- Setting up the Interview
- Patient's Perception
("before you tell: ask!")
- Patient's Invitation
- Giving Knowledge and Information
- Addressing the Patient's Emotions
- Strategy and Summary



©Sehouli 2021/Charité Berlin

www.esgo.org CHARITÉ

11



@IJGOnline

Copyright © 2020 BMJ Publishing Group Ltd, International Gynecologic Cancer Society, & European Society of Gynaecological Oncology. All rights reserved.

Video 1 Standardized evaluation tools for the peri-operative psychosexual assessment of patients with advanced ovarian cancer are now an integral part of peri-operative algorithms in an effort to routinely establish more holistic care for these patients.

to identify patients with clinically elevated levels of distress.³ Further scales such as the Hospital Anxiety and Depression Scale and the Female Sexual Function Index can supplement the diagnostic process and identify areas of need. Patient-reported outcomes may help to monitor treatment side effects.⁴ A survivorship care plan may support the patient to organize his/her life with and after cancer.

Patients with a low level of distress should be offered patient-orientated information and psychosocial consultation.⁵ Patients with a high level of distress should be seen by specialized caregivers to provide high standard psycho-oncological and psychosocial support. An armamentarium of interventions including counseling, psychoeducation, dignity-based therapy, relaxation, all creative therapies including art, music, creative writing and movement therapies, and guided imagery techniques can help to reduce patients' anxiety and improve their quality of life.^{6,7}

Love, affection, and sexuality are essential elements of life. The type and radicality of surgical treatment influence sexual function and overall quality of life, which are—in any case—often impaired by the cancer diagnosis itself, with subsequent

hormonal imbalance that may additionally be potentiated by the side effects of systemic therapy.

Almost 50% of women with cancer lack adequate information about sexual function and associated challenges during their treatment journey, such as vaginal dryness, dyspareunia, and impairment of orgasm. Therefore, prior to surgery and during treatment the patient and, where applicable, her partner should be appropriately counseled regarding potential sexual dysfunction and options of support.

Communication is one of the backbones of such a holistic care approach and of a balanced doctor–patient relationship. In this context, all members of the medical staff as well as the relatives should be involved. Breaking bad news, especially in the context of surgical complications, is a major challenge in the clinical routine that needs adequate training.

Various tools and techniques, including the 'SPIKES' model (Setting up, Perception, Invitation, Knowledge, Emotions with Empathy, and Strategy or Summary), are available to teach and learn verbal and non-verbal communication.

Educational video lecture

All the above strategies will need to be under a holistic care umbrella to fit the patient's need in an individualized approach, and also to adapt to the infrastructure and support of each given environment.

Author affiliations

¹Department of Gynecology and Obstetrics, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

²Gynecology with Center of Oncological Surgery, Charite Universitätsmedizin Berlin, Berlin, Germany

³Gynaecologic Oncology, Imperial College London Faculty of Medicine, London, UK

Presented at

Published in partnership with the European Society for Gynecologic Oncology and BMJ

Contributors All authors collected data, wrote the statements and did the videos.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

Data availability statement Data are available in a public, open access repository.

ORCID iD

Christina Fotopoulou <http://orcid.org/0000-0001-6375-9645>

REFERENCES

- 1 Mehnert A, Hartung TJ, Friedrich M, *et al*. One in two cancer patients is significantly distressed: prevalence and indicators of distress. *Psychooncology* 2018;27:75–82.
- 2 Donovan KA, Grassi L, McGinty HL, *et al*. Validation of the distress thermometer worldwide: state of the science. *Psychooncology* 2014;23:241–50.
- 3 Ma X, Zhang J, Zhong W, *et al*. The diagnostic role of a short screening tool--the distress thermometer: a meta-analysis. *Support Care Cancer* 2014;22:1741–55.
- 4 Snaith RP. The Hospital Anxiety and Depression Scale. *Health Qual Life Outcomes* 2003;1:29.
- 5 AWMF. Psychoonkologische Diagnostik, Beratung und Behandlung von erwachsenen Krebspatienten (in German). Registernummer: 032/051OL; 2014. <https://www.awmf.org/en/clinical-practice-guidelines/detail/1/032-051OL.html>
- 6 Hartung TJ, Fried EI, Mehnert A, *et al*. Frequency and network analysis of depressive symptoms in patients with cancer compared to the general population. *J Affect Disord* 2019;256:295–301.
- 7 Lee J, Choi MY, Kim YB, *et al*. Art therapy based on appreciation of famous paintings and its effect on distress among cancer patients. *Qual Life Res* 2017;26:707–15.