

2022-RA-1131-ESGO


HUMAN PAPILLOMAVIRUS GENOTYPE AND PROGNOSTIC FACTORS OF VAGINAL CANCER

¹Chyong-Huey Lai, ²Hsiu-Jung Tung, ³You-Chen Wang, ³Min-Jie Liao, ⁴Shih-Ming Jung, ⁵Lan-Yan Yang. ¹Obstetrics and Gynecology, Chang Gung Memorial Hospital, Linkou Branch and Chang Gung University, Taoyuan City, Taiwan; ²Department of Obstetrics and Gynecology, Chang Gung Memorial Hospital, Linkou Branch and Chang Gung University, Taoyuan city, Taiwan; ³Department of Obstetrics and Gynecology, Chang Gung Memorial Hospital, Keelung Branch, Keelung city, Taiwan; ⁴Department of Pathology, Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Taoyuan City, Taiwan; ⁵Clinical Trial Center, Chang Gung Memorial Hospital Linkou Branch, Taoyuan City, Taiwan

10.1136/ijgc-2022-ESGO.950

Introduction/Background The natural history of invasive vaginal cancer has been minimally investigated. We aimed to investigate HPV distribution and prognostic factors in vaginal cancer (VC).


Methodology We retrospectively reviewed medical records of patients with VC who received primary treatment between 1989 and 2020. General polymerase chain reaction (PCR) SPF1/GP6+ followed by revert-blot detection was performed for human papillomavirus (HPV) genotyping. E6 type-specific PCR of the top-5 prevalent types was performed to reconfirm HPV-negative status. P16INK4a immunohistochemistry staining was performed. Univariate and multivariate analyses were performed to identify predictors of clinical outcomes.



Abstract 2022-RA-1131-ESGO Figure 1

Results A total of 73 vaginal carcinoma patients eligible for analysis. Median follow-up time was 88.6 months (range 0.56–239.5 months). 66 patients (90.4%) were diagnosed as squamous cell carcinoma (SCC) and 7 (9.6%) as non-SCC. HPV DNA sequences were detected in 88.7% of SCC specimens, and 83.3% of non-SCC VC ($P = 0.543$). The leading types were HPV16 (51.7%), HPV52 (13.3%) and HPV58 (11.7%). HPV-positivity was associated with better 5-year cancer-specific survival (CSS) (70.8% vs 35.7%, $P = 0.005$). Because there was strong correlation between p16-positivity and HPV-positivity ($P < 0.001$), they were alternatively entered in multivariate analysis. In both models, pelvic lymph node (PLN) metastasis (HR 4.72, 95%confidence interval [CI] 1.505–14.804, $P =$

0.008 or 6.35, 95%CI 1.871–21.564, $P = 0.003$) was a significant adverse predictor of CSS, while p16 (HR 0.049, 95% CI 0.01–0.229, $P < 0.001$) or HPV-positivity (HR 0.129, 95%CI 0.036–0.466, $P = 0.002$) was related to better prognosis. International Federation of Gynaecology and Obstetrics stage (III/IV vs I/II) was significant in univariate analysis, but was not significant in either model.



Abstract 2022-RA-1131-ESGO Figure 2

Conclusion PLN metastasis was a significant adverse predictor, while p16-positivity or HPV-positivity (alternatively) was a significant factor of better prognosis.

2022-VA-1138-ESGO

LAPAROSCOPIC ANTERIOR RESECTION WITH TOTAL VAGINECTOMY

VN Raju K, Pavan kumar Jonnada, Zeeba Usofi, B Madhu, K Pradeep, Syed Nusrath. BIACHRI, hyd, India

10.1136/ijgc-2022-ESGO.951

Introduction/Background This is a video demonstration of laparoscopic anterior exenteration with total vaginectomy for vaginal squamous cell cancer in a young female.

Methodology Techniques of dissection and surgical demonstration of anterior exenteration with total vaginectomy

Results Video demonstrated in the given format

Conclusion laparoscopic anterior exenteration with total vaginectomy is an acceptable surgical procedure with minimal morbidity for the management of locally advanced vaginal cancer.

2022-VA-1152-ESGO

SIGMOID NEOVAGINA AND DOUBLE V-Y FLAP RECONSTRUCTION AFTER A TOTAL PELVIC INFRALEVATOR EXENTERATION

¹Felix Boria, ²Enrique Chacón, ³Monica Gutierrez, ²Nabil Manzour, ³Teresa Castellanos, ³Daniel Vazquez, ³Luis Chiva. ¹Gynecologic oncology department, Clinica universidad de Navarra, MADRID, Spain; ²Gynecologic oncology, Clinica Universidad de Navarra, Pamplona, Spain; ³Gynecologic oncology, Clinica Universidad de Navarra, Madrid, Spain

10.1136/ijgc-2022-ESGO.952