

study series, supporting the clinician to integrate output analysis (Morphonode-SP).

**Conclusion** Our findings indicate that Morphonode Predictive Model is a simple and observer-independent tool. It could be easily integrated in the clinical routine for preoperative stratification of vulvar cancer patients.

**2022-RA-1300-ESGO** **MULTICENTER EXPERIENCE ON SENTINEL NODE MAPPING IN VULVAR MELANOMA EVALUATION OF CLINICAL IMPACT**

<sup>1</sup>Simona Maria Fragomeni, <sup>2</sup>Angela Collarino, <sup>3</sup>Valentina Fuoco, <sup>4</sup>Fabio Martinelli, <sup>5</sup>Tina Pasciuto, <sup>1</sup>Giacomo Corrado, <sup>4</sup>Francesco Raspagliesi, <sup>2,6</sup>Vittoria Rufini, <sup>3</sup>Marco Maccauro, <sup>1,7</sup>Giovanni Scambia, <sup>7,8</sup>Giorgia Garganese. <sup>1</sup>Unità di Ginecologia Oncologica, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; <sup>2</sup>Nuclear Medicine Unit, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; <sup>3</sup>Department of Nuclear Medicine, Fondazione IRCCS Istituto Nazionale Tumori di Milano, Milan, Italy; <sup>4</sup>Department of Gynecologic Oncology, Fondazione IRCCS Istituto Nazionale Tumori di Milano, Milan, Italy; <sup>5</sup>Dipartimento Scienze della Salute della Donna, del Bambino e di Sanità Pubblica, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy; <sup>6</sup>Institute of Nuclear Medicine, Università Cattolica del Sacro Cuore, Rome, Italy; <sup>7</sup>Dipartimento Universitario Scienze della Vita e Sanità Pubblica Sezione di Ginecologia ed Ostetricia, Università Cattolica del Sacro Cuore, Rome, Italy; <sup>8</sup>Gynecology and Breast Care Center, Mater Olbia Hospital, Olbia, Italy

10.1136/ijgc-2022-ESGO.962

**Introduction/Background** Melanoma of the vulva is a rare disease, often burdened by a poor prognosis. It is essential to define the optimal treatment in early stage disease. This multicenter retrospective study investigates the role of preoperative lymphoscintigraphy and sentinel node biopsy (SNB) and the impact of SNB on loco-regional control and survival in vulvar melanoma patients with clinically negative nodes (cN0).

**Methodology** All women treated between July 2013 and March 2021 were evaluated. Inclusion criteria consisted in: (i) histologically proven vulvar invasive melanoma, (ii) a Breslow tumor thickness of 1–4 mm and (iii) cN0 at preoperative evaluation. Patients selected underwent a preoperative lymphoscintigraphy followed by SNB with or without inguinofemoral lymphadenectomy. DFS and OS were assessed by the Kaplan-Meier method.

**Results** Eighteen women were included for a total of 28 groins studied. Planar images showed 51 sentinel nodes (SNs) in the enrolled inguinal regions. SNs were identified in all cases. Metastatic SNs were found in 5 patients (27,7%) for a total of 8 metastatic nodes in 7 groins (25%). Recurrent disease was diagnosed in 10 (55,5%) patients at 3 to 30 months: 7 were SN-negative, among which no specific groin recurrence was observed; 3 were SN-positive, among which 2 patients died of disease after 26.2 and 33.8 months, respectively. The overall mortality rate was 0% for SN negative and 40% in SN positive patients. OS and DFS at 36 months were 62.5% and 19.2%, respectively. The median DFS was 18.0 months (95% CI, 10.3–30.0).

**Conclusion** Lymphoscintigraphy followed by sentinel lymph node biopsy in patients with vulvar melanoma is feasible and allows adequate assessment of the stage of disease. Negative SNB is associated with low risk of groin relapse and good survival rate. Further prospective multicenter studies are needed to evaluate the criteria for clinical application.

**2022-RA-1303-ESGO** **ANALYSIS OF HOSPITALIZATION OF PATIENTS WITH MALIGNANT VULVAR DISEASES OPERATED BETWEEN 02.2020 AND 04. 2022 IN THE DEPARTMENT OF ONCOLOGICAL GYNECOLOGY OF THE CENTRE OF ONCOLOGY, OPOLE, POLAND**

Marcin Kalus, Krzysztof Nowak, Maja Mrugala, Zofia Borowiec, Ewa Milnerowicz-Nabzdzyk. Department of Oncological Gynecology, Dept Director Associate Prof. Ewa Milnerowicz-Nabzdzyk MD PD, Centre of Oncology, Opole, Poland

10.1136/ijgc-2022-ESGO.963

**Introduction/Background**

**The objective** the analysis of patients with V-Y technique vulvar reconstruction in the treatment of malignant vulvar diseases between 02.2020 and 04. 2022 in the Department of Oncological Gynecology

**Methodology** The surgical treatment was performed in 25 patients: 21 with squamous cell carcinoma and 4 with vulvar Paget disease. 18 women underwent complete radical vulvectomy and 7 partial radical vulvectomy (hemivulvectomy or wide tumor resection). In the group with complete radical vulvectomy 6 patients were qualified to the V-Y technique vulvar reconstruction. In 12 patients the systemic inguinal lymphadenectomy (uni- or bilateral) was performed, of which 5 was diagnosed with node metastases. In 4 patients the SLN procedure was performed, with positive results (lymph node metastases) in 2 patients. 2 patients after the simultaneous reconstruction group required secondary wound suturing in the postoperative period. In the group of patients after partial or complete radical vulvectomy without the reconstruction 6 patients needed secondary surgical intervention.

**Results** The average operating time without the simultaneous reconstruction was 160 min. and with the reconstruction was 205 min. The average healing time without the reconstruction was 30 days and with the reconstruction 23 days. The mean satisfaction index without the reconstruction was 6/10 and with the reconstruction 8/10.

**Conclusion** 1. The surgical procedures in the vulvar malignant diseases with simultaneous reconstruction are more time-consuming but more effective for the patient's quality of life than the procedures without reconstruction. 2. Secondary wound suturing in the postoperative period had no impact on the patient's final treatment assessment. 3. The number of patients treated with reconstructive procedures is an obvious limitation of this analysis, but it should be considered as a pilot study. The number of patients enrolled will be growing and more detailed meta-analysis is planned for the following years.

**2022-RA-1376-ESGO** **INTRODUCTION OF CARE BUNDLE IN VULVAR CANCER**

Fatma Mohamed AlWahaibi, Heather Agnew, Ugochukwu Umeanozie, Ian Harley, Stephen Dobbs, Hans Nagar, Elaine Craig, Mark McComiskey. Gynecology Oncology, Belfast City Hospital, Belfast, UK

10.1136/ijgc-2022-ESGO.964

**Introduction/Background** Surgery for vulvar cancer (VC) is associated with high morbidity. Design of 'care bundle' of