

2022-RA-1559-ESGO

# EXPERT ULTRASOUND EXAMINATION, MRI OR ROMA FOR DISCRIMINATING BENIGN FROM MALIGNANT IN INCONCLUSIVE ADNEXAL MASSES AS DETERMINED BY IOTA SIMPLE RULES

<sup>1</sup>Julio Vara, <sup>2</sup>Enrique Chacón, <sup>3</sup>Isabel Brotons, <sup>2</sup>Ana López-Picazo, <sup>2</sup>Juan González Canales, <sup>2</sup>Alba Etxeandia, <sup>3</sup>Teresa Castellanos, <sup>3</sup>Lucía Pérez Alonso, <sup>3</sup>Felix Boria, <sup>2</sup>Nabil Manzour, <sup>3</sup>Isabel Carriles, <sup>4</sup>M Ángela Pascual, <sup>2</sup>María Arraiza, <sup>5</sup>Stefano Guerriero, <sup>3</sup>Luis M Chiva, <sup>2</sup>Juan Luis Alcázar. <sup>1</sup>Obstetrics And Gynaecology, Clínica Universidad de Navarra, Pamplona, Spain; <sup>2</sup>Clínica Universidad de Navarra, Pamplona, Spain; <sup>3</sup>Clínica Universidad de Navarra, Madrid, Spain; <sup>4</sup>Institut Universitari Dexeus, Barcelona, Spain; <sup>5</sup>University of Cagliari, Cagliari, Italy

10.1136/ijgc-2022-ESGO.180

**Introduction/Background** To determine which would be the best second step approach for discriminating benign from malignant adnexal masses classified as inconclusive by IOTA Simple Rules (SR).

**Methodology** Single center prospective study performed (January 2018-December 2021) comprising a consecutive series of patients diagnosed as having an adnexal mass classified as inconclusive according to IOTA SR by non-expert examiners. All women were underwent ROMA analysis, DC-MRI interpreted by an expert radiologist and ultrasound (US) examination by expert gynecological sonologist. Pregnant patients and patients with less than 12 months of follow-up were excluded. Cases were clinically managed according to the result of the US expert examination by either serial follow-up for at least one years or surgery. Reference standard was histology (patient was submitted to surgery if any of the tests was suspicious) or follow-up (Masses with > 12 months and no signs of malignancy were considered as benign). Diagnostic performance of all three approaches were calculated and compared. Direct cost analysis of the test used was also performed.

**Results** 80 women were included. Seventeen patients were managed expectantly and 63 patients underwent surgery. 23 masses were malignant. Diagnostic performance of all three approaches is shown in table. Both US expert examination and MRI had significantly better diagnostic performance than ROMA. There was no difference in terms of diagnostic performance between US and MRI. Direct costs were significantly lower for US than for MRI and similar to ROMA.

**Abstract 2022-RA-1559-ESGO Table 1** Diagnostic performance of ROMA, MRI, Expert US examination

Method	Sensitivity	Specificity
ROMA	26%	93%
MRI	91%	77%
Expert US	100%	91%

**Conclusion** US expert examination is the best second step approach in inconclusive adnexal masses as determined by IOTA Simple Rules.

2022-RA-1563-ESGO

# PRE-OPERATIVE ULTRASOUND ASSESSMENT OF RECTOSIGMOID INFILTRATION IN ADVANCED OVARIAN CANCER

<sup>1</sup>Silvia Gómez Carballo, <sup>1</sup>Claudia Pumarola, <sup>2</sup>Ariel Glickman, <sup>2</sup>Núria Carreras, <sup>2</sup>Agustí Núria, <sup>3</sup>Ana Luzárraga, <sup>4</sup>Pere Fusté, <sup>4</sup>Aureli Torné, <sup>1</sup>Berta Díaz-Feijoo, <sup>1</sup>Meritxell Munmany Delgado. <sup>1</sup>Hospital Clínic de Barcelona, Barcelona, Spain; <sup>2</sup>ICGON. Hospital Clínic de Barcelona, Barcelona, Spain; <sup>3</sup>Hospital Vall d'Hebron, Barcelona, Spain; <sup>4</sup>Ginecología y Obstetricia, ICGON. Hospital Clínic de Barcelona, Barcelona, Spain

10.1136/ijgc-2022-ESGO.181

**Introduction/Background** It is essential to perform a detailed preoperative description of disease's extension which can improve patient management, including preoperative work-up, operative time and postoperative care. Ultrasound (US) is a reliable method for differentiation between benign and malignant adnexal tumors and for local staging of endometrial and cervical cancers. Few studies have pointed the use of US evaluating the extent of disease in advanced ovarian cancer and evaluation of operability. The objective of this study is to assess the accuracy of US predicting rectosigmoid tumor infiltration in patients with advanced ovarian cancer.

**Methodology** This observational prospective study includes 55 patients with an US diagnosis of adnexal mass suspected of malignancy which was confirmed histologically. 39 patients underwent primary surgery and 16 interval surgery. US was performed to assess disease's extension. Rectosigmoid infiltration was evaluated by perioperative findings.

**Results** Rectosigmoid infiltration was confirmed in 36 patients. Rectosigmoid resection was performed in 12 cases and visceral peritoneum stripping in 3. In the other 21 cases bowel surgery was not performed due to unresectable disease. Rectosigmoid carcinomatosis was correctly detected by US in 24/36 patients. In 9/36 it was not detected and in 3/36 rectosigmoid wall was not assessable. In 2/24 cases miliary carcinomatosis was identified and 22/24 had nodular carcinomatosis with a nodule mean diameter of 26 mm. In 23/24 there was a Douglas lock. The Sensitivity of US in detecting rectosigmoid carcinomatosis was 72.7%, and specificity was 93.7%. Positive predictive value of 96% and negative predictive value of 62.5%. The absence of ascites, high BMI, dimensions of adnexal mass and abundant bowel content could affect the accuracy of US.

**Conclusion** US is an accurate method for the pre-operative assessment of rectosigmoid infiltration in advanced ovarian cancer and it can be used for adequately preoperative planning and predict need of surgery on rectosigmoid carcinomatosis.

2022-VA-1642-ESGO

# TRICKS TO IMPROVE THE LAPAROSCOPIC EXTRAPERITONEAL SPACE IN PARA-AORTIC LYMPHADENECTOMY

<sup>1</sup>Maria Carbonell Lopez, <sup>2</sup>Myriam Gracia, <sup>2</sup>Virginia Garcia, <sup>2</sup>Jaime Siegrist, <sup>2</sup>Elena Rodriguez Gonzalez, <sup>2</sup>Maria Alonso, <sup>2</sup>Maria Dolores Diestro, <sup>2</sup>Alicia Hernandez, <sup>2</sup>Ignacio Zapardiel. <sup>1</sup>Gynecology, Hospital Universitario La Paz, Madrid, Spain; <sup>2</sup>Hospital Universitario La Paz, Madrid, Spain

10.1136/ijgc-2022-ESGO.182

**Introduction/Background** Minimal invasive surgery in gynecological cancer offers benefits over laparotomy in terms of fewer operative complications. There are two approaches to para-aortic lymphadenectomy: transperitoneal and extraperitoneal. The transperitoneal approach offers a greater working space and familiar landmarks, but sometimes requires bowel mobilization. The advantages of the extraperitoneal approach include operative feasibility in spite of previous abdominal surgery, decreased risk of direct bowel injury, and bowel adhesion formation. The disadvantages are a small working space, limited landmarks, and the risk of becoming disoriented. The use of some techniques to increase the surgical field may be helpful by making surgery easier and faster.

**Methodology** We present a video with four surgical techniques to improve the viewing area in extraperitoneal para-aortic lymphadenectomy.

**Results** Accessory trocar for instrument insertion to raise the upper peritoneum in the form of a tent.- Placement of a clamp on the umbilical trocar placed in the peritoneal cavity to facilitate the outflow of CO<sub>2</sub> to allow further distension of the retroperitoneal area.- Pneumatic balloon or Foley catheter can be placed to prevent the escape of CO<sub>2</sub> into the intraperitoneal space in case of accidental opening of the peritoneum during entry into the retroperitoneal field.- For advanced surgeons, node dissection can be performed with an advanced sealing instrument with one hand while the other hand is used to lift the upper peritoneum in a tent to increase the working space.

**Conclusion** Laparoscopic para-aortic lymphadenectomy is a procedure with technical difficulties. The most important and basic requirements for appropriate lymphadenectomy are a correct surgical field development and a precise knowledge of anatomy to prevent accidental injuries. The use of some tricks can help to improve the surgical field to facilitate the surgical procedure.

2022-RA-1650-ESGO

#### SONOGRAPHIC ASSESSMENT OF FEATURES SUSPICIOUS OF UTERINE SARCOMA: EVALUATION OF THEIR USE IN PREOPERATIVE PREDICTION OF MALIGNANCY

<sup>1</sup>Alexandra M Knipprath-Mészáros, <sup>1</sup>Alessandra Tozzi, <sup>2</sup>Annkatrin Butenschoen, <sup>2</sup>Hubertina Reina, <sup>3</sup>Andreas Schoetzau, <sup>1</sup>Viola Heinzelmann-Schwarz, <sup>2</sup>Gwendolin Manegold-Brauer. <sup>1</sup>Gynecological Cancer Center, University Hospital Basel, Basel, Switzerland; <sup>2</sup>Department of gynecologic ultrasound and prenatal diagnostics, University Hospital Basel, Basel, Switzerland; <sup>3</sup>Ovarian Cancer Research, Department of Biomedicine, University Hospital Basel, Basel, Switzerland

10.1136/ijgc-2022-ESGO.183

**Introduction/Background** The recognition of uterine sarcoma allows correct therapy planning and choice of surgical approach. To help to distinguish benign myoma from sarcoma, we assessed the value of 6 sonographic criteria (Sarcoma Prediction Score – SPS) in a prospective cohort of consecutive patients with uterine masses.

**Methodology** Patients planned for surgery between 2015–2019 for presumed myometrial masses were prospectively evaluated with a standardized ultrasound examination. For triage, the following criteria were investigated: rapid growth i(3 months), high blood flow, atypical growth (postmenopause), irregular

lining, central necrosis, and oval solitary lesion. The evaluation of the criteria was binary, the score could range from 0 to 6. Gold standard was histological diagnosis.

**Results** 522 myomas, 14 uterine sarcomas, 2 gastrointestinal stromal tumors in connection to the uterus, and 7 other malignancies were included. In the group of malignant tumors, 75.0% of patients were postmenopausal 25.0% premenopausal, while in the myoma group, 76.8% patients were premenopausal. The median SPS for mesenchymal tumors was 2.5 (range: 0–4, mean 2.6) vs 0 for myomas (range: 0–3, mean 0.17). The most common sonographic criteria leading to a false positive score in myomas were rapid growth and high blood flow. For the detection of sarcoma/mesenchymal tumors, at a threshold of >1, sensitivity was 93.75%, specificity 97.9%, PPV and NPV 57.7% and 99.8%, respectively. The AUC was 0.95.

**Conclusion** The use of the SPS could help to distinguish between myomas and sarcomas, with a high probability of benign histology if the score is negative. A higher risk of malignancy is given when  $\geq 1$  criteria are present in postmenopausal women. For premenopausal women, rapid growth and high blood flow may lead to false positive scores; a score  $\geq 2$  increases accuracy. We suggest the use of the SPS in the triage of patients with suspected myometrial lesions.

2022-RA-1652-ESGO

#### OVARIAN CANCER TREATMENT PLANNING AND COMPUTER TOMOGRAPHY INTERPRETATION SKILLS OF ONCOGYNAECOLOGIST

Olena Postupalenko, Kateryna Kharchenko. Department of minimally invasive surgery, Kyiv City Clinical Oncology Center, Kyiv, Ukraine

10.1136/ijgc-2022-ESGO.184

**Introduction/Background** Cytoreductive surgery is the cornerstone of modern ovarian cancer treatment. Planning and treatment assessment is very important.

**Methodology** Analyze of our experience with tips and tricks of selecting patients for cytoreductive surgery.

**Results** Radiologist's report contains the basic information about disease burden. Multiplanar review may allow clinician to imagine anatomical peculiarities of advanced disease. Sometimes it may be helpful to plan the placement of ports during diagnostic laparoscopy or to navigate during searching of suspicious areas. Patients' anatomy is easier more safely to determine preoperatively, for example variants of vessel anatomy, tumor interrelation with major vessels, ureters, spleen, pancreas etc. In the case where vascular or hepatobiliary surgeon would be needed it may be done in a planned manner, not in the emergency because of accidental intraoperative finding. After cytoreductive surgery with extensive peritonectomy some specific radiologic changes may occur. When clinician knows or at least have access to operative report, he can more correctly interpret postoperative changes (different kinds irregular soft tissue fibrosis after peritonectomy, liver changes after decapsulation or atypic resection, lymph cysts, lymphadenopathy etc.).

**Conclusion** Computer tomography interpretation skills is very important for oncogynaecologist. It should be incorporated in educational programs and training programs.