

Supplemental Table 1. Surgical procedures in overall population

Variables	All patients n (%)	Group 1 MI-SCS n (%)	Group 2 LPT-SCS n (%)	p-value <sup>a</sup>
<b>All cases</b>	276	62	214	
<b>Peritonectomy</b>	156 (56.5)	24 (38.7)	132 (61.7)	<b>0.001</b>
<b>Lymphadenectomy</b>	167 (60.5)	37 (59.7)	130 (60.7)	0.496
<b>Systematic lymphadenectomy</b>				0.816
Pelvic	36 (13.0)	10 (16.1)	26 (12.1)	
Aortic	36 (13.0)	7 (11.3)	29 (13.6)	
Pelvic & Aortic	31 (11.2)	5 (8.1)	26 (12.1)	
Mixed	5 (1.8)	1 (1.6)	4 (1.9)	
<b>Bulky lymph-node</b>				0.076
Pelvic	57 (20.7)	20 (32.3)	37 (17.3)	
Aortic	62 (22.5)	9 (14.5)	53 (24.8)	
Pelvic & Aortic	32 (11.6)	5 (8.1)	27 (12.6)	
Mixed	16 (5.8)	3 (4.8)	13 (6.1)	
<b>Splenectomy</b>	30 (10.9)	3 (4.8)	27 (12.6)	0.059
<b>Liver resection</b>	14 (5.1)	2 (3.2)	12 (5.6)	0.355
<b>Bowel resection</b>				<b>&lt;0.001</b>
Small bowel	3 (1.1)	0	3 (1.4)	
Large bowel/rectum	48 (17.4)	1 (1.6)	47 (22.0)	
Both	17 (6.2)	0	17 (7.9)	
<b>Aletti score at SCS</b>				<b>&lt;0.001</b>
1	181 (65.6)	61 (98.4)	120 (56.1)	
2	78 (28.3)	1 (1.6)	77 (36.0)	
3	17 (6.2)	0	17 (7.9)	
<b>RT at SCS</b>				<b>0.026</b>
0	262 (94.9)	62 (100.0)	200 (93.5)	
> 0	14 (5.1)	0	14 (6.5)	

<sup>a</sup>*Calculated using the Chi square test or the Fisher's exact test.*

Abbreviations: MI-SCS: minimally invasive secondary cytoreduction surgery; LPT-SCS: laparotomic secondary cytoreduction surgery; SCS= secondary cytoreduction surgery; RT= residual tumour.