

Table S2. Patients' characteristics and predictive factors of sentinel lymph node failed mapping

Study	Age (Mean ± SD or median), [years +]	BMI (Mean ± SD or median), [Kg/m ²]	Menopausal status n (%)	Adenomy- osis n (%)	Prior pelvic surgery n (%)	Prior cervical surgery n (%)	Prior cesarian section n (%)	Lysis of adhesions n (%)	Deep myometrial invasion n (%)	FIGO grade 3 n (%)	FIGO stage III-IV n (%)	Non- endometrioid histotype n (%)	LVSI n (%)	Enlarged LN n (%)	LN involvement n (%)
2020 Sozzi [19]	62.6 (26-90)	29.3 (17-52)	-	23 (6.1)	169 (44.9)	-	-	51 (13.6)	125 (33.2)	77 (20.5)	58 (15.4)	51 (13.6)	108 (28.7)	32 (8.5)	-
2019 Ianieri [16]	-	-	99 (90.0)	-	37 (33.6)	3 (2.7)	5 (4.5)	-	41 (37.3)	-	18 (16.4)	12 (10.9)	30 (27.3)	-	12 (10.9)
2019 Taskin [11]	58 (38-80)	33 (21-49)	89 (88.1)	-	-	-	-	-	21 (20.8)	21 (20.8)	19 (18.9)	11 (10.9)	20 (19.8)	-	12 (11.9)
2019 Tortorella [17]	64.2 ± 10.3	34.5 ± 8.3	-	88 (26.9)	84 (25.7)	4 (1.2)	38 (11.6)	45 (13.8)	61 (18.7)	78 (23.9)	30 (9.2)	51 (15.6)	46 (14.1)	20 (6.1)	32 (9.8)
2018 Body [15]	65.5 (49-91)	31.0 (18-56)	108 (90.8)	-	-	-	11 (9.2)	-	-	25 (21.0)	17 (14.3)	18 (15.1)	38 (31.9)	-	-
2016 Eriksson [14]	62 (30-90)	30.6 (17.7–65.3)	-	-	-	-	-	-	-	-	-	55 (17.6)	-	-	-
Total	-	-	296 (89.7)	111 (15.8)	290 (35.7)	7 (1.6)	54 (12.1)	96 (13.7)	248 (27.1)	201 (21.8)	142 (13.7)	198 (14.7)	242 (23.4)	52 (7.4)	56 (10.4)

-: not available; **BMI**: body mass index; **FIGO**: International Federation of Gynecology and Obstetrics; **LVSI**: lymphovascular space invasion; **LN**: lymph node