

#174

INTEGRATED MULTI-OMIC AND CLINICOPATHOLOGICAL ANALYSIS OF VULVAR SQUAMOUS CELL CARCINOMA: IDENTIFICATION OF PREDICTIVE BIOMARKERS FOR PERSONALIZED TREATMENT

^{1,2}Tibor A Zwimpfer*, ³Flavio Lombardo, ³Natalie Rimmer, ⁴Sandra Götze, ⁴Franziska Singer, ⁴Anne Bertolini, ¹Céline Montavon, ¹Christian Kurzeder, ³Francis Jacob, ¹Viola Heinzelmann-Schwarz. ¹University Hospital Basel, Gynecological Oncology, Basel, Switzerland; ²Peter MacCallum Cancer Centre, Cancer Research, Melbourne, Australia; ³University Hospital and University Basel, Department of Biomedicine, Basel, Switzerland; ⁴ETH Zürich, NEXUS Personalized Health Technologies, Zürich, Switzerland. SIB Swiss Institute of Bioinformatics, Zuerich, Switzerland

10.1136/ijgc-2023-ESGO.773

Introduction/Background Vulvar cancer responds poorly to systemic treatment and in contrast to other gynecological cancers, targeted therapies remain at an early stage. Interactions of clinical, immune, and molecular biomarkers remain poorly understood for this rare cancer type. Here, we characterized the proteome of vulvar squamous cell carcinoma (vSCC) and provide a computational workflow to integrate publicly available data to study this rare and aggressive cancer type.

Methodology Whole exome sequencing, bulk RNA-sequencing, clinicopathological, and proteomics data from 23 patients with vSCC were analyzed. In addition, a total of 5543 women-derived TCGA genomic and transcriptomics sequencing data were obtained for this analysis.

Results Our cohort consisted of 34.8% human papillomavirus (HPV)+ and 65.2% HPV- vSCC patients. Early FIGO stage was significantly associated with HPV- status (12% (1/8) vs. 86% (12/15), $p < 0.001$). TP53 and CDKN2 were mutually exclusive to HPV+. Patients being HPV+ revealed a higher frequency in PIK3CA alterations with 37.5% vs. 6.7% in HPV-. TYMS which is considered the primary site of action for 5-fluorouracil was significantly higher expressed in HPV+ patients ($\log_2FC = 1.7497$, $p < 0.001$). We identified unique single and doublet base substitution signatures in vSCC indicating defective DNA mismatch-repair and correlation with age. Tongue SCC (tSCC) was identified as a comparison cohort supported by different clustering methods of conjoint RNA-seq and clinicopathological data, derived from TCGA. vSCC was associated with elevated activated mast cells ($p = 0.0031$), monocytes ($p = 0.0027$), and M2 macrophages ($p < 0.001$) compared to tSCC, independent of the HPV status.

Conclusion vSCC can be further distinguished by HPV status, somatic gene alterations and tumour microenvironment, while a similar cancer tissue has been identified with tSCC. Further analysis is targeted at identifying molecular signatures serving as biomarkers in vSCC and to suggest alternative treatment options of drugs already in use in tSCC or other cancer types.

Disclosures All authors declare that they have no conflicts of interest.

#251

PATIENTS WITH NEWLY DIAGNOSED ADVANCED HIGH-GRADE SEROUS CARCINOMA WITH IMPAIRED IMMUNOSUPPRESSIVE ASCITIC MICROENVIRONMENT EXHIBIT POOR OUTCOME

¹Imane Nafia, ^{2,3}Guillaume Babin, ²Frédéric Guyon, ²Mathilde Mairé, ²Laura Poetsch Beigt, ²Sabrina Croce, ^{2,3}Cécile Hartog, ¹Assia Chaibi, ^{2,4,5}Antoine Italiano, ¹Alban Bessede, ^{2,3}Coriolan Lebreton*. ¹Explicyte Immuno Oncology, Bordeaux, France; ²Institut Bergonié, Bordeaux, France; ³ARTiSt Lab, Inserm U1312, Université de Bordeaux, Bordeaux, France; ⁴Institut Gustave Roussy, Villejuif, France; ⁵Université de Bordeaux, Bordeaux, France

10.1136/ijgc-2023-ESGO.774

Introduction/Background Despite substantial progress in high-grade serous ovarian cancers (HGSOC) in recent years, the prognosis remains poor. While ascites play a significant role in HGSOC progression, their immune landscape characterization is essential to understand their impact on the immune cell response.

Methodology Ascites were collected during initial laparoscopy for advanced HGSOC and then processed, using respective multiplexed approaches of flow cytometry-based marker expression analysis and quantitation of mediators, for profiling immune cellular and cell-free fractions. Furthermore, ascites fluids were functionally screened for their biological effects on healthy monocytes, undifferentiated and under M1-polarization.

Results 20 HGSOC patients were included from September 2020 through November 2022. Median age at diagnosis was 68 years (50–73). 19 patients had stage IIIC or IV HGSOC. 7 patients had debulking surgery (2 primary DS). After a median follow-up of 13 months (5–30), median progression-free survival was 11,0 months (95% confidence interval 7,9 – 14,1). At the time of analysis 7 patients had died.

Ascites were mostly ‘enriched’ in regulatory immune cells including T and myeloid populations. T cells highly expressed immune checkpoints e.g. PD1 in T cells and TIGIT in Tregs. CD163+ TAMs were shown to express Arg1, CCR8, CCR2, and iNOS. While acellular fluids exhibited differentially-elevated levels of soluble mediators, functionally, they polarized monocytes into M2 macrophages, and even opposed their M1 polarization to switch to M2-status.

Conclusion HGSOC ascites harbor an altered immune environment where suppressive cells and mediators lead to immune cell dysfunction. Identification of ascites subgroups most favorable to the M2 phenotype would suggest a population with poor outcomes.

These findings highlight HGSOC ascites as a valuable tool for the identification of new immuno-oncological targets.

Disclosures I. Nafia, A. Chaibi and A. Bessede are employees of Explicyte Immuno Oncology.

G. Babin received personal fees from MSD and GSK

A. Italiano received research grants from AstraZeneca, Bayer, Bristol Myers Squibb, Chugai, Merck, MSD, PharmaMar, Novartis and Roche and received personal fees from Epizyme, Bayer, Lilly, Roche and SpringWorks.

C. Lebreton received personal fees from Clovis oncology, EISAI, MSD and GSK