

had significantly higher level of satisfaction with QoL. Moreover, the regression models show that 19.2% and 14.5% of variance in QoL could be explained by nurse's perceptions toward healing environment between the two wards, respectively.

Conclusion/Implications The Chinese version ASPECT was demonstrated to be suitable as a predicative toolkit for nurses' QoL in the targeted tertiary center. Future studies might examine its applicability to patients in corresponding wards.

EP211/#383

CURRENT PERCEPTIONS OF THE ROLE OF NURSES IN CANCER CLINICAL TRIALS

¹Noriko Fujiwara*, ²Izumi Kohara, ³Shuko Tamaki, ³Midori Yabuki, ⁴Mayumi Yamamoto, ⁵Hiroko Nakahama, ⁵Chiemi Kojima, ⁶Akiko Nosaki, ⁷Shimon Tashiro, ⁸Kenichi Yoshimura, ⁹Keiichi Fujiwara. ¹The University of Tokyo, Department of Palliative Medicine and Advanced Clinical Oncology, Inmut Hospital of The Institute of Medical Science, Tokyo, Japan; ²Jichi Medical University, School of Nursing, Tochigi, Japan; ³Saitama Medical University International Medical Center, Department of Nursing, Saitama, Japan; ⁴Jichi Medical University Hospital, Department of Nursing, Tochigi, Japan; ⁵National Cancer Center Hospital, Department of Nursing, Tokyo, Japan; ⁶Chiba University, Graduate School of Nursing, Chiba, Japan; ⁷Tohoku University, Graduate School of Arts and Letters, Sendai, Japan; ⁸Hiroshima University Hospital, Future Medical Center, Hiroshima, Japan; ⁹Saitama Medical University International Medical Center, Department of Gynecologic Oncology, Saitama, Japan

10.1136/ijgc-2023-IGCS.293

Introduction Clinical (bedside) nurses play a crucial role in supporting cancer patients in making decisions regarding clinical trials, but this role is currently not being fulfilled sufficiently. The purpose of this study was to clarify the current perceptions of clinical nurses regarding their role in cancer clinical trials.

Methods Nurses who participated in a study; 'Development of Learning Program to Nurses Supporting Patients' Decision Making in Cancer Clinical Trials' were surveyed using an originally developed questionnaire (Kohara.2023). Descriptive statistics of these responses were conducted using SPSS Statistics ver. 25.

Results The analysis included 101 nurses from two university hospitals in Japan, with a median clinical nursing experience was 12 years. 51% of the nurses worked for in-patient units. About half of the nurses reported experiencing the burden of communicating with patients in clinical trials, with the main reason being their inability to explain the trial properly due to insufficient understanding (36%). Furthermore, 90% of the nurses felt a lack of knowledge about clinical trials, and the fear of being able to provide proper answers to patient-nurse relationships (75%). Only 17% of nurses had the opportunity to be involved in caring for patients and making decisions regarding their participation in cancer clinical trials in the last three months.

Conclusion/Implications Clinical nurses play an important role in supporting patients' decision-making process about participating in cancer clinical trials. However, their limited knowledge and burdens might hinder their nursing care, which calls for educational programs to improve their practice in clinical research nursing.

EP212/#606

OCCUPATIONAL SAFETY AND THERAPEUTIC EFFECT OF PACLITAXEL ACCORDING TO TYPES OF FORMULATION AEROSOLIZED DURING ROTATIONAL INTRAPERITONEAL PRESSURIZED AEROSOL CHEMOTHERAPY FOR PERITONEAL METASTASIS

¹Soo Jin Park*, ²Wongeon Jung, ³Sunwoo Park, ⁴Wonhyoung Park, ²Mijin Park, ²Chungsik Yoon, ¹Hee Seung Kim. ¹Seoul National University Hospital, Obstetrics and Gynecology, Seoul, Korea, Republic of; ²Graduate School of Public Health, Seoul National University, Department of Environmental Health Sciences, Seoul, Korea, Republic of; ³Gyeongsang National University, Department of Plant and Biomaterials Science, Jinju-si, Korea, Republic of; ⁴College of Life Sciences and Biotechnology, Korea University, Department of Biotechnology, Seoul, Korea, Republic of

10.1136/ijgc-2023-IGCS.294

Introduction To evaluate the occupation safety and effect of paclitaxel based on types of formulation aerosolized during rotational intraperitoneal aerosol chemotherapy (RIPAC) in pigs

Methods In terms of occupational safety, we first conducted RIPAC using paclitaxel twice over two days (n=2), and then performed RIPAC using paclitaxel (n=3) and polymeric nanoparticle micellar paclitaxel (PM-Pac, n=3) three consecutive times a day in eight pigs for estimating airborne and surface contamination. Moreover, we tried to make ten piglets with peritoneal metastasis (PM) using SNU-008 cells. We evaluated the pattern of PM by using the modified peritoneal cancer index (PCI) score five weeks after the first inoculation. After RIPAC only on piglets with successful PM, we compared the rate of tumor reduction between paclitaxel and PM-Pac used in RIPAC.

Results The airborne detection rate of paclitaxel was 75–100% despite no detection of PM-Pac during RIPAC. The number of airborne particles increased in the abdominal closure period during RIPAC using paclitaxel despite no increase in them during RIPAC using PM-PAC. Among surface wipe samples, the concentration above the limit of detection (LOD) was more common in paclitaxel than in PM-Pac (100% vs. 66.7% for laparoscopic instruments, P=0.03; 87% vs. 3.6% for healthcare personnel equipment). On the other hand, the modified PCI score increased after PM-Pac despite no change after paclitaxel for RIPAC in seven piglets with PM.

Conclusion/Implications PM-Pac may be safe occupationally for RIPAC, whereas it may not be effective in suppressing PM of ovarian cancer when compared with paclitaxel.

AS10. Oncologic care during and post-pandemic

EP216/#619

SAFETY OF COVID-19 VACCINATION IN GYNECOLOGIC CANCER PATIENTS AT KING CHULALONGKORN MEMORIAL HOSPITAL, THAILAND

Sasivimon Ratreer*, Natacha Phooolcharoen, Somsook Santibenchakul. Faculty of Medicine, Chulalongkorn University, Obstetrics and Gynecology, Bangkok, Thailand

10.1136/ijgc-2023-IGCS.295