

paclitaxel-carboplatin-bevacizumab chemotherapy followed by BMT between 2015 and 2021. To predict the prognosis and duration of long-term BMT (≥ 10 cycles), we calculated KELIM scores after completion of three cycles of paclitaxel-carboplatin-bevacizumab. Then, we calculated the cut-off value of the KELIM score to predict progression-free interval ≥ 12 months.

Results A total of 96 patients were included, who consisted of 28 (29%) treated with secondary cytoreductive surgery (SCS) followed by chemotherapy, and 68 (71%) treated with chemotherapy alone. The cut-off value of the KELIM score for predicting PFI ≥ 12 months was 1.08. (AUC 0.82; sensitivity, 0.75; specificity, 0.76). Although SCS did not affect progression-free survival (PFS) and overall survival (OS), high-KELIM demonstrated better prognosis than low-KELIM in patients treated with SCS (PFS, median, 13.7 vs. 15.4 mos; $p=0.04$; OS, 21.7 vs. 28.8; $p=0.006$; figure 1), whereas there was no significant difference of PFS and OS according to the KELIM score in those treated with chemotherapy alone. Moreover, high-KELIM score was a favorable factor for long-term BMT in only those treated with SCS (adjusted odd ratio, 15; 95% confidence interval, 1.225–18.363).

Conclusion/Implications High-KELIM had a predictive value for better prognosis and long-term BMT in patients with first platinum-sensitive ovarian cancer who received paclitaxel-carboplatin-bevacizumab followed by BMT after SCS.

EP249/#705

PLASMA-ACTIVATED MEDIUM INHIBITS CANCER STEM CELL-LIKE PROPERTIES AND EXHIBITS A SYNERGISTIC EFFECT IN COMBINATION WITH CISPLATIN IN OVARIAN CANCER

Woo Yeon Hwang*, So Young Kim, Sun Woo Choi, Moon Yeon Hwang, Da Young Kang, Ji Eun Jung, Byung Su Kwon, Min Hyung Jung. *Kyung Hee University Hospital, Obstetrics and Gynecology, Seoul, Korea, Republic of*

10.1136/ijgc-2023-IGCS.323

Introduction Ovarian cancer stem-like cells (CSCs) have been implicated in tumor recurrence, metastasis, and drug resistance. Accumulating evidence has demonstrated the antitumor effect of plasma-activated medium (PAM) in various carcinomas, including ovarian cancer. Thus, PAM represents a novel onco-therapeutic strategy. However, its impact on ovarian CSCs is unclear.

Methods In this study, we assessed whether PAM regulates the stemness properties in a CSC-like spheroid model. Furthermore, we investigated the potential enhanced anti-cancer effects of combination treatment with conventional chemotherapeutic agents and PAM using in vitro and in vivo models.

Results PAM exhibited synergistic cytotoxicity with cisplatin (CDDP) but not with paclitaxel and doxorubicin. In a peritoneal metastasis xenograft model established via intraperitoneal spheroid injection, PAM intraperitoneal therapy significantly suppressed peritoneal carcinomatosis (tumor size and number), with a more significant decrease observed due to the combined effects of PAM and CDDP with no side effects.

Conclusion/Implications We demonstrated the anti-CSC activity of PAM and the synergistic cytotoxic effect of the PAM and CDDP combination therapy on ovarian CSCs. Analysis of intraperitoneal PAM and CDDP combination therapy in a spheroid culture xenograft model of ovarian cancer showed promising results. Further studies are needed to determine the

molecular mechanism underlying the synergistic anti-cancer effects of PAM and anti-cancer drugs to enhance their antitumor efficacy against ovarian CSCs and reduce the relevant side effects.

EP250/#544

SYNERGISTIC EFFECT OF SHETA2 AND ABEMACICLIB IN OVARIAN CANCER SPHEROIDS AND ASCITES-DERIVED SPHEROIDS

Zitha Redempta Isingizwe*, Doris Benbrook. *University of Oklahoma Health Science Center, Stephenson Cancer Center, Oklahoma City, USA*

10.1136/ijgc-2023-IGCS.324

Introduction While recent advances in ovarian cancer therapy have improved patient's lives, current therapies are highly toxic. We developed sulfur heteroarotinoid A2 (SHetA2), a non-toxic drug currently in Phase 1 trial. SHetA2 causes cyclin D1 degradation and cyclin-dependent kinases 4 and 6 (CDK4/6) from protection by heat shock cognate 70 protein, causing G1 cell cycle arrest. Abemaciclib is a cyclin-dependent kinase (CDK) 4 and 6 inhibitor used to treat breast cancer. We hypothesized that combination of SHetA2 and abemaciclib synergistically reduces cell lines- and ascites- derived spheroids viability as both drugs target different proteins in the cyclinD1/CDK4/6 complex critical for G1 to S progression.

Methods Ovarian cancer spheroids were developed from ES2 and OVCAR8 ovarian cancer cells lines using ultra-low attachment plates. Ascites-derived spheroids were collected from ovarian cancer patients who were undergoing paracentesis for ascites removal after receiving their informed consent. The half maximal inhibitory concentration (IC_{50}) for spheroids or ascites-derived spheroids was determined using CellTiter-Glo® 3D Cell Viability Assay. GraphPad Prism was used to calculate IC_{50} for single drugs, while combination indexes were determined using CompuSyn.

Results Ovarian cancer spheroids IC_{50} values ranged between 3–10 μ M and 10–15 μ M for SHetA2 and abemaciclib, respectively. There were synergistic effects when abemaciclib and SHetA2 were combined at doses above IC_{50} of both drugs for cell line and ascites-derived spheroids.

Conclusion/Implications Findings from this study support the combination of SHetA2 and abemaciclib as a novel, less toxic therapy for ovarian cancer. Animal models will be carried out to validate these findings.

EP253/#205

DISCREPANCY IN DIAGNOSIS OF ADVANCED EPITHELIAL OVARIAN CARCINOMA, TUBAL CARCINOMA AND PRIMARY PERITONEAL CARCINOMA PRIOR TO NEOADJUVANT CHEMOTHERAPY

¹Wathirada Karnchanabanyong*, ²Malee Wannissorn, ¹Boonlert Viriyapak. ¹Faculty of Medicine Siriraj Hospital, Mahidol University, Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Bangkok, Thailand; ²Faculty of Medicine, Siriraj Hospital, Mahidol University, Department of Pathology, Bangkok, Thailand

10.1136/ijgc-2023-IGCS.325

Introduction To evaluate the rate of discrepancy between initial diagnosis and surgico-pathological diagnosis in patients treated with neoadjuvant chemotherapy (NAC) followed by interval debulking surgery (IDS) of advanced epithelial ovarian