

NRLN was stratified into 2 groups: NRLN <12 and NRLN \geq 12. We analyzed the impact of the NRLN on RFS according to clinical and pathological factors.

Results The mean age was 64.93 ± 13.817 years (range, 24–104 years). Surgery consisted of a radical vulvectomy, hemivulvectomy, and pelvic exenteration in respectively 96.4%, 2.1%, and 1.6% of cases. Lymph node (LN) dissection was bilateral in 88.5% of cases and the mean number of retrieved lymph LN was 14. LN metastasis (LNM) was assessed in 67 patients (34.9%). NRLN<12 and \geq 12 were recorded in respectively 31.8% and 68.2% of cases. After a mean follow-up time of 35.48 ± 35.48 months, the 5-year RFS in patients with NRLN<12 and NRLN \geq 12 was 38.2% and 64.9% respectively ($p=0.092$). The subgroup analysis revealed that a NRLN \geq 12 was significantly associated with a better 5 years RFS compared to NRLN<12 in stage pT1 (70.1% vs 38.9%, $p=0.016$), patients staged without LNM (81% vs 46.5%, $p=0.032$), patients with 3 or more LNM (33.6% vs 0%, $p=0.027$), in case of R0 resection (74.8% vs 36.6%, $p=0.005$) and in the absence of lymphovascular space invasion (64.9% vs 28.7%, $p=0.046$).

Conclusion/Implications The removal of more than 12 LN improves VC outcomes in patients with node-positive and negative disease, pT1 stage, and complete resection.

EP419/#629

PROGNOSTIC ANALYSIS OF FIGO STAGE I AND II OF VULVAR CANCER: A RETROSPECTIVE STUDY OF 123 CASES

¹Ines Zemni, ²Houyem Mansouri*, ¹Mohamed Ali Ayadi, ¹Amani Jellali, ¹Riadh Chargui, ¹Tarek Ben Dhiab. ¹Salah Azaiz Institute, Faculty of Medicine of Tunis, University of Tunis El Manar, Department of Surgical Oncology, Tunis, Tunisia; ²Regional Hospital of Jendouba, University of Tunis ELManar, Department of Surgical Oncology, Jendouba, Tunisia

10.1136/ijgc-2023-IGCS.460

Introduction To identify clinical, pathological features and survival predictors of vulvar cancer in patients with no lymph node metastasis.

Methods A retrospective study of 123 patients who were diagnosed and treated for vulvar cancer staged I and II at the Salah Azaiez Institute of Oncology between 1994 and 2022

Results Mean age was 65.61 ± 14.081 years (range, 30–104 years) and median follow-up was 37.84 ± 40.221 months. Surgery was a radical vulvectomy, hemivulvectomy, and pelvic exenteration in respectively 96.7%, 2.4%, and 0.8% of cases. Inguinal Sentinel lymph node (LN) biopsy was performed in 10 cases (8.1%), bilateral inguinofemoral lymphadenectomy (ILND) in 101 cases (81.1%), and unilateral (ILND) in 14 cases (11.4%). The mean tumor size was 38.04 ± 20.45 mm. Tumors were classified as stage pT1a, pT1b, pT2, and pT3 in respectively 4.9%, 81.3%, 13%, and 0.8% of cases. The 5 years overall survival was 64.2% and decreased with advanced age \geq 70 years (45% vs 74% in patients younger than 70 years, $p=0.014$) and with the presence of perineural invasion (PNI) (0% vs 70.7%, $p=0.028$). The 5-year recurrence-free survival was 65.8% and decreased with a number of retrieved LN less than 12 (0% vs 45.3% in case of 12 or more LN; $p=0.001$), and the presence of PNI (33.3% vs 71.4%, $p=0.002$). Moreover, the rate of recurrence increased with the pT stage (0%, 26%, and 52% respectively in stages pT1a, pT1b, and pT2–3; $p=0.021$).

Conclusion/Implications The survival of patients with stage I and II of vulvar cancer is correlated to the number of retrieved LN and the presence of PNI.

EP420/#482

PROGNOSTIC VALUE OF PERINEURAL INVASION IN VULVAR CANCER

¹Houyem Mansouri*, ²Ines Zemni, ²Mohamed Ali Ayadi, ²Marwa Aloui, ³Nedia Boujelbene, ²Tarek Ben Dhiab. ¹Regional Hospital of Jendouba, University of Tunis ELManar, Department of Surgical Oncology, Jendouba, Tunisia; ²Salah Azaiz Institute, Faculty of Medicine of Tunis, University of Tunis El Manar, Department of Surgical Oncology, Tunis, Tunisia; ³Salah Azaiz Institute, Faculty of Medicine of Tunis, University of Tunis El Manar, Department of Pathology, Tunis, Tunisia

10.1136/ijgc-2023-IGCS.461

Introduction Perineural invasion (PNI) is considered a poor prognostic factor in various malignant tumors, however, its predictive value in vulvar (VC) cancer remains unclear. This study aimed to determine the prognostic significance of PNI in patients with VC.

Methods We retrospectively analyzed clinicopathological data on 192 patients with VC treated surgically at the Salah Azaiez Institute of Tunisia between 1994 and 2022.

Results The mean age was 64.93 ± 13.817 years (range, 24–104 years). Surgery consisted of a radical vulvectomy, hemivulvectomy, and pelvic exenteration in respectively 96.4%, 2.1%, and 1.6% of cases. Lymph node (LN) dissection was bilateral in 88.5% of cases. The mean tumor size was 42.21 ± 24.018 mm. Tumors were classified as stage FIGO I, II, III, and IV in 55.2%, 9.4%, 32.8%, and 2.6% of cases. LN metastasis was recorded in 34.9%. Perineural invasion (PNI) and lymphovascular space invasion (LVSI) were detected respectively in 13.5% ($n=15$) and 2.7% ($n=3$) of cases. The presence of PNI was associated with young age \leq 50 years (46.7% vs 18.8%), advanced stage III–VI (80% vs 35.4%, $p=0.001$), stage LN metastasis (80% vs 35.4%, $p=0.001$) with a LN ratio \geq 0.2 (46.7% vs 12.5%, $p=0.01$) and bilateral groin metastasis (33.3% vs 13.5%, $p=0.005$), the presence of LVSI (20% vs 0%, $p<0.0001$), a decreased 5 years overall survival (OS) (12.8% vs 56.9%, $p<0.0001$) and decreased 5 years recurrence-free survival (RFS) (20.8% vs 60%, $p=0.001$).

Conclusion/Implications The presence of PNI in vulvar cancer is correlated to aggressive tumors and decreased OS and RFS

EP421/#895

CHEMORADIATION FOR ADVANCED PRIMARY VULVAR CANCER – A RETROSPECTIVE SINGLE INSTITUTION ANALYSIS

Esten Nakken*, Silje Os. Oslo University Hospital, Dept of Oncology, Oslo, Norway

10.1136/ijgc-2023-IGCS.462

Introduction Patients with advanced vulvar cancer that are considered inoperable can be cured with (chemo)radiation. We present a single institution experience with treatment outcomes and factors associated with survival.

Methods A retrospective cohort study of patients with proven carcinoma of the vulva diagnosed from 2011 to 2020 at Oslo University Hospital (Norway) is presented. Data were collected from the hospital radiation registry and medical records.

Results A total of 45 patients with inoperable vulvar cancer were included. Median age at diagnosis was 75 years (range