

Conclusion/Implications The 2023 FIGO endometrial cancer staging schema is a major revision from the 2009 FIGO schema. Almost doubled enriched sub-stages based on detailed anatomical metastatic site and incorporation of histological information enable more robust prognostication.

Early Career Workshop

W001/#1412

MOLECULAR MARKERS PERFORMED ON ENDOMETRIAL BIOPSY IN CA ENDOMETRIUM (EC) PROVIDES PROGNOSTIC AND PREDICTIVE INFORMATION

Pratima Raj*, Rupinder Sekhon, Amrita Naithani. *RGIRC, Gynaecological Oncology, New Delhi, India*

10.1136/ijgc-2023-IGCS.38

Introduction This shift towards a molecular driven Endometrial Cancer classification is an important step to the future precision medicine. These biomarkers could be used in clinical practice for a more individualized management in EC and promoting a personalized therapeutic strategy to avoid over- or under-treatment.

Aim To evaluate the role of IHC markers Preoperatively in endometrial biopsies(EB) (ER, PR, HER2, p53, L1CAM, MSI) in determining prognostication in patients with EC

Methods Observational study N= 80 patents diagnosed with endometrial cancers between September 2019- September 2021 Site-Tertiary cancer centre India IHC marker expressions in preoperative EB were correlated with post operative histopathological specimen parameters

Results Correlation of IHC marker was done with various post surgery pathology parameters and it showed correlation with variable p values suggesting that certain IHC markers correlated with advanced disease and aids in prognostication. ER and PR expression showed correlation with early disease, HER2 score 3+ showed correlations with size of the lesion and Advanced disease, L1CAM expression of >10% showed correlation with para aortic nodes and distant metastases, p53 mutation showed correlation with pelvic lymphnodal involvement and advanced disease, MMR deficient- showed

correlation with >50% myometrial invasion and no distant metastases.

Conclusions At present to our knowledge this is the first ever study evaluating the role of incorporating IHC in preoperative endometrial biopsies and correlating it with final staging.

W002/#1410

NOVEL STRATEGY OF TRAINING THE ACCREDITED SOCIAL HEALTH ACTIVISTS (ASHAS) VIA TELEMEDICINE FOR CERVICAL CANCER SCREENING BY HPV SELF-SAMPLING – THE TRACK TRIAL

¹Nilanchali Singh*, ²Pranay Tanwar, ³Sandeep Mathur, ⁴Seema Singhal, ¹Jyoti Meena, ¹Anju Singh, ¹Neerja Bhatla. ¹All India Institute of Medical Sciences, Obstetrics and Gynaecology, New Delhi, India; ²All India Institute of Medical Sciences, Laboratory Oncology, New Delhi, India; ³All India Institute of Medical Sciences, New Delhi, Department of Pathology, New Delhi, India; ⁴AIIMS, New Delhi, Gynecologic Oncology, New Delhi, India

10.1136/ijgc-2023-IGCS.39

Introduction This is a novel, pilot study aimed to analyse training of ASHAs by telemedicine for counselling women for cervical cancer screening by HPV Self-Sampling.

Methods This study is a pilot, community-based, prospective, single-arm study. Physicians trained the ASHA workers over telephone using videos, e-pamphlets and video conferencing regarding self-sampling HPV testing, who in turn trained the clients in community. Self-sampling HPV kits were transported via courier.

Results 465 women of age group 30–65 years were tested by 47 tele-trained ASHA workers. The mean age of ASHA worker and clients was 39.47±6.45 and 37.26±8.38 years, respectively. Almost half (53%) of the ASHA worker were educated till intermediate standard. The time taken to train ASHA workers via telephone was 26.51±4.18 minutes. 91.7% of the ASHA worker were satisfied with the information provided during tele-counselling. Each ASHA recruited ten clients on an average. 95.8% of the ASHAs felt it as easy to explain the clients. The acceptability of this strategy among clients was 56%. The feasibility of this strategy (percentage of clients who find it easy/those who did self-sampling) was 99%. Half of the women (48.9%) cited ‘feeling uncomfortable’ as the reason for not wanting to get screened. Among those

Abstract W002/#1410 Table 1

Characteristics	ASHA Workers (n=47)	Clients (n=465)
Age (in years)	39.47±6.45	37.26±8.38
Education Illiterate Primary Secondary Graduate Postgraduate	0 (0.0%) 16 (34.0%) 25 (53.2%) 3 (6.4%) 3 (6.4%)	85 (18.28%) 189 (40.65%) 105 (22.59%) 66 (14.19%) 20 (4.3%)
Occupation Unskilled Semi-skilled Skilled	0 47(100%) 0	378 (81.3%) 77 (16.56%) 10 (2.15%)
History of cervical cancer screening in past	1(2.1%)	36 (7.7%)
Average time taken to counsel (in minutes) Number of repeat training sessions required	26.51±4.18	NA
Acceptability of self-sampling by clients counselled by different ASHA workers (Clients willing to get HPV sample/Total number of clients counselled by each ASHA)	NA	58.25%
No of ASHA workers who could counsel following percentage of women <50% 50–90% >90%	10 17 20	NA
Feasibility (percentage of clients who find it easy/those who did self-sampling)	46 (97.9%)	461 (99.14%)
Reason of refusal of self-sampling Uncomfortable to self-sample Do not rely on test Find it difficult Feel embarrassed to do the test Just don't want to do	NA	48.9% 2.1% 4.2% 36.2% 8.6%
Wants to know result of HPV test by ASHA Physician	NA	443 (95.26%) 22 (4.73%)
Positive HPV test	5 (10.6%)	51 (11%)
Visited centre for further management	5/5 (100%)	35/51 (68.6%)