

European Menopause and Andropause Society (EMAS) and International Gynecologic Cancer Society (IGCS) position statement on managing the menopause after gynecological cancer: focus on menopausal symptoms and osteoporosis

Margaret Rees,¹ Roberto Angioli,² Robert L. Coleman,³ Rosalind M Glasspool,⁴ Francesco Plotti,² Tommaso Simoncini,⁵ Corrado Terranova²

¹John Radcliffe Hospital, Oxford, UK

²Campus Bio-Medico, University of Rome, Rome, Italy

³MD Anderson Cancer Center, Houston, Texas, USA

⁴The Beatson West of Scotland Cancer Centre, Glasgow, UK

⁵Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy

Correspondence to

Dr Margaret Rees, John Radcliffe Hospital, Oxford OX3 9DU, UK; margaret.rees@st-hildas.ox.ac.uk

This statement is being simultaneously published in *Maturitas* and the International Journal of Gynecological Cancer on behalf of the European Menopause and Andropause Society (EMAS) and International Gynecologic Cancer Society (IGCS).

Accepted 9 January 2020



© IGCS and ESGO 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Rees M, Angioli R, Coleman RL, *et al.* *Int J Gynecol Cancer* Published Online First: [please include Day Month Year]. doi:10.1136/ijgc-2020-001217

HIGHLIGHTS

- Worldwide, it is estimated that about 1.3 million new gynecological cancer cases are diagnosed each year.
- Management of menopausal symptoms and the prevention and treatment of osteoporosis in women with gynecological cancer needs to be individualized, with or without menopausal hormone therapy, according to tumor type and stage, and the woman's age.
- In women who are taking anti-estrogenic therapies such as aromatase inhibitors, estrogen-based therapies are contraindicated and non-hormonal options are recommended as initial therapy.

ABSTRACT

Worldwide, it is estimated that about 1.3 million new gynecological cancer cases are diagnosed each year. For 2018, the predicted annual totals were cervix uteri 569 847, corpus uteri 382 069, ovary 295 414, vulva 44 235, and vagina 17 600. Treatments include hysterectomy with or without bilateral salpingo-oophorectomy, radiotherapy, and chemotherapy. These can result in loss of ovarian function and, in women under the age of 45 years, early menopause. The aim of this position statement is to set out an individualized approach to the management, with or without menopausal hormone therapy, of menopausal symptoms and the prevention and treatment of osteoporosis in women with gynecological cancer. Our methods comprised a literature review and consensus of expert opinion. The limited data suggest that women with low-grade, early-stage endometrial cancer may consider systemic or topical estrogens. However, menopausal hormone therapy may stimulate tumor growth in patients with more advanced disease, and non-hormonal approaches are recommended. Uterine sarcomas may be hormone dependent, and therefore estrogen and progesterone receptor testing should be undertaken to guide decisions as to whether menopausal hormone therapy or non-hormonal strategies should be used. The limited evidence available suggests that menopausal hormone therapy, either systemic or topical, does not appear to be associated with harm and does not decrease overall or disease-free survival in women with non-serous epithelial ovarian cancer and germ cell tumors. Caution is required with both systemic and topical menopausal hormone therapy in women with serous and granulosa cell tumors because of their hormone dependence, and

non-hormonal options are recommended as initial therapy. There is no evidence to contraindicate the use of systemic or topical menopausal hormone therapy by women with cervical, vaginal, or vulvar cancer, as these tumors are not considered to be hormone dependent.

INTRODUCTION

Worldwide, it is estimated about 1.3 million new gynecologic cancer cases are diagnosed each year. For 2018 the predicted annual totals were cervix uteri 569 847, corpus uteri 382 069, ovary 295 414, vulva 44 235, and vagina 17 600.¹

Depending on tumor type and stage, treatments include hysterectomy with or without bilateral salpingo-oophorectomy, radiotherapy, and chemotherapy. These can result in loss of ovarian function and, in women under the age of 45 years, early menopause, which increases the risk not only of osteoporosis but also of cardiovascular disease and cognitive decline.^{2,3} Surgically induced menopause often leads to the immediate onset of vasomotor symptoms, which may be more severe than after natural menopause.⁴ Vasomotor symptoms may last for many years after natural or surgical menopause.^{5–7} Other symptoms, such as those related to vulvovaginal atrophy, are lifelong.^{8,9}

The management of menopausal symptoms in gynecological cancer survivors depends on their age, tumor type, and stage, as well as the use of anti-estrogen therapies (for cancers considered to

Joint statement

be hormone dependent) and concomitant morbidities. The aim of this position statement is to provide an individualized approach to the management of menopausal symptoms and the prevention and treatment of osteoporosis.¹⁰

Hormonal and Non-hormonal Management Strategies

In women without cancer, administration of systemic estrogen-based menopausal hormone therapy for menopausal symptoms and osteoporosis has a favorable risk–benefit profile for those under the age of 60 years or up to 10 years after menopause.^{8 11–14} Systemic menopausal hormone therapy can be administered orally or transdermally. Estrogen alone is given to women who have undergone hysterectomy. Progestogens and the selective estrogen receptor modulator bazedoxifene are added in regimens for women with an intact uterus to limit the increase in risk of endometrial hyperplasia and carcinoma which occurs with unopposed estrogen.^{8 15} Tibolone is a synthetic steroid compound that is, in itself, inert, but whose metabolites have estrogenic, progestogenic, and androgenic actions. It is classified as menopausal hormone therapy.¹⁶ Availability of different menopausal hormone therapy preparations varies worldwide.

In women with early or premature menopause, systemic estrogen-based menopausal hormone therapy is recommended at least until the average age of natural menopause. Anecdotally, young women may need higher doses of estrogen initially to alleviate menopausal symptoms than their older counterparts.¹² Some young women may find taking combined oral contraception more acceptable. Menopausal hormone therapy at very low doses or non-estrogen-based therapies should be considered for older women.¹² Symptoms due to vulvovaginal atrophy can be managed with low-dose topical estrogen. There are no data on the use of ospemifene or prasterone in this context.^{17 18}

The efficacy and safety of different regimens have not been examined in many studies of the use of systemic menopausal hormone therapy after gynecological cancer. While the data regarding the use of topical vaginal estrogen after gynecological cancer are sparse, it must be remembered that with current low-dose options, for example estradiol (10 µg twice weekly), absorption is very low and estrogen levels remain in the postmenopausal range.¹⁹ The total administered vaginal dose per year is similar to one daily dose of systemic oral therapy, that is, 1 mg.

In women who are taking anti-estrogenic therapies such as aromatase inhibitors, estrogen-based therapies are contraindicated.²⁰ Here, non-hormonal options are recommended as initial therapy. For vasomotor symptoms the pharmacological options include selective serotonin reuptake inhibitors and serotonin norepinephrine reuptake inhibitors, clonidine and gabapentin. Clinicians should be aware of potential drug interactions with anticancer and adjuvant therapies (eg, see refs 21 22). Cognitive behavioral therapy may also improve menopause symptoms.²³ For problems related to vulvovaginal atrophy, a variety of lubricants and bioadhesive moisturizers are available. Laser therapy for vulvovaginal atrophy is a new approach, but larger, long-term studies are required to explore its efficacy and safety before definite conclusions can be drawn.¹²

The main pharmacologic options to consider for the prevention and treatment of osteoporosis are bisphosphonates, denosumab, and parathyroid hormone.¹² As calcium and vitamin D play a key role in bone metabolism, correction of nutritional deficiencies is

advised as part of osteoporosis management.²⁴ Strategies need to be holistic and include maintaining a healthy weight, diet, exercise, and lifestyle.^{25 26} This statement will not consider herbal supplements and botanicals as there is a lack of data regarding safety and efficacy.²⁷ In addition, some products may contain compounds with estrogenic activity or may interact with anticancer therapies.

Management Options by Tumor Type

Endometrial Cancer

While most cases of endometrial cancer are diagnosed after the menopause it can occur in younger women, such as those with Lynch syndrome or polycystic ovary syndrome or who are obese. The majority of endometrial cancers are diagnosed at an early stage (Federation of Gynecology and Obstetrics (FIGO) stage I–II) and so have a good overall prognosis, with a 5-year survival rate of over 85%. Treatment usually involves hysterectomy and bilateral oophorectomy. Studies of menopausal hormone therapy after endometrial cancer are limited to one randomized trial undertaken in 1236 women recruited between 1997 and 2003 with a mean follow-up of 35.7 months²⁸ and small observational retrospective cohort or case-control studies.^{29–35} All studies were undertaken in women with early-stage disease. The randomized trial did not specify which type of menopausal hormone therapy was used (estrogen alone or estrogen plus progestogen). The observational studies documented a variety of preparations: systemic menopausal hormone therapy with estrogen alone or combined with progestogen delivered orally or transdermally, as well as topical vaginal estrogens. No studies are available for women with Lynch syndrome, who are also at increased risk of other cancers.³⁶

In 2018 a Cochrane systematic review concluded that there is insufficient high-quality evidence to inform women considering menopausal hormone therapy after treatment of endometrial cancer. However, the evidence does not suggest significant harm after surgical treatment for early-stage disease based on FIGO classification.³⁷ There is no information available regarding the use of menopausal hormone therapy in higher-stage endometrial cancer. The National Comprehensive Cancer Network Panel states that estrogen replacement is a reasonable option for patients who are at low risk of tumor recurrence, but that initiating such therapy should be individualized and discussed in detail with the patient.³⁸ Furthermore, if adjuvant treatment is carried out, there should be a 6–12-month waiting period before starting menopausal hormone therapy.

Summary Recommendation

Thus, the limited data suggest that women with low-grade, early-stage endometrial cancer may consider systemic or topical estrogens. However, menopausal hormone therapy may stimulate tumor growth in patients with more advanced disease or high-risk early-stage tumors, and non-hormonal approaches to management of menopausal symptoms are recommended. In addition, there are no long-term data regarding the safety of menopausal hormone therapy in women with Lynch syndrome, who are also at increased risk of other cancers whose treatment may lead to premature or early menopause. With regard to atypical endometrial hyperplasia, it would not be unreasonable to consider menopausal hormone

therapy in women who have undergone hysterectomy, despite the paucity of data.

Uterine Sarcoma

Stromal or mesenchymal sarcomas are rare tumors, accounting for less than 5% of all uterine cancers. While most cases are diagnosed after the menopause, these tumors can occur in younger women. The most common types are low-grade endometrial sarcomas, high-grade endometrial sarcomas, undifferentiated uterine sarcomas, and uterine leiomyosarcomas.³⁸ As these tumors may be hormone dependent, estrogen and progesterone receptor testing should be undertaken to guide decisions as to whether menopausal hormone therapy or non-hormonal strategies should be used for the management of menopausal symptoms and the prevention and treatment of osteoporosis. Low-grade stromal sarcomas may be sensitive to aromatase inhibitors or progestogens (such as megestrol acetate or medroxyprogesterone acetate). Gonadotropin-releasing hormone analogs are also an option. Randomized controlled trials have shown that progestogens are effective in treating hot flashes.^{39 40} There are no data regarding the use of menopausal hormone therapy in non-hormone-dependent tumors. In addition, there are no studies regarding the use of menopausal hormone therapy in smooth muscle tumors of uncertain malignant potential.⁴¹

Summary Recommendation

Uterine sarcomas may be hormone dependent, and therefore estrogen and progesterone receptor testing should be undertaken to guide decisions as to whether menopausal hormone therapy or non-hormonal strategies should be used. No clinical trial data are available to inform practice in women whose tumors are steroid receptor negative or who have smooth muscle tumors of uncertain malignant potential.

Ovarian, Fallopian Tube and Peritoneal Cancers

The three major types of ovarian cancer are epithelial, accounting for 90% of cases, germ cell (3%), and sex cord-stromal (2%).⁴² As fallopian tube cancer, primary peritoneal cancer, and epithelial ovarian cancer are indistinguishable and share the same genomic signature, the three are considered together.

Epithelial, fallopian tube and peritoneal cancer While these cancers often occur after the menopause, they also affect a significant number of premenopausal women.^{43–45}

Epithelial cancers are subdivided into five histotypes: high-grade serous carcinoma, low-grade serous carcinoma, endometrioid carcinoma, clear cell carcinoma, and mucinous carcinoma.^{46 47} The different histotypes are now considered to be different diseases. While serous tumors are mostly high grade, which are characterized by involvement of both ovaries, aggressive behavior, late-stage diagnosis, and low survival rates, the other subtypes tend to affect only one ovary. It is thought that serous tumors originate in the epithelial cells of the fallopian tube as microscopic preliminary lesions that subsequently migrate to the ovaries and/or peritoneum. However, endometrioid and clear cell tumors are thought to originate in the endometrium, and mucinous tumors in the ovaries or fallopian tube peritoneal junction. One of the risk factors for ovarian cancer is prior use of menopausal hormone therapy but

the association appears to be confined to serous and endometrioid histotypes.⁴⁸

Two randomized trials as well as prospective and retrospective cohort and case-control studies have shown no adverse effect menopausal hormone therapy on survival in women who have been treated for ovarian cancer.^{49–55} They used a variety of regimens: estrogen alone or combined with a progestogen or testosterone. The randomized trial by Guidozi and Daponte, in which 130 women with invasive epithelial ovarian carcinoma were followed up for 48 months, used oral continuous conjugated equine estrogen. It did not distinguish between subtypes.⁴⁹ The authors reported median overall survival of 44 months (95% CI, 10 to 112 months) and 34 months (95% CI, 8 to 111 months) in the menopausal hormone therapy and control groups, respectively. The differences in disease-free interval ($p=0.785$) and overall survival ($p=0.354$) between the two groups were not statistically significant. Eeles et al⁵⁰ studied 150 premenopausal and postmenopausal women who had been diagnosed with epithelial ovarian cancer (any FIGO stage) nine or fewer months previously. They were randomized to either menopausal hormone therapy or not for 5 years. The choice of menopausal hormone therapy for individual patients was pragmatic and was determined according to consultant preference, with guidelines to recommend that premenopausal women receive higher doses than perimenopausal/postmenopausal women. The median follow-up of patients still alive was 19.1 years: overall and relapse-free survival was greater in the menopausal hormone therapy than in the control group.

A retrospective cohort study using the Manitoba Cancer Registry and Drug Program Information Network of 357 women found that use of menopausal hormone therapy ($n=94$) for non-serous epithelial ovarian cancer was not associated with harm and did not decrease overall or disease-free survival.⁵⁵ It found that in menopausal hormone therapy users under 55 years of age, disease-free survival was longer but there was no statistical difference in overall survival for this age group. No associations between menopausal hormone therapy use and overall survival or disease-free survival were found among women aged 55 years or more.

With regard to **endometrioid ovarian cancers**, which are potentially estrogen-sensitive, menopausal hormone therapy does not appear to have adverse effects. However, while menopausal hormone therapy appears to be safe in early-stage disease, this may not be the case in women with more advanced cancers, who commonly have residual, potentially hormone-responsive disease after surgery.^{54 55} As there is no clear evidence of benefit of aromatase inhibitors in the treatment of **clear cell and mucinous carcinomas**, estrogen replacement is a reasonable option for patients who are at low risk of tumor recurrence, but initiating such therapy should be individualized. Given the benefits seen with menopausal hormone therapy with letrozole, anastrozole, tamoxifen, and leuprolide acetate after primary cytoreductive surgery and platinum-based chemotherapy in women with stage II to IV **low-grade serous carcinoma** of the ovary or peritoneum, estrogen-based therapies are currently not recommended in advanced disease of these types.⁵⁶ There is a paucity of evidence to inform practice for **high-grade serous carcinoma**.

Borderline malignant tumors or tumors of low malignant potential most often affect younger women. Histologic types include serous, mucinous, endometrioid, clear cell, and transitional

Joint statement

cell (or Brenner) tumor.⁵⁷ Five-year survival rates are greater than 98%. There is a paucity of data regarding the use of menopausal hormone therapy, but it would not be unreasonable to consider it for women with completely resected disease (ie, without invasive implants). As always, the benefits of menopausal hormone therapy for women who have undergone premature menopause through cancer treatment need to be balanced against the risks.

The **BRCA1 and BRCA2 gene mutations** are associated with increased risk of developing invasive epithelial ovarian cancer. Risk-reducing salpingo-oophorectomy is therefore recommended. However, this will lead to early/premature menopause. Data on menopausal hormone therapy after prophylactic oophorectomy are sparse, but short-term use seems to be safe.⁵⁸

Ovarian germ cell tumors commonly affect girls and young women between 10 and 30 years of age. In most cases, fertility-preserving staging surgery is followed by platinum-based combination chemotherapy, which may lead to ovarian failure. The prognosis is excellent and 5-year survival is more than 85%.⁵⁹ There is currently no evidence to suggest that these young women should not take menopausal hormone therapy.

Granulosa cell tumors are the most common ovarian sex cord stromal tumors. They secrete steroid hormones and commonly present with symptoms of hyperestrogenism, as they secrete estrogens as well as other hormones. They may have an indolent course and can recur up to 20 years after initial diagnosis. It is generally believed that estrogens should not be used, as these tumors are estrogen dependent. Hormone recurrence therapy includes aromatase inhibitors, leuprolide and tamoxifen.⁵⁹ No study, however, has demonstrated a deleterious effect of menopausal hormone therapy.

Summary Recommendation

Menopausal hormone therapy, either systemic or topical, does not appear to be associated with harm and does not appear to decrease overall or disease-free survival in women with non-serous epithelial ovarian cancer and germ cell tumors, although the evidence is limited. The regimen (estrogen or estrogen combined with progestogen) will depend on whether hysterectomy has been undertaken. Duration of therapy will depend on the age of the woman. Caution is required for both systemic and topical menopausal hormone therapy in women with serous and granulosa cell tumors because of their hormone dependence.⁶⁰

Cervical, Vaginal and Vulvar Cancers

Given that none of these cancers are considered to be hormone dependent, there is no evidence to contraindicate the use of systemic or topical menopausal hormone therapy. However, estrogen receptors are frequently (39%) expressed in cervical adenocarcinomas, even though their expression does not correlate with clinicopathologic parameters and does not influence overall and disease-free survival.^{10 61 62} In the limited studies available, no significant difference in recurrence rate or survival^{63–66} has been linked with menopausal hormone therapy use after treatment for cervical squamous cell carcinomas. In women who have been treated with radiotherapy, rather than hysterectomy, for cervical cancer, opposed estrogen therapy to prevent stimulation of residual endometrium should be used.⁶⁷

With regard to previous use of menopausal hormone therapy, a case-control study found that exogenous estrogens, especially unopposed estrogens, increased the risk of adenocarcinomas but not squamous cell carcinomas.⁶⁸ The study involved 124 women with adenocarcinomas, 139 women with squamous cell carcinomas, and 307 healthy community controls matched on age, ethnicity, and residence. Only 13 women with adenocarcinoma (10.5%), seven with squamous carcinoma (5%), and 20 controls (6.5%) had used non-contraceptive hormones; most use was short-term former use. Ever-use was associated with adenocarcinomas (OR 2.1, 95% CI 0.95 to 4.6) but not squamous carcinomas (OR 0.85, 95% CI 0.34 to 2.1). Unopposed estrogens were positively associated with adenocarcinomas (OR 2.7, 95% CI 1.1 to 6.8). However, the authors of a Women's Health Initiative study concluded that its randomized trial data on cervical cancer were too limited to suggest there was any association with estrogen plus progestin therapy use in 8506 women compared with 8102 women taking placebo.⁶⁹

As women exposed in utero in the 1950s to 1960s to diethylstilbestrol are ageing, the issue of the safety of menopausal hormone therapy in this group is becoming more important.⁷⁰ Diethylstilbestrol is associated with an increased risk of clear cell cancers of the vagina and cervix. Currently no safety data are available about the use of systemic or topical estrogen in these women.

Summary Recommendation

Menopausal hormone therapy is not contraindicated and the regimen (unopposed or opposed estrogen) depends on whether or not hysterectomy has been undertaken.

CONCLUSIONS

An individualized approach to the management of menopausal symptoms and prevention and treatment of osteoporosis after gynecological cancer is required. It should take into account age, tumor type, and stage, and concomitant therapies and morbidities. It is best undertaken by a multidisciplinary team of health and allied health professionals. It is of concern that there is a paucity of data. Therefore, there is a need for randomized trials and analysis of data registries to provide a stronger evidence base to inform practice.

Twitter Margaret Rees @emas_online and Robert L. Coleman @rcoleudede

Acknowledgements The authors thank the board and council members from both societies for their helpful comments.

Contributors MR prepared the initial draft, which was circulated to all other named authors for comments and approval before review and endorsement by the EMAS board and IGCS council members. Production was coordinated by MR.

Funding No funding was sought or secured for this position statement.

Competing interests MR reports personal fees from Sojourn, Inc., outside the remit of the submitted work. RC reports grants from NIH, grants from Gateway Foundation, grants from VFoundation, during the conduct of the study; grants and personal fees from AstraZeneca, grants from Merck, personal fees from Tesaro, personal fees from Medivation, grants and personal fees from Clovis, personal fees from Gamamab, grants and personal fees from Genmab, grants and personal fees from Roche/Genentech, grants and personal fees from Janssen, personal fees from Agenus, personal fees from Regeneron, personal fees from OncoQuest, outside the remit of the submitted work. RMG reports grants from Boehringer Ingelheim, grants from Lilly/Ignyta, personal fees, non-financial support and other from AstraZeneca, personal fees and other from Tesaro/GSK, personal fees and other from Clovis, personal fees from Sotio, personal fees from Immunogen, outside the remit of the submitted work. TS reports personal fees from Abbott, Actavis, Bayer, and

Estetra, as well as research support from Gedeon Richter, outside the remit of the submitted work. RA,FP and CT report no conflict of interest.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

Author note This statement is being simultaneously published in *Maturitas* and the *International Journal of Gynecological Cancer* on behalf of the European Menopause and Andropause Society (EMAS) and International Gynecologic Cancer Society (IGCS).

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- Faubion SS, Kuhle CL, Shuster LT, et al. Long-term health consequences of premature or early menopause and considerations for management. *Climacteric* 2015;18:483–91.
- Shuster LT, Rhodes DJ, Gostout BS, et al. Premature menopause or early menopause: long-term health consequences. *Maturitas* 2010;65:161–6.
- Rodriguez M, Shoupe D, Menopause S. Surgical menopause. *Endocrinol Metab Clin North Am* 2015;44:531–42.
- Avis NE, Crawford SL, Greendale G, et al. Duration of menopausal vasomotor symptoms over the menopause transition. *JAMA Intern Med* 2015;175:531–9.
- Stuursma A, van Driel CMG, Wessels NJ, et al. Severity and duration of menopausal symptoms after risk-reducing salpingo-oophorectomy. *Maturitas* 2018;111:69–76.
- Wilson LF, Pandeya N, Byles J, et al. Hot flushes and night sweats symptom profiles over a 17-year period in mid-aged women: the role of hysterectomy with ovarian conservation. *Maturitas* 2016;91:1–7.
- National Institute for Health and Care Excellence (NICE). Menopause: diagnosis and management NICE guideline [NG23] Published date: November 2015 Last updated: December 2019, 2015. Available: <https://www.nice.org.uk/guidance/ng23> [Accessed 7 Jan 2020].
- Panay N, Palacios S, Bruyniks N, et al. Symptom severity and quality of life in the management of vulvovaginal atrophy in postmenopausal women. *Maturitas* 2019;124:55–61.
- Angioli R, Luvero D, Armento G, et al. Hormone replacement therapy in cancer survivors: Utopia? *Crit Rev Oncol Hematol* 2018;124:51–60.
- de Villiers TJ, Hall JE, Pinkerton JV, et al. Revised global consensus statement on menopausal hormone therapy. *Maturitas* 2016;91:153–5.
- Armeni E, Lambrinoudaki I, Ceausu I, et al. Maintaining postreproductive health: a care pathway from the European Menopause and Andropause Society (EMAS). *Maturitas* 2016;89:63–72.
- Shifren JL, Gass MLS, NAMS Recommendations for Clinical Care of Midlife Women Working Group. The North American Menopause Society recommendations for clinical care of midlife women. *Menopause* 2014;21:1038–62.
- Cobin RH, Goodman NF, AACE Reproductive Endocrinology Scientific Committee. American Association of Clinical Endocrinologists and American College of Endocrinology position statement on MENOPAUSE-2017 update. *Endocr Pract* 2017;23:869–80.
- Pickar JH, Boucher M, Morgenstern D. Tissue selective estrogen complex (TSEC): a review. *Menopause* 2018;25:1033–45.
- Formoso G, Perrone E, Maltoni S, et al. Short-term and long-term effects of tibolone in postmenopausal women. *Cochrane Database Syst Rev* 2016;10.
- Di Donato V, Schiavi MC, Iacobelli V, et al. Ospemifene for the treatment of vulvar and vaginal atrophy: a meta-analysis of randomized trials. Part II: evaluation of tolerability and safety. *Maturitas* 2019;121:93–100.
- Sauer U, Talaulikar V, Davies MC. Efficacy of intravaginal dehydroepiandrosterone (DHEA) for symptomatic women in the peri- or postmenopausal phase. *Maturitas* 2018;116:79–82.
- Vagifem 10 micrograms vaginal tablets. Available: <https://www.medicines.org.uk/emc/product/5719/smpc> [Accessed 7 Jan 2020].
- Santen RJ, Stuenkel CA, Davis SR, et al. Managing menopausal symptoms and associated clinical issues in breast cancer survivors. *J Clin Endocrinol Metab* 2017;102:3647–61.
- Desmarais JE, Looper KJ. Managing menopausal symptoms and depression in tamoxifen users: implications of drug and medicinal interactions. *Maturitas* 2010;67:296–308.
- Rogala BG, Charpentier MM, Nguyen MK, et al. Oral anticancer therapy: management of drug interactions. *JOP* 2019;15:81–90.
- Atema V, van Leeuwen M, Oldenburg HSA, et al. Design of a randomized controlled trial of Internet-based cognitive behavioral therapy for treatment-induced menopausal symptoms in breast cancer survivors. *BMC Cancer* 2016;16:920.
- Cano A, Chedraui P, Goulis DG, et al. Calcium in the prevention of postmenopausal osteoporosis: EMAS clinical guide. *Maturitas* 2018;107:7–12.
- Marshall S, Rees M. Managing menopause and post-reproductive health: beyond hormones and medicines. In: Pérez-López F, ed. *Postmenopausal diseases and disorders*. Springer, 2019.
- Armstrong MEG, Lacombe J, Wotton CJ, et al. The associations between seven different types of physical activity and the incidence of fracture at seven sites in healthy postmenopausal UK women. *J Bone Miner Res* 2019;14: [e pub ahead of print]. doi:10.1002/jbmr.3896].
- About Herbs, Botanicals & Other Products Memorial Sloan Kettering Cancer Center. Available: <https://www.mskcc.org/cancer-care/diagnosis-treatment/symptom-management/integrative-medicine/herbs> [Accessed 7 Jan 2020].
- Barakat RR, Bundy BN, Spirtos NM, et al. Randomized double-blind trial of estrogen replacement therapy versus placebo in stage I or II endometrial cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2006;24:587–92.
- Ayhan A, Taskiran C, Simsek S, et al. Does immediate hormone replacement therapy affect the oncologic outcome in endometrial cancer survivors? *Int J Gynecol Cancer* 2006;16:805–8.
- Baker DP. Estrogen replacement therapy in patients with previous endometrial carcinoma. *Compr Ther* 1990;16:28–35.
- Bryant GW. Administration of estrogens to patients with a previous diagnosis of endometrial adenocarcinoma. *South Med J* 1990;83:725–6.
- Chapman JA, DiSaia PJ, Osann K, et al. Estrogen replacement in surgical stage I and II endometrial cancer survivors. *Am J Obstet Gynecol* 1996;175:1195–200.
- Creasman WT, Henderson D, Hinshaw W, et al. Estrogen replacement therapy in the patient treated for endometrial cancer. *Obstet Gynecol* 1986;67:326–30.
- Lee RB, Burke TW, Park RC. Estrogen replacement therapy following treatment for stage I endometrial carcinoma. *Gynecol Oncol* 1990;36:189–91.
- Suriano KA, McHale M, McLaren CE, et al. Estrogen replacement therapy in endometrial cancer patients: a matched control study. *Obstet Gynecol* 2001;97:555–60.
- Lynch Syndrome UK. Endometrial (uterine) cancer in Lynch syndrome. Available: <https://www.lynch-syndrome-uk.org/womb> [Accessed 7 Jan 2020].
- Edey KA, Rundle S, Hickey M. Hormone replacement therapy for women previously treated for endometrial cancer. *Cochrane Database Syst Rev* 2018;5:CD008830.
- NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines R). Uterine Neoplasms. NCCN evidence blocks TM version 4.2019 – September 16, 2019. Available: https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf [Accessed 7 Jan 2020].
- Loprinzi CL, Levitt R, Barton D, et al. Phase III comparison of depomedroxyprogesterone acetate to venlafaxine for managing hot flashes: The North Central Cancer Treatment Group Trial N99C7. *J Clin Oncol* 2006;24:1409–14.
- Goodwin JW, Green SJ, Moinpour CM, et al. Phase III randomized placebo-controlled trial of two doses of megestrol acetate as treatment for menopausal symptoms in women with breast cancer: Southwest Oncology Group Study 9626. *J Clin Oncol* 2008;26:1650–6.
- Gadducci A, Zannoni GF, Musarò A. Uterine smooth muscle tumors of unknown malignant potential: a challenging question. *Gynecol Oncol* 2019;154:631–7.
- American Cancer Society (ACS). Cancer facts and figures. special section: ovarian cancer, 2019. Available: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2018/cancer-facts-and-figures-special-section-ovarian-cancer-2018.pdf> [Accessed 7 Jan 2020].
- Liao C-I, Chow S, Chen L-may, et al. Trends in the incidence of serous fallopian tube, ovarian, and peritoneal cancer in the US. *Gynecol Oncol* 2018;149:318–23.
- Goodman MT, Shvetsov YB. Rapidly increasing incidence of papillary serous carcinoma of the peritoneum in the United States: fact or artifact? *Int. J. Cancer* 2009;124:2231–5.
- Goodman MT, Shvetsov YB. Incidence of ovarian, peritoneal, and fallopian tube carcinomas in the United States, 1995–2004. *Cancer Epidemiol Biomarkers Prev* 2009;18:132–9.

Joint statement

- 46 Kurman RJ, Carcangiu ML, Herrington CS, *et al.* *WHO classification of tumours of female reproductive organs*. 4th edn. Lyon: IARC, 2014.
- 47 Peres LC, Cushing-Haugen KL, Anglesio M, *et al.* Histotype classification of ovarian carcinoma: a comparison of approaches. *Gynecol Oncol* 2018;151:53–60.
- 48 Beral V, Gaijskell K, Hermon C, *et al.* Menopausal hormone use and ovarian cancer risk: individual participant meta-analysis of 52 epidemiological studies. *Lancet* 2015;385:1835–42.
- 49 Guidozzi F, Daponte A. Estrogen replacement therapy for ovarian carcinoma survivors: a randomized controlled trial. *Cancer* 1999;86:1013–8.
- 50 Eeles RA, Morden JP, Gore M, *et al.* Adjuvant hormone therapy may improve survival in epithelial ovarian cancer: results of the AHT randomized trial. *JCO* 2015;33:4138–44.
- 51 Eeles RA, Tan S, Wiltshaw E, *et al.* Hormone replacement therapy and survival after surgery for ovarian cancer. *BMJ* 1991;302:259–62.
- 52 Bebar S, Ursic-Vrscaj M. Hormone replacement therapy after epithelial ovarian cancer treatment. *Eur J Gynecol Oncol* 2000;21:192–6.
- 53 Ursic-Vrscaj M, Bebar S, Zakelj MP. Hormone replacement therapy after invasive ovarian serous cystadenocarcinoma treatment: the effect on survival. *Menopause* 2001;8:70–5.
- 54 Mascarenhas C, Lambe M, Bellocco R, *et al.* Use of hormone replacement therapy before and after ovarian cancer diagnosis and ovarian cancer survival. *Int J Cancer* 2006;119:2907–15.
- 55 Power L, Lefas G, Lambert P, *et al.* Hormone use after nonserous epithelial ovarian cancer: overall and disease-free survival. *Obstet Gynecol* 2016;127:837–47.
- 56 Gershenson DM, Bodurka DC, Coleman RL, *et al.* Hormonal maintenance therapy for women with low-grade serous cancer of the ovary or peritoneum. *J Clin Oncol* 2017;35:1103–11.
- 57 Gershenson DM. Management of borderline ovarian tumours. *Best Pract Res Clin Obstet Gynaecol* 2017;41:49–59.
- 58 Vermeulen RFM, Korse CM, Kenter GG, *et al.* Safety of hormone replacement therapy following risk-reducing salpingo-oophorectomy: systematic review of literature and guidelines. *Climacteric* 2019;22:352–60.
- 59 NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines R). Ovarian cancer including fallopian tube cancer and primary peritoneal cancer. NCCN evidence blocks TM version 3.2019 – November 26, 2019. Available: https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf [Accessed 7 Jan 2020].
- 60 Rousset-Jablonski C, Selle F, Adda-Herzog E, *et al.* Fertility preservation, contraception and menopause hormone therapy in women treated for rare ovarian tumours: guidelines from the French national network dedicated to rare gynaecological cancers. *Eur J Cancer* 2019;116:35–44.
- 61 Bodner K, Laubichler P, Kimberger O, *et al.* Oestrogen and progesterone receptor expression in patients with adenocarcinoma of the uterine cervix and correlation with various clinicopathological parameters. *Anticancer Res* 2010;30:1341–5.
- 62 Martin JD, Hähnel R, McCartney AJ, *et al.* The influence of estrogen and progesterone receptors on survival in patients with carcinoma of the uterine cervix. *Gynecol Oncol* 1986;23:329–35.
- 63 Kuhle CL, Kapoor E, Sood R, *et al.* Menopausal hormone therapy in cancer survivors: a narrative review of the literature. *Maturitas* 2016;92:86–96.
- 64 Rauh LA, Pannone AF, Cantrell LA. Hormone replacement therapy after treatment for cervical cancer: are we adhering to standard of care? *Gynecol Oncol* 2017;147:597–600.
- 65 Ploch E. Hormonal replacement therapy in patients after cervical cancer treatment. *Gynecol Oncol* 1987;26:169–77.
- 66 Singh P, Oehler MK. Hormone replacement after gynaecological cancer. *Maturitas* 2010;65:190–7.
- 67 Moss EL, Taneja S, Munir F, *et al.* Iatrogenic menopause after treatment for cervical cancer. *Clin Oncol* 2016;28:766–75.
- 68 Lacey JV, Brinton LA, Barnes WA, *et al.* Use of hormone replacement therapy and adenocarcinomas and squamous cell carcinomas of the uterine cervix. *Gynecol Oncol* 2000;77:149–54.
- 69 Anderson GL, Judd HL, Kaunitz AM, *et al.* Effects of estrogen plus progestin on gynecologic cancers and associated diagnostic procedures: the Women's Health Initiative randomized trial. *JAMA* 2003;290:1739–48.
- 70 Harris RM, Waring RH. Diethylstilboestrol - a long-term legacy. *Maturitas* 2012;72:108–12.