

# Covid-19 global pandemic: options for management of gynecologic cancers. The experience in surgical management of ovarian cancer in the second highest affected Italian region

Emilia-Romagna represents the second highest affected Italian region for covid infections (more than 20 000 cases). Under these circumstances, guidelines for the management of gynecologic cancer during the covid epidemic<sup>1</sup> suggest postponing/canceling surgery, especially for ovarian cancer. This implies a reduction in the quality of treatment of the cancer.<sup>2</sup> The solution to this issue could be to identify some covid-free areas far from hospitals with patients infected with the virus. To respond to this problem, our regional policy has transformed private hospitals in cancer centers of the NHS, and the Gynecological Oncology Unit of the Sant'Orsola Hospital, Regional Ovarian Cancer Center, has been transferred into one of these hospitals to guarantee surgical procedures at the lowest risk of infection. Before admission patients awaiting surgery were: (1) contacted by telephone and checked for symptoms related to the covid virus and submitted to swabs 48 hours before surgery (the result was available after 24 hours); (2) in the case of a negative swab, the interview was repeated by medical staff on the day of admission before surgery; (3) no relatives were admitted into the ward; and (4) at hospital

discharge patients were recommended to avoid contact for 1 month. Treatment was postponed/canceled only in cases of suspected covid infection. To date, this strategy has permitted us to adhere to surgical protocols for ovarian cancer.

**Anna Myriam Perrone** <sup>1</sup>, **Alessandra De Palma**,<sup>2</sup> **Pierandrea De Iaco**<sup>1</sup>

<sup>1</sup>Gynecologic Oncology Unit. Azienda Ospedaliero-Universitaria, Policlinico di Sant'Orsola, Bologna, Italy

<sup>2</sup>Forensic Medicine and Integrated Risk Management Unit. Azienda Ospedaliero-Universitaria, Policlinico di Sant'Orsola, Bologna, Italy

**Correspondence to** Dr Anna Myriam Perrone, Azienda Ospedaliero-Universitaria di Bologna Policlinico Sant'Orsola-Malpighi, Bologna, Italy; myriam.perrone@aosp.bo.it

**Contributors** All authors contributed equally.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.



**OPEN ACCESS**

**Open access** This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© IGCS and ESGO 2020. No commercial re-use. See rights and permissions. Published by BMJ.



**To cite** Perrone AM, De Palma A, De Iaco P. *Int J Gynecol Cancer* Published Online First: [please include Day Month Year]. doi:10.1136/ijgc-2020-001489

Accepted 15 April 2020  
*Int J Gynecol Cancer* 2020;0:1.  
doi:10.1136/ijgc-2020-001489

#### ORCID iD

Anna Myriam Perrone <http://orcid.org/0000-0003-3140-4772>

#### REFERENCES

- Ramirez PT, Chiva L, Eriksson AGZ, *et al.* COVID-19 global pandemic: options for management of gynecologic cancers. *Int J Gynecol Cancer* 2020;ijgc-2020-001419.
- Colombo N, Sessa C, Bois AD, *et al.* ESMO-ESGO consensus conference recommendations on ovarian cancer: pathology and molecular biology, early and advanced stages, borderline tumours and recurrent disease. *Int J Gynecol Cancer* 2019;ijgc-2019-000308.