



# Laparoscopic pelvic and lumbo-aortic lymphadenectomy and hysterectomy by total left lateral approach

Catarina Castro Vieira <sup>1</sup>, Pauline Chauvet,<sup>2,3</sup> Feras Senny,<sup>2,3</sup> Nicolas Bourdel<sup>2,3</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/ijgc-2021-002823>).

<sup>1</sup>Department of Gynecology and Obstetrics, Hospital de Braga, Braga, Portugal

<sup>2</sup>Department of Obstetrics and Gynecology, CHU Estaing Pôle Femmes et Enfants, Clermont-Ferrand, Auvergne-Rhône-Alpes, France

<sup>3</sup>Faculty of Medicine, Université Clermont Auvergne, Clermont-Ferrand, Auvergne-Rhône-Alpes, France

## Correspondence to

Dr Catarina Castro Vieira,  
Department of Gynecology and Obstetrics, Hospital de Braga,  
Braga 4710-243, Portugal;  
[catarina.cv.16@gmail.com](mailto:catarina.cv.16@gmail.com)

Accepted 31 August 2021

This video shows a complete left lateral laparoscopic surgical approach for the treatment of endometrial carcinoma in a patient with major pelvic adhesions due to previous surgeries.

In a French university tertiary care hospital, a 73-year-old patient presented with post-menopausal bleeding and was later diagnosed with a grade 3

serous endometrial carcinoma with suspected uterine serosa, cervix, and iliac and latero-aortic lymph node involvement. It was decided to perform a total laparoscopic hysterectomy, bilateral adnexectomy, and pelvic and lumbo-aortic lymphadenectomy.

Previous abdominal interventions, including colon resection because of a volvulus and the placement of

INTERNATIONAL JOURNAL OF  
**GYNECOLOGICAL CANCER**

## LAPAROSCOPIC PELVIC AND LUMBO-AORTIC LYMPHADENECTOMY AND HYSTERECTOMY BY TOTAL LEFT LATERAL APPROACH

Catarina Vieira<sup>1</sup>, Pauline Chauvet<sup>2,3</sup>, Feras Senny<sup>2,3</sup>, Nicolas Bourdel<sup>2,3</sup>

1 - Hospital de Braga, Rua Sete Fontes, 4710-243 São Victor, Braga, Portugal

2 - Department of Obstetrics and Gynecology, CHU Estaing, 1 rue Lucie et Raymond Aubrac, 63003 Clermont Ferrand, France.

3 - Université Clermont Auvergne, Faculty of Medicine, Place Henri Dunant, 63001 Clermont Ferrand, France.



© IGCS and ESGO 2021. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Vieira CC, Chauvet P, Senny F, *et al.* *Int J Gynecol Cancer* Published Online First: [please include Day Month Year]. doi:10.1136/ijgc-2021-002823



Copyright © 2021 BMJ Publishing Group Ltd, International Gynecologic Cancer Society, & European Society of Gynecological Oncology. All rights reserved.

## Video 1



**Figure 1** Image showing the placement of all four trocars on the left side of the abdomen.

two abdominal wall meshes for a midline hernia and a left Spiegel hernia, considerably augmented the risk of adhesions. For that reason, a retroperitoneal approach was considered.

A total left lateral laparoscopic approach, with placement of all four trocars on the left side of the abdomen (Figure 1), and total hysterectomy with bilateral adnexectomy, bilateral pelvic and lumbo-aortic lymphadenectomy until the level of the left renal artery were carried out, always maintaining the left retroperitoneal approach (Online Supplemental File 1).

Laparoscopy is possible for the treatment of endometrial carcinoma, even in the presence of major pelvic adhesions due to multiple previous abdominal surgeries.<sup>1</sup> A retroperitoneal approach, classically used for lumbo-artic lymph node

dissection, can also enable pelvic lymph node dissection and total hysterectomy.<sup>2</sup>

**Contributors** NB was responsible for the recruitment of the patient and the conception and design of the study. CCV was responsible for data collection. All authors contributed to the data analysis and interpretation and manuscript preparation. Final approval of the study was made by NB.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

#### ORCID iD

Catarina Castro Vieira <http://orcid.org/0000-0001-7068-9577>

#### REFERENCES

- 1 Fowler JM, Backes FJ. Pelvic and paraaortic lymphadenectomy in gynecologic cancers, 2020. Available: [https://www.uptodate.com/contents/pelvic-and-paraaortic-lymphadenectomy-in-gynecologic-cancers?sectionName=Extraperitoneal%20laparoscopic%20access&search=laparoscopic%20extraperitoneal%20endometrial%20cancer&topicRef=3251&anchor=H352816808&source=see\\_link#H352816808](https://www.uptodate.com/contents/pelvic-and-paraaortic-lymphadenectomy-in-gynecologic-cancers?sectionName=Extraperitoneal%20laparoscopic%20access&search=laparoscopic%20extraperitoneal%20endometrial%20cancer&topicRef=3251&anchor=H352816808&source=see_link#H352816808)
- 2 Gałczyński K, Chauvet P, Canis M, *et al*. Laparoscopic extraperitoneal lumboaortic lymphadenectomy in 10 steps – Let's make it easier! *Gynecol Oncol* 2016;143:448–9.

## Material

Urinary catheter
One 12mm trocar + three 5mm trocars
0° camera
Graspers
Bipolar
Aspirator
Scissors (monopolar energy)
Endobag
Clermont-Ferrand Uterine manipulator
PDS 0 suture
Staples